One assumes that the women were told of possible metabolic side effects from continuous intake of synthetic hormones, including amenorrhoea after stopping the regimen—as also can occur with the 12-week contraceptive injection.

In view of the increasing incidence of post-pill amenorrhoea following the monthly pill-taking regimen would it not have been more ethical in this experiment to have excluded women of parity O?

ELIZABETH ELLIOTT

Wisbech, Cambs

* * We sent a copy of this letter to Dr Loudon and her colleagues, whose reply is printed below.—Ed, *BMJ*.

SIR,—In reply to Dr Elliott, the women in the study were given the same information about oral contraceptives as women starting on the standard 21/7-day oral contraceptive regimen.

We are alert to the possibility of post-pill amenorrhoea in all women who stop oral contraceptives. We are in the process of following up the subsequent menstrual pattern of women who stopped the tri-cycle regimen irrespective of their age and parity, but as yet the data are incomplete. Details of subsequent pregnancies in those women who stopped the tri-cycle regimen in order to become pregnant were included in the paper.

NANCY B LOUDON

Family Planning Services,
Lothian Health Board,
Edinburgh

R V SHORT

Medical Research Council,
Unit of Reproductive Biology,
Edinburgh

Painless thyroiditis

SIR,—Before a syndrome becomes a disease the natural history of the illness needs to be known. Painless thyroiditis (leading article, 6 August, p 348) is probably often clinically silent, but its course and duration are not usually known. It sometimes starts as subacute thyroiditis (SAT), when scattered granulomas form in the thyroid and parenchymal cells are focally damaged. This syndrome is rarely if ever due to a viral infection. Transient hyperthyroidism may develop at first, but later biopsies have shown lymphocytic thyroiditis, with variable amounts of fibrosis. Out of 108 patients reported by Woolmer *et al* 16% subsequently became myxoedematous. One patient reported by Copperman,² however, still had granulomas in her thyroid when she died of myxoedema coma.

SAT apparently quite often progresses into Hashimoto's thyroiditis and it is now usually considered to be an immune disturbance. A reason for the immune reaction is evident, in some cases. I found concurrent hyperthyroidism in more than 20 cases of sarcoidosis,³ and the serum in three of them contained long-acting thyroid stimulator (LATS) or LATS protector.⁴ Some of these patients had giant-cell granulomas in their thyroids,⁵ and the same lesion was also reported in a patient with sarcoidosis and SAT.⁶ Moreover, Hashimoto's thyroiditis and myxoedema are sometimes found,⁶ and the cellular⁸ and humoral⁹ immunity is raised in this disease.

Sarcoidosis appears to induce SAT and chronic thyroiditis immunologically whenever

the granulomas invade the gland. Although 10% of the population have subclinical thyroiditis,⁸ it is most unlikely that the granulomas in these patients had reactivated a pre-existing inflammatory process. If thyroid biopsies and Kveim tests were performed earlier, and more often, more cases of sarcoid thyroiditis would be diagnosed.

GERALD A MACGREGOR

St Luke's Hospital,
Guildford, Surrey

¹ Woolner, L B, McConehey, W M, and Beahrs, O H, *Journal of Endocrinology and Metabolism*, 1957, **17**, 1202.

² Copperman, I J, *Journal of the Irish Medical Association*, 1968, **61**, 244.

³ MacGregor, G A, *Proceedings of the Royal Society of Medicine*, 1974, **67**, 221.

⁴ Hancock, B W, and Millard, L G, *British Journal of Diseases of the Chest*, 1976, **70**, 129.

⁵ Karlish, A J, and MacGregor, G A, *Lancet*, 1970, **2**, 330.

⁶ Birchall, G, *British Journal of Clinical Practice*, 1966, **20**, 586.

⁷ Caspary, E A, and Field, E J, *British Medical Journal*, 1971, **2**, 143.

⁸ James, G D, Neville, E, and Walker, A, *American Journal of Medicine*, 1975, **59**, 388.

⁹ Rosser, R, *British Journal of Psychiatry*, 1976, **128**, 61.

The perilous skateboard

SIR,—It is highly regrettable that the BBC television programme *Nationwide* should encourage and publicise the dangerous sport of skateboarding with a national skateboard competition.

The evidence of serious injury in those attempting this hobby is at present impression only, but I predict that it will soon amass and label the skateboard as highly dangerous. The skateboard can achieve speeds of up to 40 mph (64 kph) and the skill and fun in its use lies in its inherent instability. Collision with passers-by in public places has already resulted in its banishment from several town centres, but the real danger lies in fractures and soft-tissue injuries to the participants with which the medical profession will have to cope.

It was not coincidence that the *Nationwide* personality detailed to learn the sport and have his progress monitored appeared on television after his first attempt with a torn ligament of his ankle joint. Furthermore, it was not wise to recommend the use of helmets and then to show film of expert skateboarders performing without, though the flimsy helmets are probably insufficient protection and the knee and arm pads recommended will prevent skin abrasion but not bony injury.

The skateboard is dangerous and its use should be restricted to supervised rinks where training can be given.

ADRIAN ROGERS

Stratford-upon-Avon,
Warwick

Late infection after total hip replacement

SIR,—Your leading article on this subject (23 July, p 213) and the several responses to it lead us to emphasise the need for careful bacteriological studies if meaningful conclusions are to be drawn. It is common practice for surgeons to deal with infected cases last on the list and thus material for examination is often not available until after normal laboratory hours and possibly left until the following day. As a result oxygen-sensitive anaerobes may be lost or chance contaminants multiply to a level at which they

are erroneously believed to be causative of the infection.

In an effort to answer some of the unresolved questions we have removed the bacteriological examination of prosthetic materials from the rest of the routine work and arranged that these specimens are examined as soon as possible after removal. We have also arranged for the examination to include a viable count, under aerobic and anaerobic conditions, of the numbers of organisms present on the prosthesis. As a result we are now able to exclude with some certainty the chance contaminating organism, and anaerobic species have been isolated in sufficient numbers for us to believe they are associated with the breakdown of the prosthesis.

In parallel with this study we are storing sera from patients undergoing major joint replacement. Samples are collected preoperatively and at subsequent outpatient follow-up. This enables us to examine the sera for antibodies by established methods and to develop new techniques for revealing antibodies to organisms isolated by our more searching methods. When late infection is suspected the presence of antibodies in the patient's serum gives a good indication of the time of onset of infection and provides further evidence that the organisms concerned have been involved in the failure of the prosthesis.

W J FINCHAM
JOSEPHINE COOK

Department of Clinical Pathology,
Institute of Orthopaedics,
Royal National Orthopaedic Hospital,
Stanmore, Middx

Celiac disease and malignancy

SIR,—It was with great interest that I read the case report of coeliac disease complicated by reticulum cell sarcoma by Drs A P Brooks and Christine I Harrington (17 September, p 739).

We have had a similar case recently, in which, however, there was clinical and histological response to gluten withdrawal. In fact, in our case the morphology of the jejunal mucosa improved *pari passu* with the development and manifestation of the malignancy. At post-mortem examination there was reticulum cell sarcoma in the mesenteric lymph nodes but no involvement of the bowel wall.

It is well recognised that there is an increased incidence of lymphoma and carcinoma of the gastrointestinal tract in patients with coeliac disease,¹ but, although it is generally thought that gluten withdrawal offers some degree of protection from development of malignancy,² this is by no means certain, particularly where lymphomas are concerned.³ Nor does the degree of improvement of jejunal morphology affect the risk of malignancy.

I would disagree with the authors' view that their case is unusual because the lymphoma involved the nodes and not the bowel wall. This seems to be the more usual pattern of lymphomas complicating coeliac disease, whether treated by gluten withdrawal or not, and whether or not the patient shows response to withdrawal. It is the so-called Mediterranean lymphoma that more frequently involves the bowel wall, and in a high proportion of these cases histological diagnosis can be made by jejunal biopsy.⁴ With lymphoma associated with coeliac disease, however, this is rarely the case and recourse to laparotomy is usually necessary.⁵