A prospective study was started to try to find patients who had a virus infection in the early days of pregnancy and to relate this to the occurrence of either moles or fetal abnormalities, but it is very difficult to obtain an accurate history of viral infections early in pregnancy. While there have been no severe influenza outbreaks since the "Asian flu" and there has been no recurrence of the marked increase in the incidence of hydatidiform moles, my impression is that viral infection early in pregnancy is associated with an increased incidence of fetal abnormalities.

I would be interested to know whether others have had any similar experiences, as the incidence of these conditions in any one area or practice is comparatively uncommon.

S J BARR

Fazakerley Hospital,
Liverpool

General practitioner prescribing costs

SIR,—No replies have been published to my question whether general practitioner trainees anywhere in Britain receive instructions in the economics of prescribing (30 July, p 319). This must surely stand as a serious indictment of the system by which they are trained. Advisers in general practice training, the trainers themselves, and hospital staff associated with trainees have an obligation to teach good, thrifty prescribing habits. To help in this I would suggest two practical aids.

GPs at present receive information regularly on the cost of their own prescriptions together with the local and national averages. Could this not be extended to include hospital doctors? Is it not also time that those of us working in the NHS accepted, indeed promoted, the advantages of an NHS drug formulary, probably based on the British National Formulary? Preparations not included in this could remain available but would not attract a dispensing subsidy.

Other countries, such as Australia, with public health care systems less well developed than the UK's already have this in use. Why not here?

ROBERT SCOTT

Craig Dunain Hospital,
Inverness

Medical manpower

SIR,—Professor James Parkhouse could not have expected to write and publish his very excellent paper on a simple model for medical manpower studies (20 August, p 530) without exciting a certain amount of comment. In fact he has produced a very excellent model which should enable many people to clarify their thinking on this very complex subject. If I may quote Dr Alan Maynard, "The methodology of medical manpower planning in the past has been crude to say the least."

There are some obvious comments to make on the figures produced in Professor Parkhouse's document. It is very doubtful if the losses by emigration, which he does not in fact quantify in total (although one must assume that the 2500 overseas doctors coupled with the 600 United Kingdom doctors leaving the junior hospital doctors' box in 1975 in Professor Parkhouse's diagram formed a substantial part of the total doctor emigration),

will be sustained. A further comment which requires to be made on his diagram is that he correctly states the United Kingdom graduates figure for 1975 but ignores the United Kingdom non-graduate basic medical qualifications and the graduates of the Irish universities, who enjoy the same privileges as graduates of United Kingdom universities, who traditionally have looked to the United Kingdom as an area in which it is open for them to settle, and whose numbers exceed the demand of their home market by a factor of two. This would raise the potential number of doctors entering the junior hospital doctor box by around a further thousand. (Even with access to the most accurate figures available it is impossible to state a definite figure.) If one accepts this, then the conclusions which Professor Parkhouse makes become invalid.

PAUL R J VICKERS

Gosforth,
Newcastle upon Tyne

¹ Maynard, A H. Personal communication.

Dispute in Malta

SIR,—Doctors in Britain will have read with considerable concern of the events which have taken place as a result of the dispute between the medical profession and the government in Malta. Maltese doctors have been "locked out" of their hospitals and ejected from government residential accommodation. Distinguished members of the medical profession, some of whom are external examiners in Britain, have been "sacked" for refusing to sign unacceptable undertakings required by the government. Maltese students have been unable to take their final examinations, and between 30 and 40 of them will have arrived by the time this letter is published in order to take the Conjoint examination in this country.

The BMA has taken vigorous action in support of our Maltese colleagues and has arranged assistance for students arriving here. Meanwhile there is a pressing need for financial help both for students coming to Britain and for the doctors who are victims of the government reprisals in Malta. I appeal for help for the students and for our BMA Branch in Malta, and I should be very grateful if any contributions could be sent to me, indicating whether the donation is to be used to assist the students, our colleagues in Malta, or both.

JAMES CAMERON
Chairman of Council, BMA

BMA House,
Tavistock Square,
London WC1H 9JP

Points from Letters

Shortage of anaesthetists

Dr D EYRE-WALKER (Staffordshire General Infirmary, Stafford) writes: Dr D L Freedman's letter (13 August, p 456) regarding shortage of anaesthetists and his suggested method of overcoming this problem by using local techniques does indeed suggest a way out of the difficulty but he has overlooked one important factor. These techniques take time and, from my experience in our hospital at any rate, the turnover in cases would be much reduced and our waiting lists would grow just

the same. . . . I would query also Mr Freedman's remark that "regional intravenous anaesthesia gives the operator about 20 min . . ." If carried out with two tourniquets this technique should last as long as the surgeon is prepared to leave the tourniquet on an upper limb.

Ear-piercing problems

Dr A D REDMOND (Sale, Manchester) writes: . . . Although I agree there are problems encountered with ear-piercing (Dr Ann L Jay, 27 August, p 574), they are never very serious or permanent, and to exhort the profession to discourage this practice is, I feel, somewhat over-reacting. Surely the rational approach is to encourage the correct procedure to be adhered to when ears are pierced. If carried out correctly it is relatively pain-free and free from complications.

Cough suppressants

Mr M J GILKES (Sussex Eye Hospital, Brighton) writes: Some little while ago I was appalled by correspondence in the *BMJ* in which cough-suppressant therapeutics were cavalierly dismissed by apparently caring and clinically experienced correspondents as being without effect. The overall impression was given that such preparations should be abolished from the therapeutic armamentarium, at least on the NHS. . . . Having just experienced a short and sharp but none the less very acute upper respiratory condition, which appears to be mildly epidemic and of which the major feature was convulsive, almost pertussis-like repetitive and irritative cough, which at the end of 12 hours had brought the abdominal muscles to a state of acute tenderness, I can only say, "Thank God for the presence of an effective cough suppressant, such as linctus actifed co and linctus scillae co," without which even a partial night's rest would have been demonstrably unavailable. . . .

Latin as she is wrote

Dr GILBERT WALKER (Dundee) writes: I know it is unfashionable to consider spelling to be important, but is it asking too much to have our Latin tags correctly set out? Proof-reading should have eliminated the nonsense of "res ipse loquitur" (27 August, p 569). . . . Unless my book is out of date "res" is feminine and "loquitur" is a verb. Res ipsa loquitur.

Treatment of hiccup

Mr S GOLDWATER (London NW2) writes: However an attack of hiccup commences it would seem that a vicious circle is set up consisting of hiccup producing air-swallowing, producing a further hiccup, and so on. When the attack has been going for some time there is usually a marked increase in the area of gastric resonance and treatment should surely be aimed at reducing the amount of air in the stomach. On this hypothesis I have treated many cases with charcoal tablets and have instructed patients to continue chewing them at least once an hour and in extreme cases continuously. In most cases I have met with success on this simple regimen. . . .