might be there accidentally; even if they were found constantly they might be "merely parasitic even though constant elements of contagions, as some still say of the semen-animalcules, elements that develop in the fluid, and may even be of diagnostic significance, without being the active material of the fluid or of the semen." Henle then said that only by isolating semen-animalcules (*Samenthierchen*) and seminal fluid, contagion-organisms (*Contagiumorganismen*) and contagion fluid, could the power of each be separately observed. Such an experiment, added Henle, "must no doubt be renounced."²

As Koch himself recognised when he failed to produce experimental cholera infections in animals in 1883-4, the postulates attributed to him were an oversimplification, but to compare Henle's armchair speculations with the hard-won conclusions from Koch's careful experimental work is surely absurd. It is surprising that the scholarly William Bulloch should have adopted the Fildes-McIntosh myth.³ This question has been dealt with in more detail in a recent paper.⁴

NORMAN HOWARD-JONES

Geneva, Switzerland

¹ Fildes, P. and McIntosh, J. *British Journal of Experimental Pathology*, 1920, 1, 119 and 159.

² Henle, J. *Pathologische Untersuchungen*, p 43. Berlin, Hirschwald, 1840.

³ Bulloch, W. *History of Bacteriology*, p 165. London, Oxford University Press, 1938.

⁴ Howard-Jones, N. *Medical History*, 1977, 21, 61.

Safety and danger of piped gases

SIR,—How encouraging to find Dr J V I Young (30 July, p 317) pointing out the dangers of piped nitrous oxide! Not only is it dangerous to have two gases piped to an operating theatre, it is surely quite unnecessary. Dr Young could in fact have drawn our attention to another nine reasons why, in 1977, we should dispense with the routine use of nitrous oxide in major surgery.

(1) The risk of oxygen failure is minimised since if only one carrier gas is used the efforts of a patient to breathe on an empty bag, or a ventilator to work without a proper supply, are much more obvious than if another gas such as nitrous oxide carries on when oxygen has failed.

(2) Nitrous oxide is produced chemically and therefore there is a risk of impurities. Oxygen is produced physically and any impurities present were presumably in the air from which the oxygen was produced.

(3) All inhalational anaesthetics are more flammable in nitrous oxide than in oxygen. Even halothane will burn in nitrous oxide; it is non-flammable in pure oxygen.

(4) Without nitrous oxide there is less danger from gut distension, air embolism, and bubbles in the middle ear, and the cuffs of endotracheal tubes are less liable to over-distension.

Once the fact that good anaesthesia can be produced today without the use of nitrous oxide is accepted the way is clear for a real step forward in anaesthetic technique—the use of a totally closed system. The addition of a simple ultraviolet halothane meter is all that is required to be able to enjoy a greatly simplified anaesthetic technique, which has a further five advantages.

(5) The problem of pollution is virtually eliminated. Apart from expense, present

scavenging systems have at least four risks of their own: (a) obstruction to free movement in the theatre; (b) risk of infection; (c) risk of misconnection; and (d) risks associated with the final disposal of waste gases.

(6) With oxygen as the only carrier gas capital costs are greatly reduced. There is no need for a pipeline since nitrous oxide is not required and only about 250 ml of oxygen per minute is needed.

(7) Running costs are greatly reduced.

(8) Water vapour is saved.

(9) Heat loss is reduced.

T H S BURNS

St Thomas's Hospital,
London SE1

Prolonged remission maintenance in acute myeloid leukaemia

SIR,—We were interested to read the account by Professor A S D Spiers and others (27 August, p 544) of their results in the treatment of acute myeloid leukaemia with three quadruple drug combinations in predetermined rotation: TRAP (thioguanine, daunorubicin, cytarabine, prednisolone); COAP (cyclophosphamide, vincristine, cytarabine, prednisolone); and POMP (prednisolone, vincristine, methotrexate, mercaptopurine). They state that "for maintenance, five-day courses of drugs were administered every 14-21 days and doses were increased to tolerance" (that is, if possible a 20% dose increase of rubidomycin dosage was made on each course and a similar increase in that of some of the other drugs was also attempted at each course of treatment). Cardiotoxicity or fear of this complication made them discontinue rubidomycin when their patients had been in remission for 62-122 weeks. They treated a total of 25 patients according to this protocol between July 1970 and January 1976, and on 31 January 1977 only two of the 25 patients were still in complete remission. The remissions of these two patients had lasted for 119 and 134 weeks respectively, a relatively short time. They therefore came to the somewhat pessimistic conclusion that this protocol "must still be regarded as only palliative treatment for acute myeloid leukaemia."

Since 1971 we have used a modification of Spiers *et al* TRAP-COAP-POMP rotation protocol. One of our modifications has been to keep the dosage of rubidomycin and other drugs at the original level (for example, 60 mg of rubidomycin for the average person for each course instead of the increases attempted by Spiers *et al*). During the first year we start treatment courses every two weeks, during the second year every three weeks, and during the third year every four weeks. (Two patients also received a fourth year of treatment, with courses starting every six weeks.)

From August 1971 to December 1973 56 patients were started on this treatment. Our remission rate and median survival were similar to those of Spiers *et al*. However, with our modifications we encountered no cardiotoxicity and continued our treatment for three years (in two cases for four years). Seven of our 56 patients are still in full remission 4-6 years from the start of treatment, and 1-3 years after discontinuing all treatment. Of these seven long survivors one was 17 years old at the start of treatment, one was 57, and the other five were in their 40s. One of these was a 40-year-old farmer who had asymptomatic insufficiency of rheumatic aetiology when his acute

leukaemia was diagnosed six years ago. Three years ago he developed severe cardiac failure and a Björk-Shiley disc valve prosthesis was inserted with dramatic improvement. Today his cardiac function is good and his bone marrow and peripheral blood counts are normal three years after discontinuing all leukaemia treatment. Another of our patients was a 17-year-old schoolgirl who was started on treatment six years ago. She was treated for four years and married at the time when the treatment was discontinued. One year after the treatment was discontinued a chromosome analysis showed normal findings with no chromosomal breakage. She then allowed herself to become pregnant and gave birth to a healthy daughter in August this year.

We think that the results obtained with our modification of Spiers *et al* TRAP-COAP-POMP treatment permit us to be considerably more optimistic than to regard it as only palliative treatment.

PER STAVEM
TORBJÖRN GJEMDAL
BERNT LY

Section of Haematology,
Medical Department A,
Rikshospitalet,
Oslo, Norway

An epidural service

SIR,—In reply to Mr A F Pentecost's letter (3 September, p 644) about our epidural service at this hospital (6 August, p 370), our aim has not been to provide total relief of pain in labour but to allow midwives and patients to manage the epidural more flexibly, according to how much normal sensation the patient wishes to retain. A midwife might, for example, explain to a patient who requested a top-up close to full dilatation that she would deliver herself more readily without a top-up or with a half-volume top-up at that stage. However, if the patient wanted total sensory ablation she could have it but was more likely to require assistance with forceps.

On this basis 80% of our patients were fully satisfied, and of the remaining 20% who were not some were dissatisfied with the way their epidural was managed and a few had epidurals which were not working properly. In this latter group another epidural catheter which worked was usually inserted so that overall our records show that only 2% of the total had a failed epidural which we could not correct.

I do believe that the provision of epidural analgesia should be part of the in-service training of junior obstetric staff and that the acquisition of this skill is of greater value to overseas doctors returning home than some of the techniques we teach them.

I also had difficulty in getting patients and midwives to accept the obvious benefits of this type of pain relief in labour. Resistance melted when our midwives, even some of the district midwives, realised how much more satisfying it was to manage a patient with adequate pain relief whose consciousness was not clouded with pethidine. Patients' attitudes have also changed because in a small town where the forum of communication is the local supermarket they quickly get to hear of each other's experiences in the maternity unit.

I could not provide this service on my own with the help of a medical assistant as Mr Pentecost does, which is another reason why I have involved my junior staff.