have died as a result of skateboarding in America, but how many million skateboards are in use over there?

My impression, for what it is worth, is that skateboarding is a lot safer than rugby or boxing, and infinitely safer than travelling in a car or on a motorcycle. Skateboards are like cars: they are not dangerous, but the people who use them may be. Why must doctors always be seen to be promoting the negative rather than the positive in people's lives? We live in a culture of isolation and boredom; we should be trying to do something about that before we condemn activities out of hand without the information necessary to back it up.

S GLASCOE

Royal Gwent Hospital,
Newport, Gwent

¹ *Daily Telegraph*, 1 September 1977.

Switching off

SIR,—I write to protest against the practice of asking relatives for permission to disconnect life support systems in cases where brain death has been diagnosed. This is both illogical and cruel and in many cases may prove to be counterproductive.

It is illogical because the relatives are not involved in the decision to put the patient on life support. Nor indeed could they be. The decision to initiate life support is a medical one and by the same token the decision to disconnect is a medical one. Fortunately, we now have clear guidelines on the diagnosis of brain death, which are contained in the statement issued by the honorary secretary of the Conference of Medical Royal Colleges and their Faculties in the United Kingdom on 11 October 1976.¹

It is cruel because relatives are placed in an agonising dilemma by being asked to agree to the switching off. They do not have the medical knowledge to make a scientific judgment so they are, in effect, being asked to make a moral judgment. But in asking for a moral judgment at such a time we are placing them on the horns of a dilemma. If they say yes, they may feel they are taking a life. If they say no, they may feel they are unnecessarily prolonging death or indefinitely prolonging a useless life. Without full medical knowledge they cannot give a reasoned answer and whichever answer they give they will be left with doubts. What the relatives need is support in their distress and confidence in the judgment of their medical advisers. Our duty is to make up our minds and then quietly and sympathetically inform the relatives that life support will be withdrawn because we have determined by a battery of tests that the brain is dead.

Asking the relatives for permission to switch off is likely to be counterproductive both to the patient and to others. If permission is refused and a useless life is prolonged then an intolerable burden will be placed on the relatives. Moreover, the unnecessary postponement in switching off may result in such physical deterioration of the patient that his kidneys can no longer be used to save another life.

In conclusion, I would like to emphasise that I am not suggesting that the relatives should not be involved in the decision to withdraw life support. I am suggesting that they should not be asked to make the decision. We as a medical profession should have the

courage of our convictions and recognise that it is our responsibility to make the decision on the medical evidence.

A W FOWLER

Bridgend General Hospital,
Glamorgan

¹ *British Medical Journal*, 1976, 2, 1187.

The neonatal electrocardiogram and unexpected death in infancy

SIR,—We do not agree with the negative opinion expressed by Professor J L Emery (24 September, p 833) on the value of a major ECG screening study to investigate the possible relationship between cardiac arrhythmias and unexpected infant deaths. The population in Sheffield was selected and relatively small. Furthermore, no information was given on the type of ECG or the duration of the rhythm strips carried out.

We have demonstrated, using 24-hour monitoring, that some infants with normal screening ECGs have serious cardiac arrhythmias. We still consider a prospective multi-centre study essential to determine whether cardiac arrhythmias occur regularly in infants throughout the country, whether they are dangerous, and finally whether they can cause sudden death.

D P SOUTHALL
E A SHINEBOURNE

Brompton Hospital,
London SW3

Oxytocin and neonatal jaundice

SIR,—Several UK centres have reported a recent increase in neonatal jaundice.^{1,2} Oxytocin or some as yet unidentified obstetric practice has been held responsible.^{3,4} However, there is some doubt that a genuine increase in the incidence of neonatal jaundice has occurred since serial figures^{1,5} are few and the incidence of jaundice increases with the proportion of neonates whose plasma bilirubin is measured. We have sent questionnaires to paediatricians in Europe (E) and North America (A) asking their recent experience of jaundice rates.

Sixty-six members of the European Society for Paediatric Research were circularised; 52 replied. A random 1 in 5 sample of the American Pediatric Society, total 135, were also circularised; 100 replied. Thirty-nine of the Europeans and 51 of the North Americans were in active practice in neonatology. The majority, 66% (E) and 60% (A), thought there had been no increase in neonatal jaundice since 1970. Only 23% (E) and 24% (A) thought jaundice had increased. The remainder were uncertain. The respondents' estimate of the present percentage of neonates having a peak plasma bilirubin concentration of 206 $\mu\text{mol/l}$ (12 mg/100 ml) or greater in the first 7 days after birth varied between 2 and 30%. The averages of all the replies, 14.2% (E) and 12.6% (A), were similar to the published incidence in unselected series.⁵ We asked also about the use of phenobarbitone in late pregnancy for the prophylaxis of neonatal jaundice. Only 13% (E) and 8% (A) paediatricians used this occasionally; none used it routinely.

If oxytocin were icterogenic then we would have expected most paediatricians in North America and Europe to have noticed some

increase in neonatal jaundice. The fact that they have not casts further doubt on this association.

LOUISE FRIEDMAN
PETER LEWIS
DAVID HARVEY

Institute of Obstetrics and
Gynaecology,
London

¹ Campbell, N, *et al*, *British Medical Journal*, 1975, 2, 548.

² Sims, D G, and Neligan, G A, *British Journal of Obstetrics and Gynaecology*, 1975, 82, 863.

³ Beazley, J M, and Alderman, B, *British Journal of Obstetrics and Gynaecology*, 1975, 82, 265.

⁴ Chew, W C, and Swann, I L, *British Medical Journal*, 1977, 1, 72.

⁵ Lewis, P J, *Therapeutic Problems in Pregnancy*, p 141. Lancaster, Medical and Technical Publishing, 1977.

Sealing of wounds with vacuum drainage

SIR,—Most surgeons these days make use of vacuum drainage with zimmer bottles or similar apparatus and many surgeons no doubt have learnt the value of sealing these wounds and the emergent drainage or drains with such adhesive films as Opsite or Steri-drapes. These substances are, however, expensive and I have found that simple household Cling, so useful in the kitchen, makes a most satisfactory substitute.

In view of the lack of sterility, it is best to cover the wound and emergent drain or drains with a small strip of gauze or ribbon gauze. The whole area is then generously sprayed with Onibuctane and the film of Cling spread over. This technique is of especial help when a vacuum tube has to be replaced from without in through a new stab wound. The Cling gives rise to a new and most effective vacuum seal.

C PATRICK SAMES

Bath

Raised plasma urea concentration

SIR,—Surprise was expressed by Professor D B Morgan and others (8 October, p 929) on finding heart failure to be the commonest cause of a raised plasma urea in a general hospital. In a study which I undertook on subjects suffering from acute myocardial infarction, 50% had abnormally high plasma urea concentrations and 41% had high creatinine concentrations at some time during their stay in hospital.¹ Since such cases constitute a substantial proportion of hospital admissions, the observation made by these workers is perhaps not so surprising after all, and presumably their congestive heart failure group would include subjects of this type.

Urine estimations on a small number of patients in my investigation indicated that while glomerular filtration was sometimes decreased, as judged by creatinine clearance, urea excretion over 24 hours tended to be normal or high. Although the two studies are not strictly comparable, being designed for different purposes, some similarities do exist. For example, in both cases mortality was associated with the highest levels of plasma urea, and my results suggested that daily plasma urea measurement might be helpful in monitoring progress and prognosis in acute myocardial infarction.²

Professor Morgan and his associates list decreased glomerular filtration rate as one of three general causes of a raised plasma urea, and in heart failure this may result from poor renal perfusion. Hence the uraemia occurring in this condition seems likely to have poor circulation as its primary cause, but with a mixed secondary aetiology of tissue breakdown due to stagnant tissue