bicarbonate (or sodium hydroxide) and severe metabolic alkalosis with hydrochloric acid.² When acid or alkali is administered therapeutically in 50-100-mmol quantities it is rational to express their impact on the patient in the same way.

There are two less familiar ways of expressing acidity within the SI. The most scientifically rigorous is to express acidity in terms of the chemical potential of H ion; the SI unit for energy, the joule, is used for this purpose so that normal arterial plasma acidity becomes -44.23 to -43.76 kJ/mol.³ One can easily imagine that this unit would not be acceptable in ordinary medical use. The fourth term for acidity is the concentration of free H ions in nmol/l derived from pH in a stricter manner than indicated above. Readers seeking more information about this complex matter are referred to Professor Siggaard-Andersen's contributions cited below.^{1,3}

P J N HOWORTH

Department of Chemical Pathology,
King's College Hospital,
London SE5

¹ Siggaard-Andersen, O. *Scandinavian Journal of Clinical and Laboratory Investigation*, 1977, **37**, suppl 146, p 7.

² Worthley, L I G, *British Journal of Anaesthesia*, 1977, **49**, 811.

³ Siggaard-Andersen, O. *Blood pH, Gases, and Electrolytes*, ed R A Durst, p 1. Washington, DC, National Bureau of Standards Special Publication No 450, 1977.

Tetrabenazine in Huntington's chorea

SIR,—We have just come across a most remarkable misquotation in the paper on this subject by Drs C H Hawkes and C H Nourse (28 May, p 1391).

It is said there that we "have expressed reservations about tetrabenazine in Huntington's chorea, suggesting that the risks of depression and drug-induced Parkinsonism outweigh the advantages of its antichoreic action." We made no such point in our paper¹ but on the contrary said that depression is common in Huntington's chorea and it is difficult to be certain whether it is caused by the drug. Parkinsonism was also not noted in our series. We suggested instead that postural hypotension and dysphagia were the serious side effects which require careful supervision.

C Y HUANG
C ELLIOTT

Lidcombe Hospital,
Lidcombe,
New South Wales

¹ Huang, C Y, et al, *Medical Journal of Australia*, 1976, **1**, 583.

Vertigo in children

SIR,—I should like to mention another possible cause of vertigo in children which was omitted from your leading article (5 November, p 1173)—namely, Menière's disease. I fully appreciate that this is extremely rare in childhood, but it is important because of the potential damage to hearing.

At the age of 7 I suffered a number of severe attacks of vertigo and subsequently became deaf in one ear. By the time this was brought to medical attention there was profound hearing loss, and anyway probably nothing much could have been done then. At the age of 21 tinnitus and distortion of hearing appeared in the other ear, but vertigo

did not appear until seven years later. It was only then fully appreciated that I had bilateral Menière's disease. I was not much encouraged by your leading article on "Surgery for Menière's disease" (15 January, p 124), but did have an endolymphatic sac drainage operation in March, which has been of enormous benefit. It has restored my hearing from 30-50 dB to within normal limits and abolished vertigo.

It is for this reason that I would urge colleagues presented with vertiginous children to examine the hearing also in order not to miss a condition which may now be helped in certain cases. Also, it is all too easy to dismiss vertigo as "benign," but during my childhood attacks I narrowly escaped injury on more than one occasion. Regarding treatment of other causes of vertigo, sedatives and antiemetics can do a great deal to alleviate the distress of persistent nausea and vomiting while the condition lasts.

PATRICIA M SARTER

Nantwich, Cheshire

Vertigo and the pill

SIR,—I have read with interest your leading article on the incidence of vertigo in children (5 November, p 1173).

My experience as an otolaryngologist has led me to believe that there are many causes of vertigo, especially in young people, which have not been recognised so far. The common causes of labyrinthine vertigo such as Menière's disease or labyrinthitis usually occur in the older age groups, but in my personal practice I have met with an increasing number of young women who have suffered from true vertigo. Examination of their respiratory tract has not shown evidence of disease in the ears, nor has there been very much in the way of deafness or disorder of labyrinthine function as shown on an electronystagmograph.

There has been one common thread in all these patients and that is that most of them are young women in their 20s and 30s taking a contraceptive pill. They have often related the onset of the symptoms to the time when they began taking the pill. In many cases I have persuaded the patient to discontinue the pill and perhaps adopt an alternative method of contraception and the incidence of vertigo has subsided. An attempt was made to gain the co-operation of a family planning clinic to send patients along who are on the pill with a view to testing their hearing as well as their labyrinthine function, but unfortunately most of them were unwilling to submit to such an investigation and it had to be abandoned.

J SIEGLER

Liverpool

A dangerous kitchen tool

SIR,—We wish to draw attention to a possible danger from the use of the currently popular and reputable types of freezer knife. These are used to cut frozen foods and to chip away ice, when the knife is held as a dagger. As can be seen from the illustration, this exposes the hand to injury if the grip slips down on to the blade. A housewife aged 53 attended this department with a laceration of the little finger of the right hand sustained in this way which severed both flexor tendons and one digital nerve.

This would appear to be an unacceptable risk and the knife should be either withdrawn or modified to embody a guard.

ANDREW K MARSDEN
MICHAEL W FLOWERS

Accident and Emergency Department,
General Infirmary,
Leeds

Anti-e and vertical transmission of hepatitis B surface antigen

SIR,—Okada and his colleagues¹ provided evidence that hepatitis Be antigen (HBeAg) may be used to predict transmission, and anti-HBe the absence of transmission, of hepatitis B virus from asymptomatic carrier mothers to children. The same group of workers² found that HBsAg-specific DNA polymerase activity was invariably present in serum samples of asymptomatic carriers containing HBeAg but was not detectable in serum samples containing anti-HBe. The lack of infectivity of anti-HBe-containing HBsAg-positive serum has, however, recently been questioned. Berquist and his colleagues³ were able to produce experimental hepatitis B in a chimpanzee by inoculating it with anti-HBe-containing HBsAg-positive human serum.

In Glasgow during the last 18 months we have followed up eight asymptomatic carrier mothers and their eight children over a period ranging from three to 13 months.

Of these, two babies, both male, developed HBs-antigenaemia at the age of 3 months. One baby born to a mother whose serum contained HBeAg had no clinical or biochemical evidence of hepatitis and later became a persistent HBsAg carrier. The other baby born to a mother whose serum contained anti-HBe at the time of delivery subsequently developed acute icteric hepatitis with peak serum bilirubin at 116 μ mol/l, peak aspartic (AST) and alanine (ALT) aminotransferases at 3945 and 1830 U/l respectively. He made an uneventful clinical and biochemical recovery and cleared the circulating antigen 62 days after its initial appearance. Three months later antibody to HBsAg (anti-HBs) was demonstrated in his serum. The most probable source of this infection would appear to be the mother (as the baby was subtype ay, mother subtype ay, and father negative for both antigen and antibody). Gerety⁴ also reported a similar case of hepatitis B in an infant born to an asymptomatic carrier mother whose serum contained anti-HBe. Our evidence and that of Gerety indicate that carriers of hepatitis B surface antigen who are positive for anti-HBe cannot be regarded as totally non-infectious, at least in the vertical transmission situation, although the probability of an anti-HBe-positive mother transmitting the infection to her offspring may be