susceptible to obstruction as a result of lymphoid hyperplasia until involution of this lymphoid tissue occurs. This relationship would account for the characteristic age distribution of acute appendicitis.⁵ On this basis lymphoid hyperplasia occurring during viral exanthemata and bacterial infections could lead to appendicitis. However, one study has shown that there is no greater incidence of acute appendicitis in children with these infections than there is in children who are otherwise well.⁶ An alternative explanation is that obstruction is due to muscular spasm in the appendix associated with a low-fibre diet, as is thought to occur in the colon, and this would account for the geographical distribution of the disease.^{3, 4}

L W L HORTON

Department of Surgical Pathology,
St Thomas's Hospital,
London SE1

¹ Wagensteen, O H, and Bowers, W F, *Archives of Surgery*, 1939, **34**, 496.

² Bowers, W F, *Archives of Surgery*, 1939, **39**, 362.

³ Burkitt, D P, *British Journal of Surgery*, 1971, **58**, 695.

⁴ Burkitt, D P, in *Refined Carbohydrate Foods and Disease*, ed D P Burkitt and H C Trowell, p 87. New York, Academic Press, 1975.

⁵ Bohrod, M G, *American Journal of Clinical Pathology*, 1946, **16**, 752.

⁶ Lawrence, K B, and Waring, G W, *New England Journal of Medicine*, 1949, **241**, 1.

Richter's hernia at site of insertion of laparoscope

SIR,—Mr J B Bourke's suggestion (26 November, p 1393) that this complication might be eliminated if smaller-diameter laparoscopes were used cannot be allowed to pass without comment. It is surprising that a surgeon (rather than a gynaecologist) should ever accept an unsutured 12 mm defect in the linea alba. This defect is easily closed with a figure-of-eight suture using a heavy taper-cut or cutting needle and the problem is eliminated. Smaller defects caused by smaller laparoscopes should be similarly sutured.

At present the general surgeon is usually called to deal with the complications of gynaecological laparoscopy. Natural reluctance to embarrass colleagues results in little comment being made in these circumstances and the reported complications are almost certainly only the "tip of the iceberg." As laparoscopy is used increasingly in general surgery and the surgeon treats complications of his own making I anticipate further reports of Richter's hernia if the defect in the linea alba is not sutured.

DOUGLAS STEWART

Royal Infirmary,
Edinburgh

Labetalol and urinary catecholamines

SIR,—We have recently learnt that the levels of free urinary catecholamines are considerably increased in some patients following oral labetalol and that this effect may persist for up to 10 days after stopping the drug. The demonstration of excess urinary catecholamines is often used as a criterion for the diagnosis of pheochromocytoma, but patients suspected of having this condition are not normally investigated until all antihypertensive treatment has been stopped. Nevertheless, we have recently heard of a labetalol-treated patient in France in whom pheochromocytoma was wrongly diagnosed and an unnecessary operation subsequently carried

out. It is to avoid similar mistakes that we have written this letter.

We are carrying out experiments to determine the mechanisms whereby labetalol increases urinary catecholamines. Increased sympathetic activity in response to the fall in blood pressure caused by labetalol is, of course, a probable explanation for the excess urinary catecholamines. This hypothesis and others are now being investigated in our laboratories and the results obtained will be published as soon as possible. In the meantime we wish to reassure physicians that rebound hypertension did not occur in patients in whom labetalol treatment was deliberately stopped abruptly and has not been reported in any other patient.

DAVID HARRIS
Medical Director,
D A RICHARDS

Head of Medical Department,
Allen and Hanburys Research Ltd

Ware, Herts

Shadow over Maltese medicine

SIR,—Mr J Z Adami (17 December, p 1604) says that he can provide proof that the aid offered by the BMA to Maltese students goes only to students who take an anti-Maltese Government stand and he therefore accuses the BMA of being a political organisation. The fact is that help is offered, in so far as it is possible to offer help, to all Maltese students irrespective of their political allegiances, which are of no concern to us. I have, however, made inquiries as a result of which I am informed that two students who support the Maltese Government are in this country. The first is Mr Adami himself, and he has not so far called at BMA House, nor has he requested assistance. The other student called at BMA House to request our help in obtaining a locum. He was given several addresses and I understand that he has now obtained a preregistration job in the London area, though we have not heard from him since. I believe that there are two Maltese students apart from Mr Adami who have not so far visited us at BMA House, but we are not aware of their political views. Needless to say, we shall be very pleased to see them if they would care to call.

One student who, we are told, supports the Maltese Government has sought financial help from us. Unfortunately we were unable to help him, as the fund from which we provide financial assistance is limited to Maltese nationals and this student is not Maltese. It had nothing to do with his political views, of which we were unaware at the time.

E GREY-TURNER
Secretary, BMA

BMA House,
London WC1

Prevalence of communicable diseases

SIR,—In your recent leading article on schistosomiasis in China (19 November, p 1304), the unqualified statement that "Schistosomiasis is the only communicable disease in the world whose prevalence is increasing" begs several questions. How far is it justifiable to consider the prevalence of a communicable disease as global when most such diseases occur in their own separate environments? Furthermore, those who have intimate knowledge of the occurrence of communicable diseases in

the field know the difficulties in interpreting official notifications.

Are we not in our seventh pandemic of cholera? Are viral jaundices on the decline? Is malaria control equally effective in all parts of the tropics? Is not the "epidemic" of Bancroftian filariasis in the Far East still under way? Although transmission of onchocerciasis (associated with river blindness) may have been controlled in the Haute Volta, what is happening in the rest of Africa? And is the control of sleeping sickness effective in all parts of Africa?

In sum, sir, the increase in the prevalence of communicable diseases may by no means be confined to schistosomiasis.

WILLIAM KERSHAW

Hesketh Bank,
Preston, Lancs

Emergency in emergency departments

SIR,—I would like to endorse wholeheartedly Mr C C Slack's anxiety regarding the future of staffing of accident and emergency departments throughout the country (19 November, p 1359). In his letter he raises one or two important points regarding the staffing of such departments.

(1) He points out that the present junior hospital doctors' contract has made the work in an accident and emergency department less attractive than before. I am sure that he does not state this point forcefully enough, as most departments operate a 40-hour week, creating a situation in that doctors moving to an accident department job from a surgical specialty will often suffer a cut in salary of up to 30%.

(2) He raises the point of subsequent default and alleges that some hospitals adopt a policy that work in the accident and emergency department must be the first undertaken in any surgical rotation. This is not our practice in Hull, but it is certainly accepted both by the senior house officers and by the employing consultants that one reason for senior house officers wishing to work in the department is that they use it as a time for taking examinations, as a consequence of which up to a month of the six months spent in the department may be taken as holiday, study leave, and leave for examinations.

(3) Many doctors entering Britain now have little option but to take jobs in accident and emergency departments, as few surgical specialties will accept senior house officers unless they have already passed their primary fellowship examination. This is a blueprint for disaster, as accident and emergency departments are fraught with dangers of diagnosis, medicolegal problems, hazards with communication, etc, and it is of course the last place that a doctor new to the country should elect to work.

Having now had experience of managing an accident and emergency department for several years I would like to propose (1) that it is possible to run both a training programme and a career grade structure of doctors parallel to one another in the same department; and (2) that the junior staffing levels of a department should be based on the number of patients seen per annum in that department. It is accepted now that the specialty of accident and emergency work has come to stay and it therefore follows that all major accident