Experimental demonstration of the antiherbivore effects of silica in grasses: impacts on foliage digestibility and vole growth rates

Fergus P. Massey* and Sue E. Hartley

Department of Biology and Environmental Science, University of Sussex, Falmer, Brighton,
East Sussex, BN1 9QG, UK

The impact of plant-based factors on the population dynamics of mammalian herbivores has been the subject of much debate in ecology, but the role of antiherbivore defences in grasses has received relatively little attention. Silica has been proposed as the primary defence in grasses and is thought to lead to increased abrasiveness of foliage so deterring feeding, as well as reducing foliage digestibility and herbivore performance. However, at present there is little direct experimental evidence to support these ideas. In this study, we tested the effects of manipulating silica levels on the abrasiveness of grasses and on the feeding preference and growth performance of field voles, specialist grass-feeding herbivores. Elevated silica levels did increase the abrasiveness of grasses and deterred feeding by voles. We also demonstrated, for the first time, that silica reduced the growth rates of both juvenile and mature female voles by reducing the nitrogen they could absorb from the foliage. Furthermore, we found that vole feeding leads to increased levels of silica in leaves, suggesting a dynamic feedback between grasses and their herbivores. We propose that silica induction due to vole grazing reduces vole performance and hence could contribute to cyclic dynamics in vole populations.

Keywords: plant defence; population cycles; *Microtus agrestis*; feeding preference; herbivore performance; induction

1. INTRODUCTION

Grasses dominate many terrestrial ecosystems and are of great ecological and economic importance, as are the grazing mammals associated with them. Although coevolution between grasses and grazers has received a great deal of attention (Stebbins 1981; McNaughton 1984; Jernvall & Fortelius 2002; Prasad *et al.* 2005), the ecological mechanisms by which grasses defend themselves against herbivores remain poorly understood (O'Reagain & Mentis 1989; Vicari & Bazely 1993). The silica content of grasses can be considerably higher than in many other plants and has been proposed as an antiherbivore defence strategy (McNaughton *et al.* 1985; Gali-Muhtasib *et al.* 1992; Vicari & Bazely 1993). Silica is deposited as phytoliths in leaves. These are harder than tooth enamel (Baker *et al.* 1959), so they could potentially increase the abrasiveness of leaves and deter feeding (Vicari & Bazely 1993). However, currently the evidence supporting the effects of silica on the palatability of grasses to vertebrate herbivores is limited: there have been relatively few experimental studies and none which manipulate silica content and measure the subsequent changes in abrasiveness of leaves. For example, Gali-Muhtasib *et al.* (1992) found prairie voles (*Microtus ochrogaster*) preferred leaves of grasses with lower silica content when comparing cut leaves of two *Bromus* species, but the effect of silica on abrasiveness was not tested. It has also been proposed that phytoliths may disrupt microbial action in the gut of herbivores, reducing the digestibility of grasses (Vicari & Bazely 1993), but almost nothing is known

about the effects of silica on either grass leaf digestibility or herbivore performance.

Many small mammalian herbivores show strong selectivity in feeding preference both within and between plant species. In voles, these preferences have previously been shown to relate to both the nutrient content (e.g. nitrogen) of their food plants and the levels of physical and chemical antiherbivore defences (Batzli & Cole 1979; Marquis & Batzli 1989; Hartley *et al.* 1995; Hjalten *et al.* 1996). Despite this evidence of the importance of food quality for vole behaviour, the effects of silica, the principal defence in grasses, have not been fully investigated.

For small, fast growing and short lived mammalian herbivores such as voles, food quality is particularly significant because of their relatively high metabolic rates (Demment & Van Soest 1985), and because they have a limited capacity to increase rates of food consumption to compensate for poor quality diets (Zynel & Wunder 2002). In addition, the growth rates early in development are highly dependant upon nutrient intake. Therefore in voles, food quality has the potential to dictate the time taken to reach sexual maturity and the onset of breeding each year (Krebs & Myers 1974; Ergon *et al.* 2001). There has been considerable interest in the factors influencing the growth rates and reproductive output in voles because of the impact these parameters have on individual fitness and potentially on vole population dynamics (Agrell *et al.* 1995; Klemola *et al.* 2000a; Turchin & Batzli 2001; Ergon *et al.* 2001, 2004). Changes in food quality, if they have significant effects on vole growth and reproduction, may play a role in these population processes.

* Author for correspondence (f.p.massey@sussex.ac.uk).

Table 1. (i) The chemical composition of grass leaves under high and low silica treatments (% dry mass; mean values $\pm$ s.e.m.) and (ii) the leaf silica content of study species from field sites (minimum–maximum values in plants collected from 10 different natural plant communities). (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$)

(i)	silica treatment	leaf silica content	leaf nitrogen content	total phenolic content	(ii) field silica levels
<i>Festuca ovina</i>	low	0.52 $\pm$ 0.04	2.18 $\pm$ 0.18	1.27 $\pm$ 0.05	0.65–3.54
	high	2.44 $\pm$ 0.17	2.18 $\pm$ 0.18	1.25 $\pm$ 0.11	
<i>Lolium perenne</i>	low	0.54 $\pm$ 0.10	3.10 $\pm$ 0.10	1.39 $\pm$ 0.05	0.58–4.52
	high	4.68 $\pm$ 0.34	3.12 $\pm$ 0.10	1.38 $\pm$ 0.05	
ANOVA ^a	species	***	***	n.s.	
	silica	***	n.s.	n.s.	
	species $\times$ silica	***	n.s.	n.s.	

^a n.s. = not significant

Here, for the first time, we test both the proposed mechanisms by which silica acts as a defence, namely its ability to increase abrasion and its effect on foliage digestibility and herbivore performance. We measure the effects of silica on the feeding preference and growth rates of field voles (*Microtus agrestis* L.). We focus on the effects of silica on the growth of female field voles as their reproductive performance, which is linked to their growth, is crucial to population growth and they are more likely to be affected by changes in food quality than males (Ostfeld 1985; Agrell *et al.* 1995). By manipulating the levels of silica in two species of grass we tested whether silica is an effective defence against voles. We hypothesized that silica will: (i) increase the abrasiveness of grass leaves reducing feeding by voles; (ii) reduce the digestibility of grasses to voles resulting in reduced growth rates of both juvenile and adult voles. We also tested whether vole feeding would induce silica defences by measuring the effect of vole damage on silica levels.

2. MATERIAL AND METHODS

(a) Study species and plant growth conditions

Short-tailed field voles (*M. agrestis* L.) are abundant in grasslands across Europe (Corbet & Harris 1991) and feed selectively on a range of grass species. We selected two grass species (*Festuca ovina* L. and *Lolium perenne* L.) found abundantly in European grasslands, which vary in their natural levels of foliar silica, nutrients and palatability to voles (table 1). Leaf samples of each grass species were collected from 10 replicate plants in 10 sites around East Sussex in August 2005, selected to represent a range of soil types and habitats, to confirm that our manipulated silica levels were within the range occurring in natural communities (table 1).

Grasses were grown in inert growth media (perlite), under glasshouse conditions for 12–28 weeks (15–25 °C, 16/8 h L/D). Plants for preference trials were grown individually into 5 $\times$ 5 $\times$ 5 cm plugs, while plants for vole growth performance trials and leaf digestibility experiments were grown in sward trays (20 $\times$ 30 $\times$ 5 cm). Grasses were watered every 3 days with 25 ml (individual plants) or 300 ml (sward trays) of Hoagland's solution. To half of the plants, 150 mg l⁻¹ of soluble silica was added as NaSiO₃·9H₂O to the Hoagland's solution (Cid *et al.* 1990).

(b) Feeding preference

Voies used in feeding trials were all non-lactating adults from a captive bred colony fed on a standard dried diet (N = 2.97%, C : N = 16.4, SiO₂ = 2.64% dry matter: rabbit

maintenance diet, B&K Feeds Universal) and fresh leaves of *Triticum avenae* daily. Voies were limited to water and a dried diet for 24 h prior to the trials.

Intraspecific feeding preference trials ($n = 10$) with captive-bred, non-lactating adult voies used three individual grass plants (12 weeks old) per silica treatment per trial (*F. ovina* and *L. perenne*). The leaf area of each plant was measured before and after each trial (AM-200 leaf area meter, ADC BioScientific). Plants were randomly arranged in a 2 $\times$ 3 grid, placed in a vole cage with sawdust to the base of grasses. A single vole was added to each cage until approximately 50% leaf area was eaten.

(c) Growth of juvenile voies

Twenty-four newly weaned (between 7 and 11 g), captive-bred, female voies were randomly assigned to a specific diet treatment (*F. ovina* or *L. perenne*; low or high silica). Voies were caged individually in glass tanks (45 $\times$ 30 $\times$ 30 cm, containing sawdust, cotton wool bedding, water and a standard dried diet) and 12 g of fresh grass from the particular treatment added daily. Voies were weighed every 3–4 days over a two month period. Faecal samples were collected after one month, dried and analysed for total nitrogen content (see below).

(d) Growth of mature voies

Ten adult female voies (between 20 and 24 g) were caught from a field site close to the University of Sussex and kept for two months prior to digestion efficiency trials. Voies were kept individually (as above) and assigned sequentially to each diet treatment (*F. ovina* or *L. perenne*; low or high silica) for 7 days in a random order. Voies were given 15 g of fresh grass leaves daily. At the end of each diet treatment, voies were weighed and samples of faeces removed from each cage, dried and analysed for nitrogen content (see below).

(e) Silica induction

Festuca ovina and *L. perenne* plants were grown in compost under standard greenhouse conditions for three months. Plants were then randomly assigned to damaged or undamaged treatments ($n = 10$). For the vole damage treatment, plants were placed in a cage with a single vole until they were eaten to approximately 5 cm above ground level, which took approximately 5 h. The damage treatment was repeated every three weeks for 12 months. Plants were then left to regrow for six weeks before collecting leaf samples for silica analysis.

(f) Chemical and physical analyses

Foliar silica content ($n=10$) was determined by fusing oven dried leaf samples (approx. 0.2 g) in sodium hydroxide followed by analysis using the colorimetric silicomolybdate technique (Allen 1989). Total foliar phenolic content of dried leaf samples ($n=10$) was determined using the Folin-Ciocalteu method (Massey *et al.* 2005), using a tannic acid standard. Foliar and faecal nitrogen ($n=10$) was analysed using flash combustion of dried leaf samples (approx. 2.5 mg) followed by gas chromatographic separation (Elemental Combustion System; Costech Instruments) calibrated against a standard of composition $C_{26}H_{26}N_2O_2S$. Specific leaf area (equal to dry leaf mass per unit area) and leaf water content were measured for each species and silica treatment ($n=10$). Abrasiveness of grass samples ($n=10$) was determined using a development of the method described by Hammond & Ennos (2000). Fixed areas of grass leaves were ground against a Perspex plate using a modified Martindale abrasion and pilling tester (Model 404, James H. Heal, Halifax). The degree of abrasion by phytoliths on the Perspex was quantified using a laser perthometer to measure the depth of grooves scratched in the surface.

(g) Statistical analysis

Grass leaf characteristics (leaf abrasiveness, silica, nitrogen and phenolic contents) were compared between species and silica treatments using two-way ANOVAs. Feeding preference was calculated as the mean leaf area removed per treatment as a proportion of the total leaf area removed per trial. These data were then transformed using arcsine square-root to achieve normality, before being compared across silica treatments using paired t -tests. Growth rates of juvenile voles (derived from the slope of linear regressions lines: $r^2 = 73.9\text{--}96.8\%$) were compared between grass species and silica treatments using two-way ANCOVA, with initial vole weight as covariate. The proportion of nitrogen absorbed from grass (grass N – faecal N/grass N) was compared between species and silica treatments using two-way ANOVA. For adult voles, growth rates relative to initial weight and the proportion of nitrogen absorbed from grass were compared between species and silica treatments using two-way ANOVAs with vole as a random blocking term. For the induction experiment, the silica content of grasses after 16 months were compared between species and damage treatments using a two-way ANOVA.

3. RESULTS

(a) Leaf abrasiveness and feeding preference

The presence of silica in grass leaves led to increases of 29 and 42% in leaf abrasiveness in *F. ovina* and *L. perenne* (figure 1a), resulting in a reduction of 75 and 63% in feeding by field voles on the two grass species, respectively (figure 1b).

(b) Vole growth performance and leaf digestibility

Juvenile female voles reared for two months on leaves of *F. ovina* and *L. perenne* containing high silica levels had 41 and 35% lower growth rates, respectively, than voles reared on low silica leaves (figure 2a, ANCOVA: initial weight (cov) $F_{1,19}=0.03$ and $p=0.872$, species $F_{1,19}=11.07$ and $p=0.004$, silica $F_{1,19}=35.42$ and $p<0.001$, and species $\times$ silica $F_{1,19}=0.05$ and $p=0.831$). The same trend was evident in mature female voles, where silica

Figure 1. (a) Leaf abrasiveness ($\pm$ s.e.) of *Festuca ovina* and *Lolium perenne* from high and low silica treatments. Letters above bars denote significant differences between treatments (Tukey; $p<0.05$). (b) Vole feeding preference comparing grasses from high and low silica treatments based on trial means ($\pm$ s.e.), paired t -test: *** $p<0.001$.

reduced their growth rate by over 50% during a period of only 7 days of exposure (figure 2b, ANOVA: vole $F_{9,27}=0.41$ and $p=0.917$, silica $F_{1,27}=4.51$ and $p=0.043$). Over the 7 day period there was no difference in the response on different grass species (vole $F_{9,27}=0.41$ and $p=0.917$, species $F_{1,27}=1.68$ and $p=0.206$).

Comparisons of foliar and faecal nitrogen levels between silica treatments revealed that juvenile voles reared on high silica diets absorbed proportionally less nitrogen than voles reared on low silica diets (figure 2c ANOVA: species $F_{1,20}=125.43$ and $p<0.001$, silica $F_{1,20}=8.64$ and $p=0.008$, species $\times$ silica $F_{1,20}=0.24$ and $p=0.63$). For mature voles, high silica diets resulted in 38% and 19% less nitrogen absorbed from *F. ovina* and *L. perenne* leaves, respectively (figure 2d ANOVA: vole $F_{9,27}=6.83$ and $p<0.001$, species $F_{1,27}=67.41$ and $p<0.001$, silica $F_{1,27}=52.97$ and $p<0.001$, species $\times$ silica $F_{9,27}=4.36$ and $p=0.046$). As the silica treatments had no detectable effects on either leaf nitrogen or phenolic contents (table 1), these results suggest that it was the presence of silica that led to the reduced digestibility of grass leaves.

Figure 2. Growth rates ($\pm$ s.e.) of (a) juvenile voles over two months and (b) mature voles over 7 days on high and low silica diets of *F. ovina* and *L. perenne*. Proportion of nitrogen absorbed from grass leaves by (c) juvenile and (d) mature voles (nitrogen absorbed = $\text{N in grass} - \text{N in faeces} / \text{N in grass}$). Letters above bars denote significant differences between treatments (Tukey; $p < 0.05$).

(c) Silica induction

Feeding by voles led to an increase in foliar silica content of over 400% in both *F. ovina* and *L. perenne* compared with undamaged plants (figure 3, ANOVA: species $F_{1,36} = 12.88$ and $p < 0.001$, damage $F_{1,36} = 134.47$ and $p < 0.001$, species $\times$ damage $F_{1,36} = 5.22$ and $p = 0.028$).

4. DISCUSSION

We have demonstrated that higher silica levels in grass leaves resulted in both increased abrasiveness of leaves and reduced feeding by voles. These results suggest that the elevated abrasiveness due to silica is an effective feeding deterrent to vertebrate herbivores and support the suggestion that silica may have played a significant role in driving the evolution of dental adaptations found in mammalian grazers (Janis & Fortelius 1988; Jernvall & Fortelius 2002; Prasad *et al.* 2005). Silica levels in natural grassland systems display intraspecific spatio-temporal variation, reflecting variation in biotic factors, such as grazing levels (McNaughton & Tarrants 1983; Brizuela *et al.* 1986), and abiotic factors, such as pH and soil type (O'Regain & Mentis 1989). Hence, in addition to interspecific variation in silica content, there is potentially a large degree of intraspecific variation in silica between environments. We propose that voles can detect and respond to these variations in silica content between plants and, in combination with the highly selective nature of vole feeding, this could ultimately result in changes in plant species abundance and hence the composition of

Figure 3. Levels of silica in vole grazed and undamaged control plants of *F. ovina* and *L. perenne*. Values are means ($\pm$ s.e.); bars not sharing a common letter differ significantly (Tukey's test $p < 0.05$).

plant communities (Moen *et al.* 1993; Howe & Brown 1999; Clay 2001; Howe & Lane 2004).

We have also demonstrated, for the first time, that silica in grass leaves can reduce the growth performance of voles, and our results strongly suggest that the mechanism by which this occurs is a reduction in the amount of

nitrogen voles can absorb from grasses. Despite the nitrogen levels, phenolic concentration and water content of grass species being unaffected by the silica treatments, mature voles absorbed 38 and 19% less nitrogen when fed on high silica plants of *F. ovina* and *L. perenne*, respectively. The presence of silica reduced the nitrogen available to voles in food where nitrogen is already limiting for a herbivore with a small body mass, a high metabolic rate and a low capacity to increase consumption in the face of declining food quality (Zynel & Wunder 2002).

The substantial reductions in the growth rates of juvenile voles we observed could have implications for the time taken to reach sexual maturity, and hence for population growth. Similarly, despite the relatively short time-scale of the feeding trials with mature voles, we still found evidence that silica in grasses significantly affected body mass and hence, potentially, reproductive output. It is well-established that female body mass and growth rate are prime determinants of both the number of litters and their size in voles as well as adult survival (Krebs & Myers 1974; Cole & Batzli 1979; Ergon *et al.* 2001; Ylonen & Eccard 2004). For example, when studying the factors affecting body mass of voles in a reciprocal transplant experiment between populations of varying body masses, Ergon *et al.* (2001) found that the immediate environment was the major predictor of vole condition. The factors in the immediate environment responsible were not identified, but we have shown that food quality, specifically the silica content of grasses, is worthy of further investigation as a potential mechanism for these effects.

Vole populations, as with many small mammals, are characterized by cyclical shifts in local population densities (Agrell *et al.* 1995; Klemola *et al.* 2000a), but the mechanisms underpinning these fluctuations are still a topic of debate (Agrell *et al.* 1995; Klemola *et al.* 2000b; Jiang & Shao 2003; Stenseth *et al.* 2003; Hornfeldt *et al.* 2005). Previously, many studies have focused on predator–vole interactions as the prime mechanism determining vole cycles (Hansson 1987; Henttonen *et al.* 1987; Hanski *et al.* 1993; Klemola *et al.* 2000a). However, more recent evidence has suggested that changes in environment, including reductions in food quality, may also play a significant role (Agrell *et al.* 1995; Turchin & Batzli 2001; Ergon *et al.* 2001). Some aspects of food quality have been measured with respect to vole population growth (Agrell *et al.* 1995; Klemola *et al.* 2000a), but no studies have examined the effects of silica, the principal defence in their food plants. Although previous studies have found positive correlations between grazing levels and grass silica content (McNaughton & Tarrants 1983; McNaughton *et al.* 1985; Brizuela *et al.* 1986; Banuelos & Obeso 2000), we provide the first experimental demonstration that feeding by voles leads directly to dramatic increases in the levels of silica in grass leaves. Hence, this study highlights how silica induction due to grazing could contribute to cyclic dynamics in voles: in years of high population densities, high grazing pressure will lead to induction of silica defences and a subsequent reduction in vole body mass, reproductive outputs and survival.

The role of defences in the dynamic feedbacks between woody plants and mammalian herbivores has been well studied (Bryant *et al.* 1991), but to date, no such plant-based mediation has been demonstrated for grasses and their herbivores. Further, no previous studies have

investigated the effects of silica on herbivore performance, despite it being the principal defence in grasses. Our study demonstrates that silica is induced by 400% in response to vole grazing, and that vole feeding is reduced by 75% and growth rates by 40% on diets containing silica. Effects of this magnitude seem likely to have significant ecological impacts on vole populations if they occur in the field. Silica-based defences thus have the potential to impact on both grassland community composition and on the population dynamics of mammalian herbivores.

We thank R. Ennos and the University of Manchester for conducting abrasion tests and silica analyses, and S. Valbonesi for C/N-analyses. L. Bromham, J. Davison, E. John, X. Lambin, K. Massey, T. Roper, N. Smith and two anonymous referees commented on drafts of the manuscript. Funding was provided by the Natural Environment Research Council, UK (NER/A/S/2001/01144).

REFERENCES

- Agrell, J., Erlinge, S., Nelson, J., Nilsson, C. & Persson, I. 1995 Delayed density-dependence in a small-rodent population. *Proc. R. Soc. B* **262**, 65–70.
- Allen, S. E. 1989 *Chemical analysis of ecological materials*, 2nd edn. Oxford, UK: Blackwell Science.
- Baker, G., Jones, L. H. P. & Wardrop, I. D. 1959 Cause of wear in sheep's teeth. *Nature* **184**, 1583–1584.
- Banuelos, M. J. & Obeso, J. R. 2000 Effect of grazing history, experimental defoliation, and genotype on patterns of silicification in *Agrostis tenuis* Sibth. *Ecoscience* **7**, 45–50.
- Batzli, G. O. & Cole, F. R. 1979 Nutritional ecology of microtine rodents—digestibility of forage. *J. Mammal.* **60**, 740–750.
- Brizuela, M. A., Detling, J. K. & Cid, M. S. 1986 Silicon concentration of grasses growing in sites with different grazing histories. *Ecology* **67**, 1098–1101. (doi:10.2307/1939834)
- Bryant, J. P., Provenza, F. D., Pastor, J., Reichardt, P. B., Clausen, T. P. & Dutoit, J. T. 1991 Interactions between woody-plants and browsing mammals mediated by secondary metabolites. *Annu. Rev. Ecol. Syst.* **22**, 431–446. (doi:10.1146/annurev.es.22.110191.002243)
- Cid, M. S., Detling, J. K., Whicker, A. D. & Brizuela, M. A. 1990 Silicon uptake and distribution in *Agropyron smithii* as related to grazing history and defoliation. *J. Range Manage.* **43**, 344–346.
- Clay, K. 2001 Symbiosis and the regulation of communities. *Am. Zool.* **41**, 810–824.
- Cole, F. R. & Batzli, G. O. 1979 Nutrition and population—dynamics of the prairie vole *Microtus ochrogaster*, in Central Illinois. *J. Anim. Ecol.* **48**, 455–470.
- Corbet, G. B. & Harris, S. 1991 *The handbook of British mammals*. Oxford, UK: Blackwell Science.
- Demment, M. W. & Van Soest, P. J. 1985 A nutritional explanation for body-size patterns of ruminant and nonruminant herbivores. *Am. Nat.* **125**, 641–672. (doi:10.1086/284369)
- Ergon, T., Lambin, X. & Stenseth, N. C. 2001 Life-history traits of voles in a fluctuating population respond to the immediate environment. *Nature* **411**, 1043–1045. (doi:10.1038/35082553)
- Ergon, T., Speakman, J. R., Scantlebury, M., Cavanagh, R. & Lambin, X. 2004 Optimal body size and energy expenditure during winter: why are voles smaller in declining populations? *Am. Nat.* **163**, 442–457. (doi:10.1086/381940)

- Gali-Muhtasib, H. U., Smith, C. C. & Higgins, J. J. 1992 The effect of silica in grasses on the feeding-behavior of the prairie vole *Microtus ochrogaster*. *Ecology* **73**, 1724–1729. (doi:10.2307/1940024)
- Hammond, T. A. & Ennos, A. R. 2000 *Mechanical testing of the abrasiveness of grass leaves. Third plant biomechanics conference proceedings*. Stuttgart: Georg Thieme Verlag.
- Hanski, I., Turchin, P., Korpimäki, E. & Henttonen, H. 1993 Population oscillations of boreal rodents—regulation by mustelid predators leads to chaos. *Nature* **364**, 232–235. (doi:10.1038/364232a0)
- Hansson, L. 1987 An interpretation of rodent dynamics as due to trophic interactions. *Oikos* **50**, 308–318.
- Hartley, S. E., Nelson, K. & Gorman, M. 1995 The effect of fertilizer and shading on plant-chemical composition and palatability to Orkney voles *Microtus arvalis orcadensis*. *Oikos* **72**, 79–87.
- Henttonen, H., Oksanen, T., Jortikka, A. & Haukioja, V. 1987 How much do weasels shape microtine cycles in the northern Fennoscandian taiga. *Oikos* **50**, 353–365.
- Hjalten, J., Danell, K. & Ericson, L. 1996 Food selection by two vole species in relation to plant growth strategies and plant chemistry. *Oikos* **76**, 181–190.
- Hornfeldt, B., Hipkiss, T. & Eklund, U. 2005 Fading out of vole and predator cycles? *Proc. R. Soc. B* **272**, 2045–2049. (doi:10.1098/rspb.2005.3141)
- Howe, H. F. & Brown, J. S. 1999 Effects of birds and rodents on synthetic tallgrass communities. *Ecology* **80**, 1776–1781. (doi:10.2307/176568)
- Howe, H. F. & Lane, D. 2004 Vole-driven succession in experimental wet-prairie restorations. *Ecol. Appl.* **14**, 1295–1305.
- Janis, C. M. & Fortelius, M. 1988 On the means whereby mammals achieve increased functional durability of their dentitions, with special reference to limiting factors. *Biol. Rev. Camb. Philos. Soc.* **63**, 197–230.
- Jernvall, J. & Fortelius, M. 2002 Common mammals drive the evolutionary increase of hypsodonty in the Neogene. *Nature* **417**, 538–540. (doi:10.1038/417538a)
- Jiang, L. & Shao, N. 2003 Autocorrelated exogenous factors and the detection of delayed density dependence. *Ecology* **84**, 2208–2213.
- Klemola, T., Koivula, M., Korpimäki, E. & Norrdahl, K. 2000a Experimental tests of predation and food hypotheses for population cycles of voles. *Proc. R. Soc. B* **267**, 351–356. (doi:10.1098/rspb.2000.1008)
- Klemola, T., Norrdahl, K. & Korpimäki, E. 2000b Do delayed effects of overgrazing explain population cycles in voles? *Oikos* **90**, 509–516. (doi:10.1034/j.1600-0706.2000.900309.x)
- Krebs, C. J. & Myers, J. H. 1974 Population cycles in small mammals. *Adv. Ecol. Res.* **8**, 267–399.
- Marquis, R. J. & Batzli, G. O. 1989 Influence of chemical factors on palatability of forage to voles. *J. Mammal.* **70**, 503–511.
- Massey, F. P., Press, M. C. & Hartley, S. E. 2005 Long- and short-term induction of defences in seedlings of *Shorea leprosula* (Dipterocarpaceae): support for the carbon: nutrient balance hypothesis. *J. Trop. Ecol.* **21**, 195–201. (doi:10.1017/S0266467404002111)
- McNaughton, S. J. 1984 Grazing lawns—animals in herds, plant form, and coevolution. *Am. Nat.* **124**, 863–886. (doi:10.1086/284321)
- McNaughton, S. J. & Tarrants, J. L. 1983 Grass leaf silicification—natural-selection for an inducible defense against herbivores. *Proc. Natl Acad. Sci. USA* **80**, 790–791.
- McNaughton, S. J., Tarrants, J. L., McNaughton, M. M. & Davis, R. H. 1985 Silica as a defense against herbivory and a growth promotor in African grasses. *Ecology* **66**, 528–535. (doi:10.2307/1940401)
- Moen, J., Gardfjell, H., Oksanen, L., Ericson, L. & Ekerholm, P. 1993 Grazing by food-limited microtine rodents on a productive experimental plant community—does the green desert exist. *Oikos* **68**, 401–413.
- O'Reagain, P. J. & Mentis, M. T. 1989 Leaf silicification in grasses—a review. *J. Grassl. Soc. South Afr.* **6**, 37–43.
- Ostfeld, R. S. 1985 Limiting resources and territoriality in microtine rodents. *Am. Nat.* **126**, 1–15. (doi:10.1086/284391)
- Prasad, V., Stromberg, C. A. E., Alimohammadian, H. & Sahni, A. 2005 Dinosaur coprolites and the early evolution of grasses and grazers. *Science* **310**, 1177–1180. (doi:10.1126/science.1118806)
- Stebbins, G. L. 1981 Coevolution of grasses and herbivores. *Ann. Mo. Bot. Gard.* **68**, 75–86. (doi:10.2307/2398811)
- Stenseth, N. C., Viljugrein, H., Saitoh, T., Hansen, T. F., Kittilsen, M. O., Bolviken, E. & Glockner, F. 2003 Seasonality, density dependence, and population cycles in Hokkaido voles. *Proc. Natl Acad. Sci. USA* **100**, 11 478–11 483. (doi:10.1073/pnas.1935306100)
- Turchin, P. & Batzli, G. O. 2001 Availability of food and the population dynamics of arvicoline rodents. *Ecology* **82**, 1521–1534. (doi:10.2307/2679797)
- Vicari, M. & Bazely, D. R. 1993 Do grasses fight back—the case for antiherbivore defenses. *Trends Ecol. Evol.* **8**, 137–141. (doi:10.1016/0169-5347(93)90026-L)
- Ylonen, H. & Eccard, J. A. 2004 Does quality of winter food affect spring condition and breeding in female bank voles (*Clethrionomys glareolus*)? *Ecoscience* **11**, 1–5.
- Zynel, C. A. & Wunder, B. A. 2002 Limits to food intake by the prairie vole: effects of time for digestion. *Funct. Ecol.* **16**, 58–66. (doi:10.1046/j.0269-8463.2001.00601.x)