Resuscitation using albumin in critically ill patients

Research in patients at high risk of complications is now needed

Research p 1044

Intravenous albumin has been widely used in the treatment of shock since the early 1940s. In a recent large European study on the occurrence of sepsis in acutely ill patients, 11% of patients received albumin at some point during their stay in intensive care.¹ Despite such widespread use, there are few randomised controlled trials on albumin administration in critically ill patients, and even fewer data on which subgroups of patients are likely to benefit most. Because low baseline serum albumin concentrations are associated with worse outcomes,²⁻³ data on the use of albumin in patients with low albumin are of considerable interest. In this week's *BMJ* a subgroup analysis of the saline versus albumin fluid evaluation (SAFE) study⁴ assesses the effect of baseline serum albumin concentration on outcomes of resuscitation.⁵

A meta-analysis published in the *BMJ* in 1998 suggested that resuscitation with albumin solutions rather than crystalloids increased the absolute risk of death by 6%.⁶ In 2004, prompted in part by the meta-analysis, the saline versus albumin fluid evaluation (SAFE) study compared 4% albumin with normal saline as the initial fluid for resuscitation in 7000 patients in intensive care.⁴ It found no significant difference in outcomes between treatments at 28 days.⁴

The results of the SAFE study are often interpreted to mean that albumin administration does not result in better outcomes than saline; however, the aim of the study was only to show that albumin administration is safe. All patients who needed a fluid challenge—for whatever reason—were therefore included, regardless of the underlying condition. In any unit using a logical clinical decision process, most patients randomised to the SAFE study would not receive albumin. It is therefore unreasonable to expect the study to show a benefit from albumin administration, except perhaps in subgroup analyses. The question that we need to answer is which subgroups of people are likely to benefit from such administration.

In this week's *BMJ*, a further analysis of the SAFE study assesses whether outcomes of resuscitation with saline or albumin are related to baseline serum albumin concentrations, predefined as serum albumin less than or greater than 25 g/l.⁵ The analysis confirmed the findings of previous studies,²⁻³ that a low albumin concentration is significantly associated with increased mortality after adjusting for baseline risk factors. This confirms the importance of assessing the effects of albumin administration in such patients. Patients with low serum albumin who received albumin

tended to have a lower mortality rate than those who received saline (23.7% v 26.2%, odds ratio 0.87, 95% confidence interval 0.73 to 1.05), but the difference was not significant.

Some aspects of the study merit consideration. About 40% of patients had an albumin concentration below 25 g/l. Such patients were older and were more likely to have severe sepsis, acute respiratory distress syndrome, or renal replacement therapy than people with higher albumin values. They were also more likely to be admitted for surgical reasons (which may explain why they were also less likely to be treated by mechanical ventilation). Surprisingly they did not have higher acute physiology and chronic health evaluation (APACHE II) scores than patients with higher albumin concentrations. It would have been useful if sequential organ failure assessment scores had been reported so that the degree of organ dysfunction in the two groups could be compared.

The next logical question would be whether administration of albumin was beneficial in patients with hypoalbuminaemia and a high risk of complications, such as those with severe sepsis or a high sequential organ failure assessment score, or both. This would be an analysis of a subgroup within a subgroup, however, with double the limitations of subgroup analyses. Unfortunately, this latest study provides little information on the use of albumin in such patients.⁵

The authors are correct when they reject the routine use of albumin in adult patients in intensive care, but their interpretation of the data may be too restrictive when they conclude that albumin and saline have similar effects across the range of albumin concentrations. The results may suggest that patients with a low serum albumin who receive albumin have a lower mortality rate than those who receive saline. It should be remembered that the SAFE study was not designed to show the beneficial effects of albumin administration in any particular group of patients.

Imagine a study with an identical design that evaluated the safety of blood transfusions. Such a study would probably show no benefit and perhaps a harmful effect of blood transfusions in an unselected group of patients. This would not indicate that blood transfusions should be abandoned, but rather that studies should focus on the subgroup of patients with anaemia and a high risk of complications.

It is now time to conduct a randomised controlled trial to evaluate the effects of albumin versus saline

administration in patients with hypoalbuminaemia and an increased risk of complications. In the meantime, insufficient evidence exists to support a ban on albumin administration in critically ill patients with hypoalbuminaemia.

J L Vincent *professor*

(jlvincen@ulb.ac.be)

Department of Intensive Care, Erasme University Hospital, 1070 Brussels, Belgium

Competing interests: None declared.

- 1 Vincent JL, Sakr Y, Reinhart K, Sprung CL, Gerlach H, Ranieri VM. Is albumin administration in the acutely ill associated with increased mortality? Results of the SOAP study. *Crit Care* 2005;9:R745-54.

- 2 Vincent JL, Dubois MJ, Navickis RJ, Wilkes MM. Hypoalbuminemia in acute illness: is there a rationale for intervention? A meta-analysis of cohort studies and controlled trials. *Ann Surg* 2003;237:319-34.
- 3 Goldwasser P, Feldman J. Association of serum albumin and mortality risk. *J Clin Epidemiol* 1997;50:693-703.
- 4 Finfer S, Bellomo R, Boyce N, French J, Myburgh J, Norton R. A comparison of albumin and saline for fluid resuscitation in the intensive care unit. *N Engl J Med* 2004;350:2247-56.
- 5 Saline versus Albumin Fluid Evaluation (SAFE) Study Investigators. Effect of baseline serum albumin concentration on outcome of resuscitation with albumin or saline in patients in intensive care units: analysis of data from the saline versus albumin fluid evaluation (SAFE) study. *BMJ* doi: 10.1136/bmj.38985.398704.7C.
- 6 Cochrane Injuries Group. Human albumin administration in critically ill patients: systematic review of randomized controlled trials. *BMJ* 1998;317:235-40.

doi: 10.1136/bmj.39029.490081.80

Glomerular filtration rate

Screening cannot be recommended on the basis of current knowledge

Low glomerular filtration rate predicts cardiovascular disease, end stage renal disease (a requirement for dialysis or transplantation), and death.¹ In a population based study reported in this week's issue, Hallan and colleagues describe the yield of different screening strategies aimed at identifying people with low glomerular filtration rates.¹ The guidelines of the United States kidney disease outcomes quality initiative define chronic kidney disease as glomerular filtration rate persistently lower than 60 ml/min/1.73 m², persistent urine abnormalities, or clinically relevant anatomical abnormalities, and they recommend measuring serum creatinine to determine estimated glomerular filtration rate in high risk groups.² The Kidney Disease Improving Global Outcomes Group (an international organisation that seeks to improve care and outcomes for people with kidney disease) and draft UK guidelines endorse the classification but do not, at present, look at screening.^{3 4}

All screening programmes cause physical, psychological, and social harm through the screening test itself, and through the investigation and treatment of abnormal results. No trials to show whether the benefits of screening for low glomerular filtration rate outweigh the harms have been conducted. So how should we proceed in the absence of such a trial?

The figure shows some areas in which the information needed to define a screening strategy is missing. Because low glomerular filtration rate is associated with cardiovascular disease, screening patients with established cardiovascular disease or risks for disease rather than unselected populations will lead to lower numbers needed to screen to identify one person.¹ However, many people identified this way will have indications for interventions that randomised controlled trials have shown to prevent cardiovascular disease, end stage renal disease, or death, which limits the potential for differential intervention (application of treatments that would not have been used otherwise). Lower numbers needed to screen can also be achieved by screening older people,¹ but the same consideration applies.

The screening test and "screen positive" threshold also need to be defined. Many authorities recommend using the estimated glomerular filtration rate.²⁻⁴ In the study by Hallan and colleagues, 20% of people with estimated glomerular filtration rates lower than 30 ml/min/1.73 m², but only 1-2% of those with values of 30-60 ml/min/1.73 m², progressed to end stage renal disease over eight years. Many people in the community (especially elderly people) with low glomerular filtration rate have disease that does not progress, which further limits the potential benefit of screening.^{1 5}

Identifying people with a low glomerular filtration rate may lead to harm from labelling. For example, in people screened for hypertension at work, positive results can result in absenteeism and reduced income.⁶

Little research has been conducted into which tests should be done to diagnose possible underlying kidney diseases in people with low glomerular filtration rate.

Because estimated glomerular filtration rate and serum creatinine are continuous variables, and because

Research p 1047

BMJ 2006;333:1030-1

Information needed to define a screening strategy for low glomerular filtration rate

of uncertainty about normal values and how they change with age, many people with results just above the screening threshold cannot be told that they have normal values. These people cannot be fully reassured, but do not benefit from additional treatment, and they will only be harmed by screening.

The purpose of screening is to identify people who should be treated differently. Angiotensin converting enzyme inhibitors,⁷ angiotensin II receptor blockers (in people with diabetes⁸), and low protein diets⁹ help prevent end stage renal disease in people with low glomerular filtration rate. Evidence of the benefit from angiotensin converting enzyme inhibitors and low protein diets comes largely from studies of people referred to nephrologists and may not generalise to people at lower risk.¹⁰ Two further issues exist—additional benefits from angiotensin converting enzyme inhibitors and angiotensin II receptor blockers will be limited to people not already taking them for other indications; and the intensive dietary education to prevent malnutrition that has been used in studies on low protein diets is not practical for large numbers.

Given the obstacles to the design of a screening strategy, what are the research priorities? More information is needed on how age, proteinuria, kidney function, diabetes, and blood pressure combine to predict progression to end stage renal disease, cardiovascular disease, and death in the general population. The proportion of people in candidate risk groups in whom treatment would be modified on the basis of data on kidney function must be defined. Modelling studies would identify whether potentially cost effective screening strategies exist; if so, these should be tested in randomised clinical trials.

In the meantime what should clinicians do? The study by Hallan and colleagues found that screening people with hypertension, diabetes, or age over 55 was the most effective strategy—93.2% (95% confidence interval 92.4% to 94.0%) of all cases of chronic kidney disease were detected, and the number needed to screen to find one case was 8.7 (8.5 to 9.0). Rather than screening, case finding should therefore continue in patients who have risks for or evidence of kidney disease, who are taking drugs that affect kidney

function, or whose comorbidities make knowledge of kidney function important. Wide use of angiotensin converting enzyme inhibitors in people who have cardiovascular disease or diabetes and another cardiovascular disease risk factor will automatically treat many people with low glomerular filtration rate and will prevent cardiovascular disease and death.¹¹ Maximising the proportion of hypertensive people who attain conventional blood pressure targets will prevent cardiovascular disease and death and is also likely to prevent end stage renal disease.

Catherine M Clase *associate professor of medicine*

(clase@mcmaster.ca)

Department of Clinical Epidemiology and Biostatistics, McMaster University, Hamilton, ON, L8N 1Y2 Canada

Competing interests: None declared.

- Hallan SI, Dahl K, Oien CM, Grootendorst DC, Aasberg A, Holmen J, et al. Screening strategies for chronic kidney disease in the general population: follow-up of cross sectional health survey. *BMJ* 2006 doi: 10.1136/bmj.39001.657755.BE.
- Levey AS, Coresh J, Balk E, Kausz AT, Levin A, Steffes MW, et al. National Kidney Foundation practice guidelines for chronic kidney disease: evaluation, classification, and stratification. *Ann Intern Med* 2003;139:137-47.
- Levey AS, Eckardt KU, Tsukamoto Y, Levin A, Coresh J, Rossert J, et al. Definition and classification of chronic kidney disease: a position statement from kidney disease: improving global outcomes (KDIGO). *Kidney Int* 2005;67:2089-100.
- Taal M, Tomson C. *Clinical practice guidelines: chronic kidney disease*. Petersfield: Renal Association, 2006. www.renal.org/guidelines/module1.html.
- John R, Webb M, Young A, Stevens PE. Unreferred chronic kidney disease: a longitudinal study. *Am J Kidney Dis* 2004;43:825-35.
- Johnston ME, Gibson ES, Terry CW, Haynes RB, Taylor DW, Gafni A, et al. Effects of labelling on income, work and social function among hypertensive employees. *J Chronic Dis* 1984;37:417-23.
- Jafar TH, Schmid CH, Landa M, Giatras I, Toto R, Remuzzi G, et al. Angiotensin-converting enzyme inhibitors and progression of nondiabetic renal disease. A meta-analysis of patient-level data. *Ann Intern Med* 2001;135:73-87.
- Brenner BM, Cooper ME, de Zeeuw D, Keane WF, Mitch WE, Parving HH, et al. Effects of losartan on renal and cardiovascular outcomes in patients with type 2 diabetes and nephropathy. *N Engl J Med* 2001;345:861-9.
- Fouque D, Laville M, Boissel JP. Low protein diets for chronic kidney disease in non diabetic adults. *Cochrane Database Syst Rev* 2006;(2):CD001892.
- Rahman M, Pressel S, Davis BR, Nwachuku C, Wright JT Jr, Whelton PK, et al. Renal outcomes in high-risk hypertensive patients treated with an angiotensin-converting enzyme inhibitor or a calcium channel blocker vs a diuretic: a report from the antihypertensive and lipid-lowering treatment to prevent heart attack trial (ALLHAT). *Arch Intern Med* 2005;165:936-46.
- Yusuf S, Sleight P, Pogue J, Bosch J, Davies R, Dagenais G. Effects of an angiotensin-converting-enzyme inhibitor, ramipril, on cardiovascular events in high-risk patients. The Heart Outcomes Prevention Evaluation Study Investigators. *N Engl J Med* 2000;342:145-53.

doi 10.1136/bmj.39030.594259.BE

Tackling childhood obesity

Requires a shift in social norms, not just an exercise programme

Research p 1041

Despite heightened interest in preventing obesity in childhood, the evidence base for the effectiveness of local school based programmes and changes in policy to combat childhood obesity is poor.¹⁻³ Even less research has been devoted to improving nutrition and physical activity in preschool and childcare settings, where further challenges exist, such as lack of uniformity between programmes and fragmented oversights by regulatory authorities.^{1,3} Yet increasing evidence shows that preschool practices can influence the overall energy expenditure of young children.^{4 w1}

In this issue of the *BMJ*, Reilly and colleagues address this gap in the literature with a rigorously designed, cluster randomised controlled trial of an intervention to promote physical activity in 36 Scottish nursery schools.⁵ Body mass index and physical activity were not affected by the intervention at six and 12 months. These results contrast with recent studies showing that increasing children's physical activity can reduce weight gain,^{6 w2} and that changes in the quality

Extra references w1-w4 are on bmj.com

BMJ 2006;333:1031-2

and quantity of school based physical education can increase children's activity,^{7 w3} although it is not the only trial to find no benefit on body composition.^{8 w4}

One weakness of the study is in the main outcome measure used. Body mass index does not correlate closely with changes in body composition (amount and distribution of body fat, bone mineral density, muscularity) in adults and school aged children. The association is probably even more tenuous in very young and rapidly growing children, especially as the authors used body mass index standard deviation scores, rather than the more widely used age and sex specific body mass index centiles.⁹ In addition, the authors acknowledge that the "dose" of physical activity they used may be too low to influence the overall amount or intensity of physical activity, the study's primary independent variable. And it is not clear whether the study was powered to disaggregate the data by sex and overweight status, which may reveal effects diluted across the full sample.

Ultimately, the success of interventions disseminated to a variety of settings hinges on effective implementation—selection and training of staff, organisational culture and support, continuing education, and technical assistance.¹⁰ The pilot study for Reilly and colleagues' trial did produce increases in physical activity.¹¹ This may be because the intervention was delivered by head teachers in the pilot study, whereas it was delivered by nursery staff in the larger trial to increase generalisability. In addition, it is not clear whether the socioeconomic status of the children in the pilot and main study were similar. If the children studied in the larger trial came from more affluent families than those in the pilot study, a ceiling effect might have prevented modest effects being apparent.

Despite these criticisms, we should not lose sight of the bigger picture. Reilly and colleagues' findings show once again that such interventions have only weak and transient positive effects on behaviour and ultimately weight. To promote activity in youth and redress childhood obesity, adults must act to influence children's daily environments, as decision makers, gatekeepers, opinion leaders, service providers, and role models. This will require widespread and comprehensive changes in social norms and values in nurseries, schools, and beyond.^{1 12} In today's fast paced society—where children aged 4-5 years are pushed in buggies

rather than allowed to walk—temporary or circumscribed environmental changes are unlikely to have "permanent" effects.

However, preschool settings could be useful as part of a comprehensive strategy to prevent obesity. Preschool organisational practices and policies vary greatly^{4 w1} and may influence young children's fitness.^{7 13} While childcare settings may have less developed infrastructures than primary and secondary schools, they also have more flexible schedules and curriculums and tend to be less bureaucratic, which may help in adaptation and dissemination of interventions.⁴ Continuing investment and investigation are needed to identify approaches that work.

Antronette Yancey *associate professor of health services*

(ayancey@ucla.edu)

Department of Health Services, UCLA School of Public Health, Box 956900, Los Angeles, CA 90095-6900, USA

Competing interests: None declared.

- 1 Institute of Medicine (US). *Progress in preventing childhood obesity: how do we measure up?* Washington, DC: National Academies of Sciences, 2006.
- 2 Katz DL, O'Connell M, Yeh MC, Nawaz H, Njike V, Anderson LM, et al. Public health strategies for preventing and controlling overweight and obesity in school and worksite settings: a report on recommendations of the Task Force on Community Preventive Services. *MMWR Recomm Rep* 2005;54:1-12.
- 3 Story M, Kaphingst KM, French S. The role of child care settings in obesity prevention. *Future Child* 2006;16:143-68.
- 4 Dowda M, James F, Sallis JF, McKenzie TL, Rosengard P, Kohl HW 3rd. Evaluating the sustainability of SPARK physical education: a case study of translating research into practice. *Res Q Exerc Sport* 2005;76:11-9.
- 5 Reilly JJ, Kelly L, Montgomery C, Williamson A, Fisher A, McColl JH, et al. Physical activity to prevent obesity in young children: cluster randomised controlled trial. *BMJ* 2006 doi: 10.1136/bmj.38979.623773.55
- 6 LeMura LM, Maziakas MT. Factors that alter body fat, body mass, and fat-free mass in pediatric obesity. *Med Sci Sports Exerc* 2002;34:487-96.
- 7 Datar A, Sturm R. Physical education in elementary school and body mass index: evidence from the early childhood longitudinal study. *Am J Public Health* 2004;94:1501-6.
- 8 Trost SG. School physical education in the post-report era: an analysis from public health. *J Teaching Physical Educ* 2004;23:216-35.
- 9 Centers for Disease Control and Prevention. National health and nutrition examination survey. CDC growth charts. www.cdc.gov/nchs/about/major/nhanes/growthcharts/datafiles.htm.
- 10 Yancey AK, Ory MG, Davis SM. Dissemination of physical activity promotion interventions in underserved populations. *Am J Prev Med* 2006;31(4S):82-91.
- 11 Reilly JJ, McDowell ZC. Physical activity interventions in the prevention and treatment of paediatric obesity: systematic review and critical appraisal. *Proc Nutr Soc* 2003;62:611-9.
- 12 Yancey AK, Fielding JE, Flores GR, Sallis JF, McCarthy WJ, Breslow L. Creating a public health infrastructure for physical activity promotion: a challenge to chronic disease control policy. *Am J Prev Med* (in press).
- 13 Fitzgibbon ML, Stolley MR, Schiffer L, van Horn L, Kaufer Christoffel K, Dyer A. Two-year follow-up results for Hip-Hop to Health Jr: a randomized controlled trial for overweight prevention in preschool minority children. *J Pediatr* 2005;146:618-25.

doi: 10.1136/bmj.39021.605683.BE