Jugular Foramen Arteriovenous Shunt with Subarachnoid Hemorrhage

Abstract—The authors report the case of a 37-year-old man with an extracerebral arteriovenous fistula at the skull base, revealed by subarachnoid and intraventricular hemorrhage. The malformation was fed by the neuromeningeal trunk of the ascending pharyngeal artery and drained into left laterobulbar veins. Embolization with bucrylate was performed and occluded totally the shunting zone. A 1-year follow-up angiogram confirmed the good stability of the result, the patient being asymptomatic. This case emphasizes the quality of results that can be obtained with bucrylate in arteriovenous fistulas presenting with hemorrhage. It confirms that the external carotid artery must be studied when dealing with intracranial hemorrhage. On the other hand, magnetic resonance imaging and angiography may depict vascular abnormalities but do not always indicate the shunting area, thus the pathologic type of the malformation.

Subarachnoid hemorrhage (SAH) in the adult is mainly due to rupture of intracranial aneurysm. Other causes, such as trauma, tumors, arteriovenous malformations, and blood dyscrasias, have also been described. We report the case of an extracerebral arteriovenous shunt (AVS) of the skull base revealed by SAH and treated by embolization, in which the exact location, extradural or intradural, could not be assessed either by magnetic resonance imaging (MRI) or angiography.

CASE REPORT

A 37-year-old man, without previous medical complaints, experienced during sexual intercourse sudden and violent headaches and vomiting, followed by problems of consciousness. He was admitted to the emergency room of the hospital on the same night, where subarachnoid bleeding was clinically suspected. This was confirmed by computed tomography that showed the hemorrhage mainly localized in the posterior fossa around the brainstem and into the ventricular system (Fig. 1). The patient recovered slowly and his neurologic examination was normal 48

hours after the acute onset of the symptoms. A four-vessel cerebral angiography was performed and failed to show evidence of intracranial aneurysm. However, a localized spasm on the anterior communicating artery was suspected. The patient was submitted to MRI 25 days after his acute episode; serpiginous abnormal vascular structures were visualized on the left side of the medulla oblongata in an extra-axial position (Fig. 2). No pathologic foci, either in T₁ nor in T₂ WI could be described in the brainstem.

A cerebral vascular malformation was suspected and the patient was referred to our group. Following the regional protocol,¹ angiography demonstrated an AVS closely related to the left hypoglossal canal and fed by the neuromeningeal trunk of the ascending pharyngeal artery (Fig. 3). The venous drainage occurred into the laterobulbar veins with reflux into the basal veins bilaterally and into the vein of Galen. Left vertebral angiogram was considered normal.

No definitive proof could be obtained as to whether the AVS was located on the dura or in the subarachnoid space.² However it was believed that this information was not necessary to treat the patient safely. Endovascular

Figure 1. Nonenhanced computed tomography scan showing intraventricular and subarachnoid hemorrhage.

treatment of this fistula was proposed and realized 8 days later under general anesthesia. A 4 F catheter was positioned in the neuromeningeal trunk and 0.3 cc of a 50% mixture of NBCA and Lipiodol with tantalum powder was injected in situ. Control angiogram in the ascending pharyngeal artery failed to opacify the vascular malformation (Fig. 4). The venous pouches were still injected during left vertebral angiography via the normal interterritorial anastomoses, but no evidence of venous filling into the posterior fossa was demonstrated. The procedure was well tolerated clinically.

Control MRI performed 8 days later revealed a frank decrease in the size of the laterobulbar venous pouches without assessing their precocious thrombosis. One month following embolization, a selective angiogram failed to demonstrate any residual shunt and assessed the thrombosis of the previously noted venous pouches. The stability of the occlusion was confirmed by a final angiogram 1 year later (Fig. 5). The patient is actually doing well and is free of any neurologic symptoms.

DISCUSSION

Our case emphasizes the good quality of results that may be obtained by embolization in AVS. It also illustrates that so-called angiographically occult arteriovenous malformations may not be discovered because of incomplete angiographic protocols. The external carotid artery (ECA)

Figure 2. Magnetic resonance imaging studies of the brain in sagittal (A: T₁ WI) and axial (B: T₂ WI) views. Flow-void serpiginous structures are detected in the subarachnoid space on the left side of the medulla oblongata (small arrow). The brainstem appears normal. These pathologic structures were thought to correspond to abnormal vascular channels localized near the foramen occipitalis.

Figure 3. Selective injection of the left ascending pharyngeal artery in lateral view. The neuromeningeal trunk appears enlarged (small arrow) and feeds an arteriovenous malformation localized near the jugular foramen that drains into congestive laterobulbar veins (arrows). Reflux into posterior fossa veins is also noticed with opacification of the vein of Galen (double arrow) and the straight sinus (triple arrow).

should always be studied and included in the investigation of subarachnoid hemorrhage if no aneurysm is found. The notion that a branch of the ECA may vascularize an AVS complicated by intradural hemorrhage expresses an etiopathogenic and anatomic concept.

These hemorrhages only happen in AVS draining into cerebral veins. This type of complication is never seen if the venous drainage is exclusively sinusal. Passive reflux into cortical veins may happen because of obstructed or thrombosed sinus; this may lead to retrograde venous hyperpression leading to neurologic deficits or to venous rupture, causing intracranial bleeding.³⁻⁵ Only some rare situations drain of necessity and directly into cerebral veins.⁶

The fact that a meningeal artery vascularizes an AVS does not obligatorily prove that this is localized on the dura. There are two types of meningeal vessels: the pure meningeal and the neuromeningeal arteries. If a pure meningeal artery vascularizes a cerebral lesion, it proves the pathologic communication between the lesion itself and the dural structures (meningeal adherences). On the contrary, the neuromeningeal arteries vascularize normal transdural structures (radicular arteries or cranial nerve arteries); they anastomose on the surface of the central nervous system with their homologues arising from pial cerebral arteries. In these cases, their arterial participation with an AVS draining into cerebral veins is consistent with

Figure 4. Immediate angiographic control in the left ascending pharyngeal artery after embolization of the neuromeningeal trunk. The shunt is no longer opacified.

Figure 5. One year follow-up control angiogram in the left ascending pharyngeal artery (A) and left vertebral artery (B). The stability of the total occlusion of the arteriovenous shunt (AVS) is assessed. The venous pouches are not opacified and totally thrombosed. Figure continued on next page)

an intradural extracerebral lesion. This also means that superficial cerebral AVS may be fed only by ECA branches. Reciprocally, when a prominent pial branch accompanies the transdural structure, a meningeal lesion can be exclusively vascularized by a cerebral artery. In the case we report, although we are able to identify the arteriovenous junction because of the alteration of the vascular caliber, we cannot localize precisely the lesion on the dura as being outside or inside the dura, on the hypoglossal nerve, or even on the lateral wall of the medulla oblongata. The major point in the analysis of the angioarchitecture of the malformation is the identification of the arteriovenous junction. Its exclusive and limited occlusion guarantees the therapeutic safety, this being surgical or endovascular with NBCA. Furthermore, the dural or intradural localization is only of surgical interest because this difference

does not influence the endovascular approach. It is the reason why our embolization protocol of dural arteriovenous fistulas has been followed¹ and permitted in this case, as in others, an anatomic cure of the malformation. This total exclusion must always be looked for in cases of arteriovenous malformations with hemorrhagic accidents. In our experience⁷ this could only be obtained in a definitive and stable way with liquid agents (NBCA or bucrylate). Embolization with particles can obtain this result in a few selected situations. In the great majority of cases, it leads to transient occlusions of the catheterized and embolized arterial feeders, without occlusion of the proximal draining vein. Therefore recruitment of adjacent arteries and secondary recanalization of the previously occluded ones occurs, leading to progressive complication of the angioarchitecture and of therapeutic challenge.

Figure 5, cont. The venous phase of the vertebral angiogram can be considered as normal (C), the veins previously draining the AVS drain normal neural tissue.

REFERENCES

1. Lasjaunias P, Berenstein A: Surgical neuroangiography, vol 1. Functional Anatomy of the Craniofacial Arteries. Berlin: Springer-Verlag, 1987
2. Lasjaunias P, Berenstein A, Moret J: Signification de la vascularisation durale des lésions du système nerveux central. *J Neuroradiol* 10:31-42, 1983
3. Lasjaunias P, Chiu M, Terbrugge K, Tolia A, Hurth M, Bernstein M: Neurological manifestations of intracranial dural arteriovenous malformations. *J Neurosurg* 64:724-730, 1986
4. Malik GM, Pearce JE, Ausman JL, Mehta B: Dural arteriovenous malformations and intracranial hemorrhage. *AJNR* 6:982, 1985
5. Harding AE, Kendall B, Leonard JJK, Johnston MH: Intracerebral hemorrhage complicating dural arteriovenous fistula. Report of 2 cases. *AJNR* 7:188, 1986
6. Piske RL, Lasjaunias P: Extracranial dural arteriovenous malformations. Report of 3 cases. *Neuroradiology* 30:426-432, 1988
7. Lasjaunias P, Berenstein A: Dural arteriovenous malformations (DAVMs) In: Surgical Neuroangiography, vol 2. Endovascular Treatment of Craniofacial Lesions. Berlin: Springer-Verlag, 1987, pp 273-315