Acute Postoperative Hydrocephalus Following Translabyrinthine Craniotomy for Acoustic Neuroma Resection

Abstract—Acute mental status changes following craniotomy for acoustic tumors demand prompt evaluation and treatment to avoid serious morbidity and mortality. Two cases of acute obstructive hydrocephalus complicating the postoperative period following translabyrinthine craniotomy are presented. Diagnosis is made with noncontrast computed tomography scanning. Treatment is rendered at the bedside with placement of a ventriculostomy. Diagnosis and management implications for acoustic tumor patients are discussed. (*Skull Base Surgery*, 5(3): 143–148, 1995)

The mortality rate following resection of acoustic neuromas has dropped dramatically over the past three decades. Advanced microsurgical techniques and experience make life-threatening intraoperative complications rare. Postoperative deterioration of mental status, however, remains an occasionally encountered emergency that demands prompt, appropriate evaluation and treatment.

The differential diagnosis of mental status changes in the postoperative period includes metabolic, medication induced, ischemic, infectious, venous insufficiency, and mass effect etiologies.^{1,2} Interruption of blood supply to brain parenchyma or hematoma have, in our experience, accounted for most cases of mental status changes in the postoperative period. Although rare, acute hydrocephalus should be included in the differential diagnosis as well. We present the first two cases in a series of nearly 3500 acoustic neuromas in which acute hydrocephalus in the immediate postoperative period has been responsible for rapid patient deterioration.

CASE REPORTS

Case 1

A 62-year-old woman with a 3 cm left superior vestibular nerve neuroma underwent uncomplicated translabyrinthine craniotomy for tumor resection. Preoperatively, the patient was devoid of brainstem signs except for ipsilateral decreased corneal sensation. She awoke from surgery in a normal fashion, responding appropriately to questions and with normal facial nerve function. Postoperative medications included only intramuscular nalbuphine (6 mg) and intravenous Phenergan (12.5 mg). By late evening, she became increasingly somnolent and was unresponsive to all but painful stimuli by 2 AM. Gaze was dysconjugate, but pupils were small, symmetrical, and reactive. She remained hemodynamically stable but displayed disordered respiration. Endotracheal intubation followed. Third degree nystagmus persisted un-

changed throughout the first 2 postoperative days. Electrolytes, glucose, and arterial blood gases were normal. An emergent computed tomography (CT) scan showed blood in the fourth ventricle, aqueduct, and third ventricle with no cerebellopontine angle hematoma or shift. The ventricular system was dilated compared with the preoperative magnetic resonance imaging scan, consistent with acute obstructive hydrocephalus (Figs. 1, 2). Preoperative ventricle size was within normal limits (1.3 and 1.6 cm).

Emergent ventriculostomy relieved the elevated intraventricular pressure. Within several minutes, the patient's mental status returned to normal. Repeat CT scanning with the patent ventriculostomy in place showed reduction in ventricular size (Fig. 3).

Drainage was maintained for 72 hours and then clamped. Repeat CT with the clamped drain in place 24 hours later showed no recurrence of ventricular dilation. The ventriculostomy was removed, and the remainder of the hospitalization was without event. She had no residual neurologic deficit and maintained normal facial function.

Case 2

A 75-year-old woman with a 2.5 cm left acoustic neuroma originating on the inferior vestibular nerve under-

went uncomplicated translabyrinthine craniotomy. Preoperatively, the patient had no brainstem signs. She awoke normally after surgery with intact facial nerve function and was taken to the intensive care unit. Despite normal vital signs, mental status deterioration began 6 hours postoperatively. Over the next 2 hours, the patient progressed from somnolence to unresponsive to all but painful stimuli. Hemodynamic and respiratory stability continued. Nystagmus in all fields of gaze persisted. Electrolytes, glucose, and arterial blood gases were normal. A CT scan revealed dilated ventricles and blood in the third and fourth ventricles with abdominal fat filling the surgical defect (Figs. 4, 5).

Ventriculostomy to the right lateral horn was performed with return of cerebrospinal fluid (CSF) under high pressure. Mental status promptly returned to normal with the patient responding appropriately to questions within a few minutes. A CT performed with the patent ventriculostomy in place showed a marked reduction in ventricular size (Fig. 6).

CSF drainage was maintained for 72 hours. The ventriculostomy was then clamped for 24 hours and repeat CT showed no increase in ventricular size. The catheter was then removed and repeat scanning in 72 hours showed stable ventricular size. The remainder of the hospital course was uneventful, and the patient had no residual neurologic deficit.

Figure 1. Case 1. Acute postoperative obstructive hydrocephalus with ventricular dilation. (Reprinted with permission of House Ear Institute.)

Figure 2. Case 1. Surgical defect with abdominal fat graft. No cerebellopontine angle blood is present. Hematoma is seen in the fourth ventricle. (Reprinted with permission of House Ear Institute.)

Figure 3. Case 1. Ventriculostomy in place with reduction of ventricular size. (Reprinted with permission of House Ear Institute.)

Figure 4. Case 2. Acute postoperative obstructive hydrocephalus with ventricular dilation. (Reprinted with permission of House Ear Institute.)

Figure 5. Case 2. Surgical defect with abdominal fat graft protruding slightly into the cerebellopontine angle. Blood is seen in the fourth ventricle. (Reprinted with permission of House Ear Institute.)

Figure 6. Case 2. Decompression of ventricles with ventriculostomy catheter in place. (Reprinted with permission of House Ear Institute.)

DISCUSSION

Immediate evaluation of acute postoperative mental status changes following neurologic surgery is paramount to keep morbidity and mortality to a minimum. Acute hydrocephalus presents in a fashion similar to other causes of encephalopathy and should be included in the differential diagnosis. Both of these cases developed problems on the night of surgery 6 to 8 hours after extubation. Noncontrast CT scans allowed identification of acute hydrocephalus and exclusion of cerebellopontine angle hematoma.

Rapidly deteriorating mental status in the postoperative period should prompt immediate evaluation. The differential diagnosis includes several different entities capable of producing encephalopathy. Metabolic causes such as hyponatremia, hyperglycemia, hypoglycemia, hypoxia, and hypercapnia are examples. The prudent physician will order immediate blood studies when first notified of the problem to rule these causes out in a timely manner. Medication-induced consciousness changes should be obvious when inspecting the medication sheet. Infectious etiologies typically occur later in the postoperative course and are accompanied by fever. Meningitis infrequently causes deterioration of mental status over minutes to

hours. Venous insufficiency results in cerebral edema and a steadily progressing and sometimes rapid loss of consciousness. Rarely, a patient will awaken from anesthesia to experience a cerebrovascular accident in the postoperative period. Most cases of stroke are evolving from an intraoperative interruption of blood flow and are obvious on initial assessment in the intensive care unit.

By far, the most common cause of deterioration in mental status in the immediate postoperative period is hematoma with mass effect. An expanding hematoma in the cerebellopontine angle is a life-threatening problem for the patient. Handled correctly, however, removal of the clot and hemostasis usually leaves no residual neurologic deficit and returns consciousness to normal.

Left to enlarge, posterior fossa hematoma compresses the brainstem and the vital centers of respiration and blood pressure control or may cause herniation through the foramen magnum. The physician must be prepared to institute artificial respiration and to support circulation long enough to allow the mass to be removed. In cases of impending death, we sterilely open the wound at the bedside, allowing decompression of the posterior fossa by removing the abdominal fat placed in the mastoidectomy/labyrinthectomy defect. The cerebellopontine angle is not exposed until in the operating room. This maneuver can temporize long enough to assemble the operative team

and to prepare the operative suite for clot evacuation, saving the patient's life.

Fortunately, most cases of mental status deterioration progress at a rate that allows evaluation without such urgent lifesaving maneuvers. If the clinical situation permits, noncontrast CT scanning allows identification of the problem producing mental status changes in a high percentage of cases. Acute hydrocephalus and cerebellopontine angle hematoma produce distinct pathognomonic findings on radiographs. In these two cases, we detected clot in the third and fourth ventricles as well as the sylvian aqueduct, producing obstruction of CSF egress into the posterior fossa.

Exploration of the operative field of each of these patients was avoided. Presumably, surgical exploration would not have improved the acute obstructive hydrocephalus or the patient's mental status deterioration. Identification of acute obstructive hydrocephalus is important to allow appropriate treatment. Furthermore, treatment can be undertaken promptly at the bedside, avoiding a return to the operating room.

Both patients responded dramatically and promptly to alleviation of increased intraventricular pressure. In our patients, ventricular drainage was maintained for 48 to 72 hours to allow clot retraction or lysis to reestablish fourth ventricular to subarachnoid CSF flow. Clamping the ventriculostomy for a period of 24 hours before its removal and repeating the CT scan ensures that this pathway has been reestablished. Neither patient has had any

long-term complications. Hospitalization was not prolonged in either instance.

CONCLUSION

Acute postoperative mental status changes following posterior fossa surgery demand prompt evaluation and treatment. Acute obstructive hydrocephalus due to fourth ventricle clot can present in a similar fashion to more common complications, such as stroke or hematoma, and should be considered when evaluating such patients. CT scanning correctly identifies the problem when the clinical situation safely allows a trip to radiology. Insertion of a ventriculostomy normalized intraventricular pressure, returning mental status to normal in both of these cases. Clot retraction or lysis reestablished CSF flow from the fourth ventricle in each case by 72 hours. Removal of the ventriculostomy was possible in each of the patients without long-term sequelae.

REFERENCES

1. Kletzkner GR, Smith PG, Backer RJ, Leonetti JP: Complications in neurotological surgery. In Jackler RK, Brackmann DE (eds): *Neurotology*. St. Louis: Mosby, 1994, pp 713-727
2. Horwitz NH, Rizzoli HV: *Postoperative Complications of Intracranial Neurological Surgery*. Baltimore: Williams & Wilkins, 1982, pp 1-34