Giant-Cell Tumors of the Sphenoid Bone in Four Children: Radiological, Clinical, and Pathological Findings

ABSTRACT—We report the clinical, pathological, and radiological findings of giant cell tumor of the sphenoid bone in four children aged 10 to 16 years. The most common clinical finding was headache, followed by cranial nerve abnormalities. The computed tomographic findings of giant cell tumor consist of a lytic defect, sharply marginated with no sclerosis associated with an expansile, homogeneous mass (isodense with muscle). In the differential diagnosis, the lesion most similar to giant cell tumor is giant cell granuloma. The different histopathological features of the two lesions are discussed, along with other lesions, in the differential diagnosis. The magnetic resonance features consist of a mass with low signal intensities on T_1 - and T_2 -weighted images associated with moderate enhancement after introduction of gadolinium.

Giant-cell tumor (GCT) of bone comprises about 3-7% of bone tumors.^{1,2} GCT involves the end of long bones (70% and 90% of cases),³ the sacrum (5-10%), hands and feet (4%), vertebral bodies (2-5%), and the head and neck area, including skull (2%).^{2,4-6} In the head and neck region, GCT occurs predominantly in the mandible^{7,8} and maxilla^{9,10} and less commonly in the sphenoid bone,^{11,12} ethmoid bones,¹³ temporal bones,¹⁴⁻²¹ zygoma,²² and frontal bone.²³ According to several reviews of GCT of bone published through 1991, the total reported number of tumors involving the skull is 87.²² Twenty additional cases have been reported from 1991 up to 1996.^{14,22-35} The selective and preferential occurrence of GCT in the sphenoid bone is thought to be based on the endochondral ossification of this bone as opposed to membranous development of other skull bones. There is a female predominance. The

age at presentation ranges from the first to the eighth decade, with the majority of GCT occurring in the second to fourth decades. The tumor is believed to arise from the nonosteogenic stromal cells of the bone marrow.³⁶

Computed tomography (CT) and magnetic resonance imaging (MRI) are the optimal imaging modalities for assessment of GCT of bone, including the base of the skull.³⁷⁻⁴²

METHODS

This article describes the clinical, pathological, and radiological findings of four pediatric cases (aged 10 to 16 years) of GCT of the sphenoid bone admitted for proton beam therapy at the Radiation Medicine De-

partment of the Massachusetts General Hospital. These four patients had axial and coronal contrast CT images. MRI was performed with T₁- and T₂-weighted sequences following the administration of gadolinium (Gd). The literature of GCT in the head and neck area is reviewed, and the occurrence within the sphenoid bone is related to other sites of involvement.

CASE HISTORIES

Case 1

A 15-year-old white female complained of right frontal headache associated with an ear infection 1 year prior to admission. She developed numbness above her right eye and experienced more frequent and severe headaches. A systemic evaluation revealed no neurological findings or evidence of an endocrine disorder. Her past medical history was noncontributory. CT and MRI examinations performed in October and November 1991, respectively, demonstrated a large, homogeneous mass involving the basisphenoid, sphenoid sinus, and sella, with extension into the prepontine cistern and nasopharynx (Fig. 1). On MRI, the signal intensities were low on T₁- and T₂-weighted images. Moderate enhancement occurred after injection of Gd.

On November 14, 1991, a bifrontal craniotomy was performed, and a 5 × 3 cm mass was found in the sphenoid bone, including the sphenoid sinus. The histopathologic examination was benign GCT. The patient was reoperated on March 18, 1992, because of an increase in size of the residual disease, with further removal of tumor. Subsequently, in March 1992 combined photon and proton beam radiation was carried out.

Case 2

A 16-year-old white male was admitted with a 4-month history of progressive diplopia, headaches, nausea, and vomiting. On ophthalmological examination the left eye was paralyzed on medial and lateral gaze, and there was a lid lag. This was secondary to a partial deficit of the ocular motor nerve and trochlear nerve on the left. The remaining ophthalmological and neurological examinations were normal, and the past medical history was noncontributory. In October 1991, a CT study was carried out, which demonstrated a homogeneous, expansile mass with lytic bone destruction involving the clivus, sphenoid sinus, petrous apex, and medial aspect of the left middle cranial fossa (Fig. 2A). This study was followed by an MRI scan in April 1993, which demonstrated the large tumor in the clivus and sphenoid sinus with extension to the left cavernous sinus and adjacent middle cranial fossa. The GCT revealed low signal in-

tensity on the T₁- and T₂-weighted images with moderate enhancement after Gd introduction (Fig. 2B-E).

On May 4, 1993, the patient underwent a bilateral medial maxillotomy with partial removal of the tumor. The histopathological diagnosis was consistent with a benign GCT. One month postsurgery, there was evidence of recurrent tumor, and a second procedure was carried out on August 12, 1993. Combined photon and proton beam radiation were subsequently instituted from September 27, 1993, to November 12, 1993, in the Department of Radiation Oncology, Massachusetts General Hospital. A follow-up examination on August 21, 1996, revealed the patient to be well with no symptoms.

Case 3

An 11-year-old white female presented in November 1995 with frontal headaches associated with a sinus infection. Despite treatment with antibiotics, the headaches increased. She developed diplopia secondary to a left sixth nerve palsy. There was no abnormality in the visual fields. The remaining clinical examination, including neurological examination, was normal. The past medical history was noncontributory. A CT examination on December 7, 1995, revealed an expansile mass in the basisphenoid with erosion of the sella and extension into the sphenoid sinus (Fig. 3A, B). An MRI examination performed 2 days after the CT study demonstrated the large mass with low signal intensity on the T₁- and T₂-weighted images and marked enhancement (Fig. 3C-E). There was engulfment of the right intracavernous carotid artery with no evidence of narrowing.

On December 10, 1995, the sphenoid sinus was approached via the nasal cavity, and a subtotal resection of the tumor was accomplished. The histopathological examination revealed a benign GCT. Because of residual tumor, an extended transphenoidal approach with further removal of the tumor was carried out on March 14, 1996. The histological review at the Mayo Clinic and Massachusetts General Hospital following the second procedure was consistent with a GCT. Between April 23, 1996, and June 5, 1996, the patient underwent a course of combined photon and proton beam irradiation therapy.

Case 4

A 10-year-old female presented in August 1995 with double vision on the left secondary to sixth nerve palsy. She also complained of nausea and vomiting. Otherwise, the patient was in excellent health, and the past medical history was noncontributory. A CT scan done on August 23, 1995, demonstrated a moderately enhancing tumor arising within the basisphenoid with extension into the sphenoid sinus (Fig. 4A-C). The CT

Figure 1. Giant cell tumor of the sphenoid bone. (A) Coronal contrast CT study through the sphenoid sinus and adjacent clivus demonstrates a homogeneous, expansile mass within the clivus and sphenoid sinus. There is lateral extension with displacement and compression of the left and right cavernous sinuses. There is superior extension into the suprasellar cistern with superior displacement of the hypophysis, which is not delineated. There is inferior extension into the roof of the nasopharynx. The mass is bordered inferiorly by a thin, bony rim consistent with an attenuated floor of the sphenoid sinus–basisphenoid (arrows). There is slight enhancement of the tumor. Note destroyed bone at the lateral margin of the tumor. (B) Sagittal T₁-weighted image through the clivus demonstrates a homogeneous, low intensity mass involving the entire clivus with extension into the preoptine cistern. There is erosion of the sella turcica with superior displacement of the hypophysis (arrows). There is inferior extension into the nasopharynx and anterior extension into the sphenoid sinus and posterior third of the ethmoid sinuses. (C) Coronal T₁-weighted image through the mass in the sphenoid sinus demonstrates the previously described extension and low, homogeneous signal intensity of the tumor. (D) Coronal T₁-weighted image through the mass after Gd introduction demonstrates marked, heterogeneous enhancement of the tumor. (E) Axial T₂-weighted image through the center of the mass demonstrates low signal intensity. There is no evidence of tumor necrosis. Note the extension of this mass into the posterior ethmoid sinuses. The carotid arteries are displaced laterally but are not encased (arrows). There is bulging of the tumor into the preoptine cistern.

Figure 2. Giant-cell tumor of the sphenoid bone. **(A)** Axial CT study (bone window setting) through the clivus demonstrates lytic destruction of the clivus, petrous apex on the left, medial aspect of the middle cranial fossa, and the sphenoid sinus. Higher sections (not shown) demonstrates partial erosion of the sella turcica. **(B)** Axial T₁-weighted image demonstrates a large, low intensity mass corresponding to the large, lytic, bony defect. The carotid arteries are displaced laterally and posteriorly (arrows). The left carotid artery is partially surrounded by the tumor. On sagittal images (not shown), the mass bulged into the upper nasopharynx. **(C)** Axial T₁-weighted image (after Gd introduction) shows marked enhancement of the tumor. Again, note tumor surrounding the laterally displaced carotid arteries with engulfment of the left intracavernous carotid artery (arrow). **(D)** Coronal T₁-weighted image through the clivus shows the enhanced tumor bulging into the nasopharynx and cavernous sinuses with extension into the middle cranial fossa on the left. Note displacement of the left temporal lobe. **(E)** Axial T₂-weighted image through the clivus and sphenoid sinus reveals the tumor mass to be of low signal intensity. Note again tumor partially surrounding the displaced, left carotid artery (arrow).

Figure 3. Giant cell tumor of the sphenoid bone. (A) Axial contrast CT study (soft tissue window setting) through the sphenoid sinus and clivus demonstrates a homogeneous, moderately enhancing tumor mass. (B) Axial CT section (bone window setting) demonstrates partial erosion of the sphenoid sinus and adjacent clivus. (C) Sagittal T_1 -weighted image through the clivus demonstrates the low intensity tumor with erosion of the basisphenoid and extension into the pre-pontine cistern, sella turcica, and posterior sphenoid sinus. (D) Axial T_1 -weighted image after Gd introduction and fat suppression through the mid portion of the tumor reveals marked enhancement of the tumor. Note low signal intensity fluid in the anterior right sphenoid sinus (*asterisk*). There is lateral displacement of both intracavernous carotid arteries. (E) Axial T_2 -weighted image reveals low signal intensity of the tumor. Note high signal intensity fluid collection in the right sphenoid sinus.

Figure 4. Giant-cell tumor of the sphenoid bone. (A) Axial contrast CT study (soft-tissue window setting) demonstrates a moderately enhancing mass in the region of the basisphenoid, sphenoid sinus, and sellar region with extension into the prepontine cistern. The mass is homogeneous and bulges laterally into the cavernous sinuses. There is partial erosion of the sphenoid sinus and adjacent clivus. (B) Axial CT section through the sphenoid sinus and clivus (bone window setting) reveals partial erosion of the clivus and adjacent sphenoid sinus. (C) Coronal contrast CT study through the sphenoid sinus and clivus demonstrates a homogeneous, well-defined mass, extending into the suprasellar cistern and left and right cavernous sinuses, which are displaced laterally. (D) Sagittal T_1 -weighted image (after Gd introduction) reveals heterogeneous, marked enhancement of the tumor of the clivus.

scan was supplemented with an MRI scan on the same day, which reconfirmed the presence of a large tumor mass involving the clivus. The tumor revealed low signal intensity on T_1 - and T_2 -weighted images with moderate heterogeneous enhancement (Fig. 4D).

On September 1, 1995, she underwent surgical exploration with a transphenoidal approach. A firm, tan, relatively avascular tumor was found that invaded the dura. The pathological examination demonstrated a be-

nign GCT. The postoperative recovery was uncomplicated, and the diplopia resolved completely. An MRI study performed on October 3, 1995, and a CT study performed on October 18, 1995, revealed residual disease. This prompted a second surgical procedure with further resection of the tumor on October 27, 1995. A repeat MRI study on October 28, 1995, revealed some residual tumor. She underwent a combined course of photon and proton beam therapy from December 12,

1995, to January 27, 1996. On the last follow-up examination at the end of August 1996, the patient was doing well with no residual symptoms.

RESULTS

This article describes three female patients and one male patient, aged 10, 11, 15, and 16 years. The symptoms consisted of headache, predominantly in the frontal region (four patients), numbness in the V1 distribution (one patient), diplopia (three patients), and lid lag (one patient). The general neurological evaluations of these patients revealed no abnormalities. On pathological evaluation, these tumors, on gross inspection, were soft, red-brown in discoloration, and bled easily. On the histopathological examination, the characteristic findings were osteoclastic giant cells evenly distributed throughout the tumor tissue. The radiological examinations demonstrated a bulky tumor mass extending into the sella turcica, sphenoid sinus, cavernous sinuses, and adjacent middle cranial fossae. The intracavernous carotid arteries were displaced laterally, and in one patient the artery was engulfed by the tumor with no evidence of narrowing, however.

On CT study, the tumors were isodense with muscle and revealed slight to moderate enhancement. On MRI examination, all tumors revealed low signal intensity on the T_1 - and T_2 -weighted images. There was moderate enhancement after introduction of Gd.

DISCUSSION

Clinical and Pathological Evaluations

GCT of the sphenoid bone is usually detected when the lesion is large and has expanded into the adjacent anatomical structures. The most common finding is intermittent headaches, which were present in all four cases.^{23,32,43,44} These headaches were persistent and increased in severity over time. The extradural location of GCT with stretching of the dura is probably the cause for this clinical finding.

The expansile nature of GCT to the cavernous sinuses is responsible for the frequent involvement of the third and sixth cranial nerves. As in other skull base lesions of various etiologies, the sixth nerve was most commonly involved on ophthalmological examination, reflected by diplopia in three patients. The sixth nerve courses along the basisphenoid and through the cavernous sinus and is vulnerable to compressive lesions arising from the sphenoid bone, including the cavernous sinuses. The third cranial nerve is less commonly involved and usually follows involvement of the sixth nerve. Encroachment on the third nerve is characterized by partial ophthalmoplegia and ptosis, which was found

in one of our cases (case 2). Other ophthalmological findings consist of decreased visual acuity,⁴⁵ afferent pupillary defect secondary to compression on the optic nerves,⁴³ visual field defects,⁴⁶ ptosis,⁴⁷ and papilledema.⁴⁸ Occasionally, there may be slight proptosis, if the lesion extends into the posterior orbit.⁴⁹ None of these findings were present in our series. Concomitantly, there may be involvement of the fifth nerve (V1 and V2 branches)⁵⁰ with pain, paresthesia, or hypesthesia in the respective areas.⁴³ Not uncommonly, the tumor destroys the sella turcica with superior displacement of the hypophysis. Endocrine disorders may be a sequence of this expansion,⁵¹ as was found in one of our cases when amenorrhea developed. This, however, occurred two years after surgery and irradiation, and therefore may be related to these posttreatment events.

Other nonspecific symptoms that have been mentioned in various cases reports are changes in cognitive state,⁴³ olfactory hallucinations, grand mal seizures, zygomatic swelling, and pain on mastication. Some of the patients also suffered from nausea and vomiting, as was noted in two of the patients in this report.³⁶

On gross pathological examination, the tumor tissue is soft, red-brown, friable, and bleeds easily, which is a reflection of its increased vascularity. On cross section, some areas of yellowish discoloration may be present, which is secondary to lipid accumulation in histiocytes.²⁷

On histopathological examination, the tumor is composed of osteoclastic giant cells that are evenly distributed throughout the tumor tissue. The giant cells contain a variable amount of nuclei with an average of about 10 to 30 nuclei in most cases. The nuclei are round to oval in shape, hyperchromatic, and reveal no atypicality or nuclear pleomorphism. Scattered mitoses may be found in the tumor matrix. The bulk of the tumor is made up of fibroblastic spindle and round cells, which are short and plump, and nuclei that are identical to those seen within the giant cells.⁵² The spindle cells have abundant eosinophilic cytoplasm and produce collagen.⁵³ Hemorrhage and hemosiderin deposition are found, but are less prominent compared with the giant cell granuloma. Osteoid formation may occur and has been reported in 30% of cases.⁵² Malignancy in a GCT is difficult to assess on histopathological examination. Benign-appearing tumors on histological studies have metastasized, especially to the lungs and occasionally to the bone.⁵⁴ Malignant transformation, reflected by increased mitotic activity and atypicality of nuclei, has been reported in a small percentage of cases following radiation therapy.⁵⁵

CT and MRI Evaluations

On CT and MRI, GCT of the skull base arising from the sphenoid bone (most often basisphenoid) man-

ifests as a bulky tumor mass with lytic destruction, as was demonstrated in the reported four cases. Because of their slowly progressive expansion, a bony rim from the site of origin is often visible at the margin of the tumor (Fig. 1A, C). The bony shell shows frequent dehiscences, which is a reflection of aggressive growth. These bone changes are optimally depicted on CT. Because of the proximity of the basisphenoid to the sphenoid sinus, the GCT preferentially expands the sphenoid sinus, as was noted in our four cases (Figs. 1 to 4). Additionally, the sella was eroded in all cases, with superior displacement of the hypophysis (Fig. 1). Lateral expansion leads to encroachment on the cavernous sinuses, with lateral displacement of the cavernous carotid arteries. Infrequently, the carotid arteries become engulfed in the tumor mass, as occurred in one case (Figs. 2B–E).⁵⁶ Significant growth may produce almost complete replacement of the clivus by the lesion reflected by a variable-sized lytic defect (Figs. 2A, 3B, 4B). Lateral expansion from the clivus proceeds into the middle cranial fossa extradurally with displacement of the temporal lobes (Fig. 2B–E). The petrous apex is in close proximity to the clivus, and larger tumors cause erosion of the petrous apices and, in advanced cases, may proceed to the tympanic cavity and internal auditory canal with consequent symptoms of hearing loss, tinnitus, and facial nerve dysfunction (Fig. 2A). Frequently, there is encroachment on the optic canal, with bony erosion being responsible for visual impairment. Encroachment upon the cavernous sinuses is the cause for impairment in the function of the third, fourth, and sixth cranial nerves (Figs. 1 to 4). Invasion of the apex of the orbit with associated lytic destruction of the greater sphenoid wing may be responsible for a slight degree of exophthalmos encountered in some of the patients reported in the literature. Cystic degeneration is not a common feature in GCT of the skull base and was not demonstrated in our series of four cases. However, in other parts of the skeleton (axial and appendicular), GCT has been reported with large cystic areas.^{57,58} New bone formation or osteoid formation is not a prominent feature in GCT and was not visualized in our four cases.⁵⁸

Because of the greater contrast resolution, MRI is the preferred imaging modality to define the extent of the soft tissue component of the tumor. On MRI, the tumors were homogeneous and low in signal intensity on T₁- and T₂-weighted images. Moreover, characterization of the lesion on T₂-weighted images provides additional diagnostic information. The low signal intensity on the T₂-weighted images is not specific, but is suggestive of a very cellular lesion, including giant cell tumor. Low signal intensity, however, may also occur in other very cellular lesions, such as giant cell granuloma, lymphoma, meningioma, Ewing's sarcoma, and Primary Neuroectoderm Tumor (PNET).

All four lesions revealed moderate to marked, homogeneous enhancement after the introduction of Gd

(Figs. 1 to 4). This is again confirmatory evidence of the increased vascularity of these lesions and homogeneity of the tumor tissue on macroscopic and histological examination. The carotid arteries were displaced, but in one case the carotid artery was encased by the tumor with, however, no narrowing (Fig. 2B, C). Fluid levels have been described in GCT but were not identified in our four cases.⁵⁹

DIFFERENTIAL DIAGNOSIS

Giant-Cell Reparative Granuloma

The most common tumor-like condition that simulates a GCT is the giant-cell reparative granuloma. Giant-cell granulomas involving the sphenoid bone are uncommon and cannot be differentiated from GCT by CT or MRI examination. The lesions are bulky, cause lytic bone destruction, and have sharply defined margins with no sclerotic reaction.⁶⁰ They may display variable-sized, fluid-filled cavities containing blood degradation products. On CT and MRI, this is manifested by fluid fluid levels scattered throughout the lesion. They enhance slight to moderately following introduction of contrast material on CT or MRI. Because of their propensity for new bone formation, bone densities may be scattered throughout the lesion.⁶¹ This is in contrast to GCT, where bone is uncommonly demonstrated by radiological methods. The MRI signal intensities on T₁-weighted images are low and may also be low on T₂-weighted images, because of the high cellular content. Fluid levels are easily demonstrated, and the fluid content of these cavities is reflected by high signal intensity of the supernatant fluid on T₂-weighted images and low signal intensity of the blood degradation sediment.⁶²

Brown Tumor

The Brown tumor is associated with primary or secondary hyperparathyroidism. Laboratory values (high calcium, low phosphorus, increased parathyroid hormone and alkaline phosphatase) are important parameters in the diagnosis of Brown tumor. The lesion occurs predominantly in middle-aged patients (30 to 50 years), but can be found in any age, and there is a higher incidence in females.^{63,64} In the head and neck area, the lesion has been described in the mandible, maxilla,^{65–68} orbit,⁶³ base of skull, sinuses,^{68,69} predominantly frontal and ethmoid sinuses,^{70,71} and temporal bones. Frequently, there is hemorrhage within the tumor with hemosiderin within the giant cells. Cystic degeneration may occur within the tumor matrix, and consequently fluid fluid levels may be encountered on radiological examination.^{63,68} The CT and MRI appearances are similar to those of the giant-cell granuloma.²³

Aneurysmal Bone Cysts

Aneurysmal bone cysts do not form a separate entity but evolve from other lesions, such as GCT, giant cell granuloma,⁷² osteoblastoma, and chondroblastoma.^{73,74} Aneurysmal bone cysts have been described in the mandible, maxilla, base of skull, orbits,⁷⁵ and temporal bones.⁷⁶ They are composed of large, fluid-filled cystic spaces that cause marked, often eccentric expansion of bone. On the CT examination, a large cavitory lesion with occasional fluid fluid levels is encountered that is often surrounded by a dehiscent bony shell. On MRI, multiple cystic spaces may reveal fluid levels, which are low on T₁-weighted images, but reveal increased signal intensities of the supernatant fluid.^{52,73,76}

Langerhans Cell Histiocytosis

Among Langerhans cell histiocytosis, the eosinophilic granuloma is the most common type and represents a benign, solitary, or multiple tumor-like disorder of children or young adults. About half of the cases occur in the first decade of life.⁷⁷⁻⁷⁹ The most common sites of involvement in the skull occur in the skull table, temporal bones and orbits,⁸⁰ sphenoid bone,⁸¹ and base of skull.⁸² In the skull, including orbits and temporal bones, they are manifested by lytic lesions with irregular, but fairly well-defined margins. A variable amount of soft tissue densities are encountered within the lesion, which may bulge into the adjacent anatomical structures. The findings on CT and MRI⁸³ are nonspecific⁸³⁻⁸⁵ and frequently cannot be differentiated from GCT or other giant cell lesions. However, GCT is more expansile, reflected by a bony rim at the tumor margin.

Chordoma

Chordoma is derived from notochordal rests in the skull base, including the sphenoid bone and basiocciput. It is a rare tumor in the skull base in the pediatric age group.⁸⁶ The tumor causes an irregular, lytic defect similar to the GCT. The epicenter of the tumor is within the basisphenoid bone and/or basiocciput.

On CT, there is lytic bone destruction, which may be irregular or, in smaller lesions, well demarcated. In larger lesions, destroyed bone particles are scattered throughout the tumor matrix.⁸⁷ On MRI, the lesion is often heterogeneous, composed of low signal intensity on the T₁-weighted images and high signal intensity on the T₂-weighted images. Interspersed dark signal intensities of bone caused by remnants of destroyed bone are present within the lesion. There is marked enhancement after contrast material introduction.⁸⁷

REFERENCES

1. Dahlin DC: Bone Tumors: General Aspects and Analysis of 3987 Cases, ed 2. Springfield, IL: Charles C. Thomas, 1967. pp 78-89
2. Dahlin DC, Cupps RE, Johnson EW Jr: Giant cell tumor. A study of 195 cases. *Cancer* 25:1061-1070, 1970
3. Dahlin DC: Giant cell tumor of bone: Highlights of 407 cases. *AJR* 144:955-960, 1985
4. Silvers AR, Som PM, Brandwein M, Chong JL-M, Shah D: The role of imaging in the diagnosis of giant cell tumor of the skull base. *AJNR* 17:1392-1395, 1996
5. Findlay JM, Chiasson D, Hudson AR, Chui M: Giant cell tumor of the middle cranial fossa. *J Neurosurg* 66:924-928, 1987
6. Henderson BT, Whitwell H: Giant cell tumor of the skull: Case report. *Neurosurgery* 23:120-122, 1988
7. Berges O, Illouz C, Laudenbach P, et al: A giant cell tumor of the mandibular condyle. *Ann Radiol* 25:172-174, 1982
8. Chuong R, Kaban LB, Kozakewich H, et al: Central giant cell lesions of the jaws: A clinicopathologic study. *J Oral Maxillofac Surg* 44:708-713, 1986
9. Sentilkes C, Michaud J: Giant cell lesion of the maxilla. The diagnostic difficulties. *Rev Stomatol Clin Maxillofac* 87:102-107, 1986
10. Andersen L, Iejerskov O, Philipsen HP: Oral giant cell granuloma. A clinical and histological study of 129 new cases. *Acta Pathol Microbiol Scand* 87:606-616, 1973
11. Carmody RF, Rickles DJ, Johnson SF: Giant cell tumor of the sphenoid bone. *JCAT* 7:370-373, 1983
12. Kioumehri F, Rooholamini SA, Yaghmai I, Hoover L, Arnold A, Hannah J, Rodriguez L: Giant cell tumor of the sphenoid bone: Case report and review of the literature. *J Can Assoc Radiol* 41:155-157, 1990
13. Burnam JA, Benson J, Cohen I: Giant cell tumor of the ethmoid sinuses: Diagnostic dilemma. *Laryngoscope* 89:1415-1424, 1979
14. Rosen MP: Giant cell tumor of the temporal bone. *AJR* 156:1290-1291, 1991
15. Kinney SE, Sebek BA: Rare tumors of the skull base and temporal bone. *Am J Otol* (November suppl):135-142, 1985
16. Motomochi M, Handa Y, Makita Y: Giant cell tumor of the skull. *Surg Neurol* 23:25-30, 1985
17. Cook HF, Miller R, Yamada R: Giant cell tumor of infratemporal fossa: Report of a case. *J Oral Maxillofac Surg* 44:651-656, 1986
18. Kowalska B: A case of giant cell tumor of the temporal bone and the region of the temporomandibular joint. *Otolaryngol Pol* 43:415-419, 1989
19. Bessede JP, Moreau JJ, Sauvage JP, et al: Giant cell tumor of the temporal bone: Clinical study of a case. Histologic and therapeutic remarks. *Ann Otolaryngol Cerv Fac* 106:201-215, 1989
20. Glasscock ME, Hunt WE: Giant cell tumor of the sphenoid and temporal bones. *Laryngoscope* 84:1181-1187, 1974
21. Colclasure JB, Shea CM, Graham SS: Giant cell lesions of the temporal bone. *Am J Otol* 2:188-192, 1981
22. Rock JP, Mahmood A, Cramer HB: Giant cell tumor of the skull base. *Am J Otol* 15:268-272, 1994
23. Bertoni F, Unni KK, Beabout JW, Ebersold MJ: Giant cell tumor of the skull. *Cancer* 70:1124-1132, 1992
24. Hug EB, Fitzek MM, Liebsch NJ, Munzenrider JE: Locally challenging osteo- and chondrogenic tumors of the axial skeleton: Results of combined proton and photon radiation therapy using three-dimensional treatment planning. *Int J Radiat Oncol Biol Phys* 31:467-476, 1995
25. Reed L, Willison CD, Schochet SS, Voelker JL: Giant cell tumor of the calvaria in a child. *J Neurosurg* 80:148-151, 1994
26. Halpin SF, Britton JA, Uttley D: Giant cell tumor of the skull base: MR appearances. Case report. *Neuroradiology* 34:526-527, 1992
27. Vernet O, Ducrey N, Deruaz J-P, de Tribolet N: Giant cell tumor of the orbit. *Neurosurg* 32:848-851, 1993
28. Malone S, O'Sullivan B, Catton C, Bell R, Fornasier V, Davis A: Long-term follow-up of efficacy and safety of megavoltage radiotherapy in high-risk giant cell tumors of bone. *Int J Radiat Oncol Biol Phys* 33:689-694, 1995

29. Muszynski P, Modrzejewski M, Kotarba E: Osteoclastoma of temporal and zygomatic bone. *Otolaryngol Pol* 49:324-328, 1995
30. Saleh EA, Taibah AK, Naguib M, Aristegui M, Vassallo G, Landolfi M, Sanna M: Giant cell tumor of the lateral skull base: A case report. *Otol Head Neck Surg* 111:314-318, 1994
31. Pradhan S, Datta NR, Krishnani N, Ayyagari S, Tandon P: Giant cell tumor of the petrous bone. *Indian J Cancer* 28:177-180, 1991
32. Uttley D, Archer DJ: Giant cell tumor of the sphenoid sinus: An unusual skull base tumor. *J Laryngol Otol* 105:855-857, 1991
33. Curilovic A, Eich GF, Stallmach T: Giant cell tumor in the skull of a 9 year old child: Immunohistochemistry to confirm a diagnosis rare for age and site. *Pediatr Pathol Lab Med* 15:769-779, 1995
34. McCluggage WG, McBride GB, Primrose WJ, Cullan J, McNaboe EJ, Bharucha H, Fannin T: Giant cell tumor of the temporal bone presenting as vertigo. *J Laryngol Otol* 109:538-541, 1995
35. do Amaral CM, Julio GL, Cardoso LA, Bueno MA: Craniofacial treatment of giant cell tumors of the sphenoid bone. *J Craniofac Surg* 5:254-256, 1994
36. Ohaeghalam SC, Gupta IM: Giant cell tumor of the sphenoid bone with clival extension. *J Neurol Neurosurg Psychiatry* 40:790-794, 1977
37. Herman SD, Mesgarzadeh M, Bonakdarpour A, Dalinka MK: The role of magnetic resonance imaging in giant cell tumor of bone. *Skel Radiol* 16:635-643, 1987
38. Rogers LF, Mikhael M, Christi M: Case report 276. *Skel Radiol* 12:48-53, 1984
39. Schmidt RG, Kabbani YM, Mayer DP: Magnetic resonance imaging of giant cell tumor of bone. *J Am Podiatr Med Assoc* 83:634-636, 1993
40. Schutte HE, Taconis WK: Giant cell tumor in children and adolescents. *Skel Rad* 22:173-176, 1993
41. Aoki J, Moriya KM, Yamashita K, Fujioka F, Ishii K, Karakida O, Imai S, Sakai F, Imai Y, Sone S: Giant cell tumors of bone containing large amounts of hemosiderin: MR-pathologic correlation. *JCAT* 15:1024-1027, 1991
42. Harman SA, Pribram H: Case report 586. *Skel Radiol* 19:212-215, 1990
43. Wolfe JT, Schetthauer BW, Dahlin DC: Giant-cell tumor of the sphenoid bone. *J Neurosurg* 59:322-327, 1983
44. Watkins LD, Uttley D, Archer DJ, Wilkins P, Plowman N: Giant cell tumors of the sphenoid bone. *Neurosurg* 30:576-581, 1992
45. Doshi R, Chaudhari AB, Thompson G: Giant cell tumor of the sphenoid bone. *J Can Sci Neurol* 4:213-216, 1977
46. Ramamurthi B, Visvanathan GS, Plaii KM: Osteoblastoma of the skull. *J Neurosurg* 12:287-290, 1955
47. Geissinger JD, Siqueira FB, Ross ER: Giant cell tumor of the sphenoid bone. *J Neurosurg* 32:665-670, 1975
48. McNerney JC: Giant cell tumor of the base of the skull. *J Neurosurg* 169-174, 1949
49. Michael LA: Giant cell tumor of the sinuses. *Laryngoscope* 69:320-328, 1959
50. Potter GD, McClellan BL: Malignant giant cell tumor of the sphenoid bone and its differential diagnosis. *Cancer* 25:167-170, 1970
51. Kemeny AA, Afshar F: Giant cell tumor of the sphenoid bone with coincidental galactorrhoea—A case report. *Br J Neurosurg* 4:343-346, 1990
52. Huvos AG: Giant cell tumor of bone, in *Bone Tumors: Diagnosis, Treatment, and Prognosis*, ed 2. Philadelphia, WB Saunders, 1991, pp 425-467
53. Huvos AG: Giant cell tumor of the craniofacial bones: Giant cell "reparative" granuloma of jaw bones; cherubism, in *Bone Tumors: Diagnosis, Treatment, and Prognosis*, ed 2. Philadelphia, WB Saunders, 1991, pp 469-475
54. Rock MP, Pritchard DJ, Unni KK: Metastases from histologically benign giant cell tumor of bone. *J Bone Joint Surg (Am)* 66:269-274, 1984
55. Martins AN, Dean DF: Giant cell tumor of sphenoid bone. Malignant transformation following radiotherapy. *Surg Neurol* 2:105-107, 1974
56. Perria L, Viale GL, Gentile SL: Riesen Zell tumor (osteoklastome) der sellaegens. *Acta Neurochir* 19:183-197, 1968
57. Resnick D: *Diagnosis of Bone and Joint Disorders*, ed 3. Philadelphia, WB Saunders, 1995, pp 3785-3806
58. Epstein N, Whelan M, Reede D, Slobadan A: Giant cell tumor of the skull: A report of two cases. *Neurosurgery* 11:263-267, 1982
59. Kaplan PA, Murphey M, Greenway G, Resnick D, Sartoris DG, Harms S: Fluid-fluid levels in giant cell tumors of bone: Report of two cases. *JCAT* 11:151-155, 1987
60. Tallan EM, Olsen KD, McCaffrey TV, Unni KK, Lund BA: Advanced giant cell granuloma: A twenty-year study. *Otolaryngol Head Neck Surg* 110:413-418, 1994
61. Rhea JT, Weber AL: Giant cell granuloma of the sinuses. *Radiology* 147:135-137, 1983
62. Wiatrak BJ, Bluckman JL, Fabian RL, Wessler TA: Giant cell reparative granuloma of the ethmoid sinus. *Otolaryngol Head Neck Surg* 97:504-509, 1987
63. Parrish CM, O'Day DM: Brown tumor of the orbit. *Arch Ophthalmol* 104:1199-1202, 1986
64. Naiman J, Green WR, D'Huerle, Iliff WJ, Benedict W, Eggleston JC, Dorfman HD: Brown tumor of the orbit associated with primary hyperparathyroidism. *Am J Ophthalmol* 90:565-571, 1980
65. Shanmugham MS, Alhady SF: Hyperparathyroidism with osteitis fibrosa cystica in the maxilla. *J Laryngol Otol* 98:417-420, 1984
66. Friedman WH, Pervez N, Schwartz AE: Brown tumor of the maxilla in secondary hyperparathyroidism. *Arch Otolaryngol* 100:157-159, 1974
67. Smith PG, Marrogi AJ, Delfino JJ: Multifocal central giant cell lesions of the maxillofacial skeleton. *J Oral Maxillofac Surg* 48:300-305, 1990
68. Gelman R, Gelland FE: Brown tumor of the facial bones. *AJNR* 12:1179-1181, 1991
69. Schweitzer VG, Thompson NW, McClatchey KD: Sphenoid sinus Brown tumor, hypercalcemia and blindness: An unusual presentation of primary hyperparathyroidism. *Head Neck Surg* 8:379-386, 1986
70. Siendenbiedel H: Brauner tumor der orbita. *Klin Monatsbl Augenheilkd* 122:86, 1953
71. Ferry AP: Brown tumors (fibro-osseous bone replacement and overgrowth) of the orbit in hyperparathyroidism. *Metab Pediatr Ophthalmol* 3:67, 1979
72. Oda Y, Tsuneyoshi M, Shinohara N: "Solid" variant of aneurysmal bone cyst (extragnathic giant cell reparative granuloma) in the axial skeletal and long bones. A study of its morphologic spectrum and distinction from allied giant cell lesions. *Cancer* 70:2642-2649, 1992
73. Damjanovic J, Linder J: *Anderson's Pathology*. St. Louis, CV Mosby, 1996
74. Spjut HJ, Ayala AG: Skeletal tumors in children and adolescents. *Hum Pathol* 14:628-642, 1983
75. Yee RD, Gogan DG, Thorp TR, Schut L: Optic nerve compression due to aneurysmal bone cyst. *Arch Ophthalmol* 95:2176-2179, 1977
76. Vergel de Dios AM, Bond JD, Shives TC, McLeo RA, Unnik K: Aneurysmal bone cyst: A clinicopathologic study of 238 cases. *Cancer* 65:29:21-31, 1992
77. Krenghi M, Boccardi A, Gandini G, Negri G: Report of a rare case of Langerhans-cell histiocytosis of the skull base. *Minerva Med* 87:99-103, 1996
78. Goldsmith AJ, Myssiorek D, Balderrama E, Patel M: Unifocal Langerhans's cell histiocytosis (eosinophilic granuloma) of the petrous apex. *Arch Otolaryngol Head Neck Surg* 119:113-116, 1993
79. Bonafé A, Joomy H, Jaeger P, Frayssé B, Manelfe C: Histiocytosis X of the petrous bone in the adult: MRI. *Neuroradiology* 36:330-333, 1994
80. Erly WK, Carmody RF, Dryden RM: Orbital histiocytosis X. *AJNR* 16:1258-1261, 1995
81. Stromberg JS, Wang AM, Huang TE, Vicini FA, Nowak PA: Langerhans cell histiocytosis involving the sphenoid sinus and superior orbital fissure. *AJNR* 16:964-967, 1995
82. Boccardi A, Krenghi M, Gandini G: An unusual case of Langerhans-cell histiocytosis of the base of the skull: The computed tomographic and magnetic resonance aspects. *Radiol Med* 87:872-874, 1994

83. Beltran J, Aparisi F, Bonmati LM, Rosenberg ZS, Present D, Steiner GC: Eosinophilic granuloma: MRI manifestations. *Skel Radiol* 22:157–161, 1993
84. Davies AM, Pikoulas C, Griffith J: MRI of eosinophilic granuloma. *Eur J Radiol* 18:205–209, 1994
85. Hermans R, DeFoer B, Smet MH, Leysen J, Feenstra L, Fossion E, Baert AL: Eosinophilic granuloma of the head and neck: CT and MRI features in three cases. *Pediatr Radiol* 24:33–36, 1994
86. Benk V, Liebsch NJ, Munzenrider JE, Efird J, McNanus P, Suit H: Base of skull and cervical spine chordomas in children treated by high-dose irradiation. *Int J Radiol Oncol Biol Phys* 31:577–581, 1995
87. Weber AL, Liebsch N, Sanchez R, Sweriduk S: Chordomas of the skull base. *Neuroimaging Clin North Am* 4:515–527, 1994