Management of Zone III Missile Injuries Involving the Carotid Artery and Cranial Nerves

Abstract—Carotid and cranial nerve injuries from zone III (high cervical/cranial base) missile injuries are rare and difficult to treat. We have treated five patients with such injuries. We present our management scheme, and compare it to the management of the same injuries in other reports. Five consecutive zone III missile injuries presented to our institution. Trauma assessment by the trauma team, followed by detailed neurological assessment and radiographs (angiogram and computed tomography) were obtained on admission. All patients presented with dysphagia and carotid artery injury with good collateral flow, documented by angiogram. Two patients had facial nerve injury, one had trigeminal nerve injury, one patient presented with tongue weakness, and one patient suffered conductive hearing loss. No patient had evidence of stroke clinically or radiographically. Carotid artery injury was managed with bypass (3 of 5) or ligation (2 of 5). Cranial nerve injuries were documented and treated aggressively with surgery if needed. All patients were discharged to home. Patients presenting with zone III missile injuries should receive an expeditious neurological exam and four-vessel angiogram after initial trauma survey and resuscitation. Bypass of the injured portion of carotid artery is a valid treatment in the hemodynamically stable patient. The unstable patient should undergo ligation to stop hemorrhage and protect against immediate risk for stroke, with the option to bypass later. Cranial nerve injuries should be pursued and aggressively treated to minimize morbidity and prevent mortality.

Gunshot wounds (GSWs) to the high neck and cranial base can present complex management problems.^{1,2} Approximately 10% of patients with penetrating injuries to the neck involve the carotid and are associated with a 20% fatality rate.¹ Injuries can include the arterial and venous structures, cranial nerves (V-XII), cranio-vertebral junction (spinal cord and vertebral elements), aerodigestive tract, and mandible. Cerebrovascular injury, immediate (hemorrhage, stroke) and delayed (embolization, venous infarction, etc.) can be life-threatening. Careful assessment and surgical planning is required to sustain life, minimize morbidity, and prevent future neurological complications. We have managed five cases of zone III (angle of the jaw to the skull base) missile injuries to the neck and present a management

schema for these injuries based on our experience and this review.

CASE REPORTS

Illustrative Case of Carotid Ligation and Cranial Nerve Management

A 24-year-old male (P.J.) sustained a GSW to the left mandible. He was intubated upon arrival, with an initial blood pressure of 50 mmHg, a heart rate of 160, and a Glasgow Coma Score (GCS) of 5. He received a standard trauma survey as per Advanced Trauma and Life Support (ATLS) protocols and radiographic evalu-

Figure 1. Lateral view, digital subtraction angiogram (DSA) of common carotid artery revealing extravasation of contrast dye at the superior to the angle of the mandible (patient P.J.).

ation of cervical spine and chest. He was noted to have excessive intra-oral bleeding. Computed tomography (CT) of the head revealed bilateral condylar fossa and mandibular fractures. There were no intracerebral abnormalities. Following a prolonged resuscitation, the patient underwent a four-vessel cerebral angiogram. The left internal and external carotid arteries were transected by the bullet and there was evidence of extravasated contrast (Fig. 1). The study revealed good collateral flow from the right internal carotid artery (ICA) via the anterior communicating artery (AComm) and from the left vertebral artery via the posterior communicating artery (PComm) (Fig. 2). The cervical segment of the ICA was managed with intraluminal balloon occlusion and the external carotid artery (ECA) was occluded with platinum coils (Fig. 3). As the patient's vital signs stabilized, he became more responsive and showed no focal signs of hemispheric dysfunction. Ipsilateral facial nerve palsy and a decreased gag reflex were also noted. The following day, the patient under-

Figure 2. (Patient P.J.) Left image: lateral view, DSA of left vertebral artery demonstrates patent Pcomm with anterograde filling of MCA. Right image: Anterior to posterior (AP) view: DSA of right internal carotid also shows good collateral filling of opposite MCA via anterior communicating artery.

Figure 3. (Patient P.J.) Left image: lateral view, DSA is of the left common carotid artery after placement of an intra-arterial balloon in the left ICA and coils that were used to occlude the left maxillary artery. The right image is a nonsubtracted study that allows visualization of the balloon, coils, and shrapnel from the bullet.

went surgery for debridement of the bullet wound, direct pharyngoscopy, immobilization of the mandible, and tracheostomy. Postoperatively the patient remained stable and had no signs of cerebral ischemia. His facial weakness required the insertion of a gold weight in the upper eyelid to protect against corneal exposure and injury. Further evaluation including electromyogram (EMG) of the facial nerve noted complete transection and the patient consented to facial nerve repair during his hospital stay. He was discharged home following removal of the tracheostomy. Twelve weeks postoperatively his facial weakness improved in the inferior divisions, but he was lost to follow-up after removal of his mandibular wires.

Illustrative Case (Extracranial to Intracranial) of EC-IC Bypass and Cranial Nerve Management

A 22-year-old male (M.F.) sustained a GSW to the left posterior neck/skull base above the angle of the mandible. He was admitted to another institution and on arrival, he was hemodynamically stable, awake, alert, and oriented with a GCS of 15. He complained of difficulty swallowing, left shoulder dysesthesias and slight, left, deltoid weakness. Examination revealed swelling in the left hypopharynx with uvular and tongue deviation to the left. Cervical spine films revealed obliteration of the left foramen transversarium at C2. Dynamic

studies revealed no cervical spine instability. CT of the head and upper cervical spine revealed no intracranial pathology, and confirmation of the C2 spine injury. The patient was electively intubated for airway protection and remained sedated for a four-vessel cerebral angiogram. The angiogram revealed left vertebral artery occlusion and a left internal carotid pseudoaneurysm (Fig. 4). The left ICA remained patent despite the pseudoaneurysm and the right vertebral injection revealed adequate cross filling (Fig. 5). The patient was transferred to our institution for management.

Repeat CT scan of the head revealed no abnormality and the patient was taken to surgery for ICA bypass using the greater saphenous vein. The ligation and bypass was performed from the proximal left ICA to the left M2 segment of the middle cerebral artery (MCA). Graft patency was confirmed by intraoperative angiography and postoperative 3-D CT angiogram (Figs. 6 and 7). The patient tolerated the procedure well and following extubation, underwent evaluation of cranial nerve function including EMG, swallowing studies and direct laryngoscopy, which revealed significant weakness of the left vocal cords and potential for aspiration. The otolaryngology service performed a left thyroplasty and placed a gastrostomy using a percutaneous-endoscopic technique (PEG). The patient was discharged home receiving bolus feeding through his PEG. In follow-up, the patient is now eating normally with minor left-sided tongue weakness. The left deltoid weakness resolved without treatment.

Figure 4. (Patient M.F.) Lateral view, DSA frames show filling of the ICA pseudoaneurysm as well as good distal runoff to the distal ICA and MCA. The two images are from the same run: the left image is prior to the right in time sequence.

METHODS: ASSESSMENT AND INTERVENTION

Patient Evaluation

All patients were adults between 19 and 26 years of age, two females and three males. All five patients were admitted to a level I trauma center. Initial assessment of the extent of injury, airway management, and hemodynamic status was performed and appropriate resuscitation instituted. Neurosurgical consultation was obtained and a thorough, expeditious baseline neurological exam established documenting any localizing signs (hemispheric or spinal cord dysfunction) and any cranial neuropathy.

Two of the five patients were considered hemodynamically unstable with sustained systolic blood pressures < 90 mmHg and tachycardia (heart rate > 100).³ One required 16 units of packed red blood cells for resuscitation (P.J.). The second patient had multiple GSWs to the chest (M.B.). The remaining three had stable vital signs. Each patient underwent a four-vessel cerebral angiogram to evaluate vascular injury and collateral flow. Once the extent of the vascular injury was determined, the patients were taken to the operating room for EC-IC bypass or surgical ligation. One patient had emergent balloon occlusion of the proximal ICA and coil occlusion of the proximal ECA in the angiogram suite. He went to the operating room the following day for debridement of the wound, distal carotid occlusion by ligation of the petrous segment, tracheostomy, and intraoperative pharyngoscopy to evaluate mucosal integrity and to place a feeding tube under direct visualization. All patients were

Figure 5. (Patient M.F.) This unsubtracted, angiographic, (AP) view of MF shows good cross filling from right vertebral artery to left PCA and MCA. The bullet can be seen through the left maxilla.

Figure 6. (Patient M.F.) Intraoperative angiograms are used to confirm patency and flow of ICA bypasses. The left image shows the distal anastomosis of the saphenous vein to the M2 segment of the MCA. The right image shows the proximal anastomosis with the ICA. The large vessel on the extreme right of the field is the bypass as it ascends under the skin to the cranial anastomosis.

Figure 7. (Patient M.F.) On the first postoperative day, a routine 3D CT angiogram is obtained to evaluate patency and anatomical relationship of the bypass to the skull. The bypass is seen extracranially, passing posterior to the patients left ear and entering the craniotomy site on the left image. The right image shows the bypass entering the skull and filling the ipsilateral MCA branches.

evaluated postoperatively with swallowing studies and direct visualization of the vocal cords (Table 1).

Surgical Technique

The surgical technique for carotid artery bypass using saphenous vein autografts has been previously de-

scribed.⁴⁻⁶ The ICA was utilized for the proximal bypass anastomosis in all cases and the M2 segment of the MCA (2 of 3), or petrous ICA (1 of 3) for the distal terminus. Ligation (2 of 5 cases), when performed was performed proximal (ICA) and distal (petrous ICA) to the injured segment. The craniotomy is a small frontotemporal exposure of the Sylvian fissure to access the

Table 1. Patient Statistics on Arrival

Patient, Age	Date	Zone of Injury	Neuro Exam on Arrival	Blood Pressure on Arrival
J.S., 19	3/11/94	III Right	Right Side: V3 numbness decreased gag Left Side: Horner's syndrome X (vocal cord paralysis)* XII Deltoid weakness decreased gag	130/65 150/70
M.F., 22	3/29/95	III Left	Left Side: VII Grade III X (vocal cord paralysis) [†] VIII (conductive hearing loss) decreased gag	85/40 multiple GSWs to Chest
M.B., 23	6/9/95	III Left	Left Side: X (vocal cord paresis) Horner's syndrome decreased gag	110/65
K.R., 26	2/10/96	III Left	Left Side: VII Grade IV* decreased gag	50/0 required multiple transfusions on angiogram table

*Received PEG and thyroplasty.

[†]Developed gastroparesis and received PEG and thyroplasty.

*Facial nerve reanastomosis and gold weight implant to left upper eyelid prior to discharge.

M2 segment of the MCA. For petrous ICA exposure, a lower craniotomy is made and the petrous temporal bone is drilled, exposing the carotid. The former approach is faster, but requires opening the dura and dissection of the Sylvian fissure. A subgaleal tunnel is made behind the ear, wide enough to easily pass a finger, connecting the cranial incision with the neck incision. During the cranial portion of the surgery, the saphenous vein is harvested from the leg. The neck dissection gives access to the bifurcation and the upper cervical ICA and ECA. We prefer to perform the distal anastomosis first, to the either MCA or petrous ICA, followed by the proximal anastomosis, using a microscope for both. Heparin is given intravenously (1000–2000 Units) just prior to performing the anastomosis. The patient is monitored continuously (brain stem evoked responses, electroencephalography, somatosensory evoked responses) during the procedure and placed under moderate hypothermia (34°C), barbiturate-induced burst suppression and induced hypertension during temporary occlusion for the bypass portion of the procedure.

Adjunctive procedures for cranial nerve palsies included thyroplasty and PEG. A thyroplasty is used to medialize the paretic vocal cord to protect the airway. This requires that the contralateral vocal cord can adduct effectively to the paretic cord. A small silicon prosthesis or autologous cartilaginous wedge is inserted under the vocalis muscle while the patient is awake. The voice is tested and the implant is shaped to “tune” the voice. Patients with poor mechanical swallowing and risk of aspiration had alimentary diversion by PEG. Facial nerve injuries, when present, are evaluated and if complete transection is identified, treated with nerve grafting or direct repair. Nonocclusive tarsorrhaphy is used to protect the cornea using a gold weight inserted in the upper eyelid of the affected eye.

RESULTS

Five patients with zone III GSW injuries to the ICA were admitted to our institution from 1994 to 1996. In two out of five cases the patients presented with hemodynamic instability and massive hemorrhage. Three other patients were hemodynamically stable upon arrival. These three patients received carotid artery bypass (3 of 5). Two unstable patients either from massive hemorrhage (P.J.) or extensive chest wounds (M.B.) had carotid occlusion or ligation, respectively (Table 2).

Lower cranial nerve deficits were present in all patients. Other neurological deficits included sympathetic (2 of 5) and spinal root (1 of 5) injuries. There were no patients in this series with signs of hemispheric ischemia (i.e., hemiparesis).

Cranial nerve injuries included trigeminal, facial, glossopharyngeal-vagal complex, and hypoglossal

Table 2. Findings and Management

Patient	Angiographic Findings	Management
J.S.	pseudoaneurysm	EC-IC bypass
M.F.	pseudoaneurysm	EC-IC bypass
M.B.	pseudoaneurysm, vertebral artery transection	Ligation
K.R.	pseudoaneurysm, A-V fistula, carotid and jugular transection	EC-IC bypass and coiling of fistula
P.J.	Transection of ICA and ECA	Ligation (Balloon occlusion and coiling)

EC-IC = extracranial-intracranial; ICA = internal carotid artery; ECA = external carotid artery; and A-V = arterio-venous.

nerves. Adjunctive surgical procedures were required for cranial neuropathy in three cases. One patient (M.B.) with gastrointestinal reflux required emergent re-intubation for respiratory distress secondary to aspiration. The second patient (M.F.) had difficulty managing oral secretions. Paralysis of the vocal cord ipsilateral to injury was identified by direct laryngoscopy in both patients. Both patients were treated with thyroplasty to medialize the vocal cord and PEG for alimentation while lower cranial function recovered. The third patient had a gold weight implant for the left upper eyelid and ipsilateral facial nerve re-anastomosis (Table 1).

DISCUSSION

Zones of the Neck

Treating individuals with penetrating injuries to the neck requires a working knowledge of the anatomy. The classical approach to cervical anatomy subdivides the neck into a series of triangles. This anatomical paradigm, however, has limitations in assessing penetrating cervical injury, thus an alternate system was developed that divides the neck into three vascular zones.

The zones are numbered I to III from the root of the neck to the skull base, respectively (Fig. 8).¹ The root of the neck to the mid-portion of the neck (hyoid bone) describes zone I. Penetrating trauma in this area may cause massive hypotension secondary to great vessel injury. The aorta, subclavian, brachiocephalic, carotid, and vertebral arteries and the jugular vein are subject to injury in this zone. Zone II spans the mid-portion of the neck (hyoid bone) to the angle of the jaw. The common carotid often bifurcates and the external carotid may ramify extensively in zone II. The jugular veins as well as the carotid arteries and its branches may be injured in this zone, although blood loss is frequently limited by tamponade from surrounding myofascial compartment in zones II and III. Zone III is defined inferiorly by the angle of the jaw and superiorly by the carotid canal at the skull base. Vascular structures in this zone include

Figure 8. This illustration shows the three zones of the neck. Zone I spans the area from the clavicle to the mid-neck (hyoid bone). Zone II stretches from the mid-neck (hyoid bone) to the angle of the jaw. Zone III includes the remainder of the neck from the angle of jaw to the skull base.

cervical, petrous, and cavernous segments of the ICA as well as terminal branches of the external carotid and the vertebral artery. The paravertebral cervical plexus, jugular vein, and bulb may be compromised by penetrating injury. The majority of patients with zone III injuries will have associated cranial nerve injuries that represent a substantial source of morbidity. Careful examination at presentation and subsequent follow-up can limit potential morbidity (e.g., aspiration pneumonia) of such cranial nerve injuries.

HISTORICAL DATA AND LITERATURE REVIEW

The first report of ICA bypass for restoration of cerebral blood flow after GSW was in 1918.⁷ The wound was accessible (in what we now refer to as zone II), and anastomosis of ICA to the external carotid was the mode of bypass. A modern review of carotid injury reports a 10% incidence of penetrating injuries to the neck. The overall mortality was as high as 47% with proximal ligation or occlusion of the carotid as treatment.⁸ Anatomical studies have shown that only 20% of the population has a complete anastomotic Circle of Willis, perhaps a contributing factor for the reported poor outcome.⁹ Since the Korean War, the management of ICA injury has evolved toward preservation of the carotid circulation, usually by direct repair or interposition grafting, where feasible, with an overall mortality of 15–20%.^{1,9–15}

In 1973, Bradley⁹ introduced the stratification of patients into revascularization versus conservative therapy based on preoperative neurological condition. His conclusion was that patients who were obtunded or comatose on initial presentation have a poor prognosis, whereas awake patients, even with hemiparesis, benefit from intervention. Reports subsequent to this paper have attempted to identify the prognostic indications for aggressive versus conservative repair in such patients.^{16–19} The more recent reports demonstrate good outcomes with revascularization in the stable patient, but a management algorithm based on hemodynamic and neurological status for patients with zone III injuries has not been developed.^{10,12–14}

Direct Repair Versus EC-IC Bypass

The technical difficulties encountered for direct repair of the internal carotid in zone III injuries can be substantial. Approaches to zone III injuries require extensive surgical exposure of the distal cervical ICA and adjacent structures in the infratemporal fossa. Distal control of the ICA is a potential problem if the injury extends to the cartilaginous ring at the entrance to the carotid canal. Mandibular osteotomy or zygomatic arch osteotomy can improve access but do not address back-bleeding from the petrous carotid artery.¹

EC-IC bypass avoids the problems of direct repair while preserving the cerebral circulation, but requires a craniotomy and exposure of either the petrous or supraclinoid ICA or the MCA as the distal recipient vessel for the bypass. The complexity of such a procedure in a trauma setting has perhaps limited its utility and acceptance. Experience with vascular lesions and skull base neoplasms involving the petrous and intracavernous internal carotid have advanced the management of these complex problems.⁵ The evolution of cranial base approaches affording surgical access to the petrous and cavernous segments of the ICA permit application of these techniques to special problems such as dissection, embolization, radiation angiopathy, and acute trauma, in addition to neoplastic and vascular disease. Although EC-IC bypass is a viable option for zone III injuries, it is not universally applicable in these critically ill patients. Based on our experience and review of the literature, we have developed a management scheme to facilitate treatment planning in these circumstances.

In addition to bypass and ligation, the use of carotid stents may play a role in carotid trauma. Although current reports only include blunt trauma, they should not be excluded from the therapeutic possibilities.²⁰ In our institution, we have used carotid stents for stenosis and dissection but have no experience with traumatic pseudoaneurysms.

TREATMENT OPTIONS IN ZONE III INJURIES

The decision to perform a bypass or simply ligate the injured vessel is made after adequate trauma assessment and management of concurrent, life-threatening injuries (Fig. 9). Initial trauma assessment follows ATLS protocols to prioritize injuries and direct initial care. After initial resuscitation, the patient with a zone III injury should have an expeditious neurological survey. Careful attention is paid to hemispheric symptoms contralateral to the injury, as well as ipsilateral cranial or peripheral nerve injuries. Radiographic studies should include the standard views of the cervical spine to evaluate bony and/or soft-tissue injury. CT of the head should include thin sections (3–5 mm) through the skull base. Cervical CT may be needed to evaluate any cervical spine injuries or soft-tissue injury of the neck. A standard four-vessel cerebral angiogram should be performed to define the vascular injury and the adequacy of collateral flow.²¹ The results of the angiogram and the neurological examination are instrumental in surgical decision making regarding carotid artery management. If available, cerebral blood flow measurements using xenon CT would provide quantified values to evaluate cerebral perfusion. These supplemental studies can be useful, but are not widely available and difficult to obtain in the trauma setting.

1. Trauma team evaluation
 - a. Standard ATLS protocols with Airway, Breathing, Circulation, etc.
 - b. Initial fluid resuscitation - stabilize
2. Neurologic Evaluation
 - define hemispheric signs
 - define cranial/peripheral nerve symptoms
3. Radiographic Analysis
 - a. Plain films of neck Anterior/Posterior, Lateral, Open Mouth
 - define projectile location
 - define bony and soft tissue injury
 - Flexion/Extension (F/E) films as needed
 - b. Computed Tomography
 - Head - include thin cuts of skull base define hematomas other CNS injuries
 - Neck - when indicated by plain films- follow-up F/E as needed
 - c. Angiography
 - define vascular injury
 - define collateral flow
4. Management Plan
 - a. carotid injury
 - b. other neurologic deficits
 - gag/cough/swallowing
 - facial movement/sensation

Figure 9. Penetrating zone III injury presents to emergency room

To simplify patient evaluation, we have defined patients with carotid artery injury as “neurologically stable” or “unstable” and “hemodynamically stable” or “unstable.” Neurological stability in the face of carotid injury (transection, fistula, and pseudoaneurysm) is defined as lack of hemispheric symptoms and radiographic evidence of good collateral flow. Neurologically unstable patients have either hemispheric symptoms or poor collateral flow in the presence of carotid artery injury (transection, fistula, and pseudoaneurysm). These latter patients frequently manifest signs of carotid ischemia or have impending ischemia. Hemodynamic stability is defined as a sustained systolic blood pressure (SBP) > 90 mmHg and heart rate < 100. The decision whether to ligate or bypass the injured ICA is based on a simple table that relates neurological signs, hemodynamic status, and radiographic data, particularly collateral flow. There are four permutations to these variables (Table 3).

Neurologically Stable and Hemodynamically Stable

As defined, this group has no compromise of cerebral blood flow nor have they lost a significant amount of blood. These patients may go on to have neurological insult due to embolization, recurrent hemorrhage, or progressive thrombosis from the injury. Under these circumstances, bypass is safe and advisable. Bypass and isolation of the injured segment of the carotid artery eliminates the risk of complications and re-establishes ipsilateral blood flow in the carotid territory.^{22–24} All three patients that underwent bypass in our series fell into this category. All were hemodynamically stable and all neurological symptoms were referable to local

Table 3. Management Schema

	<i>Hemodynamically Stable</i>	<i>Hemodynamically Unstable</i>
Neurologically stable	Bypass	Ligate
Neurologically unstable	Bypass	Ligate*

“Neurologically Stable” is defined as no hemispheric symptoms and radiographic evidence of good collateral flow with ipsilateral carotid injury (transection, fistula, pseudoaneurysm or traumatic dissection).

“Neurologically Unstable” is defined as any hemispheric symptoms or poor collateral flow in the presence of carotid injury (transection, fistula, pseudoaneurysm, or traumatic dissection).

“Hemodynamically Stable” is defined as (SBP > 90 mmHg) and HR < 100 bpm > 50 bpm with a strong pulse.

“Hemodynamically Unstable” is defined as sustained hypotension (SBP < 90 mmHg) and HR > 100 bpm.

*Confers the worst prognosis from literature.

trauma rather than hemispheric ischemia. All of our patients had pseudoaneurysms, putting them at risk for rupture or embolization and stroke.

Neurologically Unstable and Hemodynamically Stable

Patients with either hemispheric symptoms or poor collateral flow with concomitant carotid artery injury are considered neurologically unstable. These patients are amenable to bypass if hemodynamically stable. Carotid artery bypass eliminates the risk of continued cerebral injury, whether embolic or ischemic.

Neurologically Stable and Hemodynamically Unstable

Neurologically stable patients who have excessive blood loss from the carotid or associated injuries need rapid stabilization to prevent hypovolemic shock. These patients presumably have adequate cerebral perfusion from patent collaterals yet cannot tolerate lengthy surgical procedures until stabilized. In such circumstances, acute ligation distal and proximal to the site of injury is advocated. This form of ligation prevents embolic phenomena distally and isolates the injured segment with proximal occlusion.

The two patients who had acute carotid ligation in our group were representative of this category. Both had good collateral flow on angiography and were hemodynamically unstable. One patient (P.J.) with ICA and external carotid transection required massive transfusion during resuscitation and angiography. Acute surgical intervention was deemed too risky and he underwent angiographic proximal occlusion. He had surgical debridement of the wound and distal ligation of the ICA the following day. The other patient (M.B.) treated with ligation, had multiple chest wounds with hemothorax. Although this second patient also had vertebral artery transection and occlusion, collateral flow from the opposite carotid and vertebral artery was adequate to prevent ischemic injury.

Neurologically Unstable and Hemodynamically Unstable

Patients who have either neurological deficit or poor collateral flow and are hemodynamically unstable are at the highest risk for morbidity. ATLS protocols have established the importance of correcting the hypovolemic state prior to organ-specific treatment. In these

circumstances, ligation of the ICA by intraluminal balloon or direct surgical exposure is recommended. Such patients may develop cerebral infarction, although with aggressive resuscitation collateral flow might prove adequate to avoid or limit cerebral ischemia. Although we advocate ligation as initial management in these acute situations, restoration of the carotid circulation after stabilization is preferable. Patients should be re-evaluated with angiography and considered for bypass procedures if appropriate.

This algorithm is primarily based on our experience with carotid artery bypass for neoplastic disease of the cavernous ICA and the ATLS protocols for patients in oligemic shock. In reviewing reports of EC-IC bypass for high ICA injury, hemodynamically stable patients who were treated with bypass did better than those treated with carotid occlusion/ligation or conservative medical therapy.^{10,12-14} The subset of hemodynamically stable patients that received "conservative therapy" (carotid sacrifice or medical therapy) had the poorest results.

A previous study examined multiple variables to predict stroke from carotid sacrifice and found no pre-operative test to be conclusive.²⁵ These patients should be followed closely for hemispheric ischemic symptoms and may need surgical intervention in the future (i.e., bypass). Such patients with reduced cerebrovascular reserve are at higher risk for future stroke from occlusive vascular disease.^{25,26}

In reviewing the indications for bypass in ICA injury given by other authors, one can find support for our suggested management scheme. Rostomily¹⁴ reported a hemodynamically stable patient with evidence of impending ischemia (with "pial collaterals" from the contralateral MCA) who was treated conservatively and developed an ipsilateral infarction and expired. Two other patients in this same series were treated conservatively for pseudoaneurysm of the ICA. The first patient showed growth of the pseudoaneurysm that required bypass 5 weeks after initial therapy.¹⁴ The second patient was obtunded on admission and conservative therapy was instituted as the patient's prognosis was thought initially to be poor. However, the patient had a "marked neurological recovery" and the decision was made to bypass an enlarging pseudoaneurysm in the cervical ICA. The aneurysm fatally ruptured prior to his planned bypass procedure. In an earlier series, a patient was treated with anticoagulation in the presence of a pseudoaneurysm. The patient had an episode of acute epistaxis requiring balloon occlusion of the carotid to control the bleeding. The patient then developed a hemiparesis, which resolved following an EC-IC bypass.¹⁰ In these and similar cases, initial treatment with and EC-IC bypass and isolation of pseudoaneurysms might have yielded either a better outcome or fewer procedures, respectively (Table 4).

Table 4. Management Schema Applied to Historical Data

<i>Patient</i>	<i>Neurological Stability</i>	<i>Hemodynamic Stability</i>	<i>Procedure</i>	<i>Outcome</i>	<i>Reference</i>
J.S.	Stable, Pseudoaneurysm	Stable	EC-IC Bypass	no hemispheric deficit	present study
M.F.	Stable, Pseudoaneurysm	Stable	EC-IC Bypass	no hemispheric deficit	present study
M.B.	Stable, Pseudoaneurysm	Unstable	Ligation	no hemispheric deficit	present study
K.R.	Stable, Fistula and Pseudoaneurysm	Stable	EC-IC Bypass	no hemispheric deficit	present study
P.J.	Stable, Transection of ICA	Unstable	Ligation	no hemispheric deficit	present study
Case #1	Stable, Pseudoaneurysm	Stable	EC-IC Bypass	no hemispheric deficit	14
Case #2	Stable, Pseudoaneurysm	Stable	Ligation	later bypassed*	14
Case #3	Unstable, Carotid Occlusion	Stable	Nothing	Massive Stroke and Death	14
Case #4	Unstable, Pseudoaneurysm	Stable	Nothing	Hemorrhage and death prior to scheduled late bypass	14
Case #1	Unstable, poor collaterals	Stable	EC-IC Bypass	no hemispheric deficit	10
Case #2	Stable	Stable	Ligation	later bypassed†	10
C.W.	Unstable, hemiparesis	Stable	EC-IC Bypass	persistent hemiparesis	12
K.M.	Unstable, hemiparesis	Stable	EC-IC Bypass	complete resolution	12
M.P.	Unstable, hemiparesis	Stable	EC-IC Bypass	complete resolution	12
S.T.	Stable	Stable	EC-IC Bypass	no hemispheric deficit	12

Reference:^{10,12,14}

*Angiogram 4 weeks after initial injury showed increased size of previous pseudoaneurysm so patient was brought back to O.R. for bypass.

†Patient initially underwent coiling of carotid-cavernous fistula and anticoagulation for pseudoaneurysm. He then developed acute epistaxis from pseudoaneurysm and underwent carotid sacrifice by balloon occlusion, but developed hemiparesis and was treated with EC-IC bypass and resolution of symptoms.

CONCLUSION

Zone III missile injuries require a multidisciplinary approach. After airway management, initial resuscitation and a stereotyped trauma survey, a complete neurological examination is required, noting any lateralizing signs and cranial nerve deficits. Angiography and CT of the neck and cranium, supplemented by other indicated studies, are performed to complete the assessment. Angiography will indicate collateral flow and define the type of injury: transection, laceration, pseudoaneurysm, dissection, occlusion or fistula, and the exact location and vessel involved.^{1,10,16,17,27} This information and the status of the patient are crucial in planning a treatment strategy.

After missile injury, maintenance of carotid circulation and isolation of injured segments of the ICA are important measures to prevent or minimize cerebrovascular insult. Zone III injuries can be successfully managed using cranial base surgical approaches and carotid artery bypass in selected patients. Neoplastic and vascular lesions of the skull base have given us experience

with large caliber, high-flow EC-IC bypass techniques and management of lower cranial nerve deficits.^{2,28,29}

The decision to perform the bypass procedure is made based on the neurological status of the patient, hemodynamic stability of the patient, and the potential for cerebrovascular insult. Unstable patients with life-threatening injury treated with proximal ligation should be considered for elective bypass when stable to restore carotid circulation. Lower cranial nerve function is often affected and may require secondary surgical procedures, to correct neurological deficit, minimize morbidity or prevent mortality.

We utilize a scheme for treatment of penetrating ICA injury that offers expedient care to patients with potentially life-threatening and highly morbid injuries. This approach preserves the well-established, prioritized ATLS assessment while collecting crucial information as to the extent of vascular and neurological injury. Such information, coupled with radiographic measures, promotes a rational and consistent management plan for patients with a variety of hemodynamic and neurological profiles.

ACKNOWLEDGMENTS

The authors thank Jennifer M. Pryll, B.F.A. for digital photography and medical illustrations.

REFERENCES

1. Sclafani SJA, Panetta T, Goldstein AS, Phillips TF, Hotson G, Loh J, Shaftan GW. The management of arterial injuries caused by penetration of Zone III of the Neck. *J Trauma* 1985;871-881
2. Fisch U, Oldring DJ, Senning A. Surgical therapy of internal carotid artery lesions of the skull base and temporal bone. *Otolaryngol Head Neck Surg* 1980;548-554
3. Samson DS. Cervical carotid injuries. *Clin Neurosurg* 1982; 647-656
4. Sekhar LN, Sen C. Interposition vein grafts: Cervical to petrous ICA bypass. *J Neurosurg* 1990;968
5. Sekhar LN, Linskey ME. Management of the internal carotid artery in cranial base neoplasms. In Johnson JT, Didolkar MS: *Head and Neck Cancer*, Vol. III, Amsterdam: Elsevier Science Publishers, 1993
6. Sen C, Sekhar LN. Direct vein graft reconstruction of the cavernous, petrous and upper cervical internal carotid artery: Lessons learned from 30 cases. *Neurosurgery* 1992;732-743
7. LeFevre M. Concerning a case involving a wound of the carotid bulb caused by a bullet treated by ligation of the common carotid and end-to-end anastomosis of the external carotid with the internal carotid. *Bull Mem Soc Chir* 1918;923-925
8. Lawrence K, Shefts L, McDaniel J. Wounds of the common carotid arteries, report of seventeen cases from World War II. *Am J Surg* 1948;29
9. Bradley EL. Management of penetrating carotid injuries: An alternative approach. *J Trauma* 1973;248-255
10. D'Alise MD, Vardiman AB, Kopitnik TA, Batjer HH. External carotid to middle cerebral bypass in the treatment of complex internal carotid injury. *J Trauma* 1996;452-455
11. Fitchett J, Pomerantz M, Butsch D. Penetrating wounds of the neck: A military and civilian experience. *Arch Surg* 1969; 307-314
12. Gewertz BL, Samson DS, Ditmore QM, Bone GE. Management of penetrating injuries of the internal carotid artery at the base of the skull utilizing extracranial-intracranial bypass. *J Trauma* 1980;365-368
13. Fry RE, Fry WJ. Extracranial carotid injuries. *Surgery* 1980; 581-587
14. Rostomily RC, Newell DW, Grady MS, Wallace S, Nicholls S, Winn HR. Gunshot wounds of the internal carotid artery at the skull base: Management with vein bypass grafts and a review of the literature. *J Trauma* 1997;123-132
15. Liekweg WG, Greenfield LJ. Management of penetrating carotid arterial injury. *Ann Surg* 1978;587-592
16. Fabian TC, George SM, Croce MA, Mangiante EC, Voeller GR, Kudsk KA. Carotid artery trauma: Management based on mechanism of injury. *J Trauma* 1990;953-963
17. McInnis WD, Cruz AB, Aust JB. Penetrating injuries to the neck: Pitfalls in management. *Am J Surg* 1975;419-420
18. Schwartz JA, Turner D, Sheldon GF, Johnson G. Penetrating trauma of the internal carotid artery at the skull base. *J Cardiovasc Surg* 1987;542-545
19. Weaver FA, Albert EY, Wagner WH, Brooks SH, Weaver AA, Milford MA. The role of arterial reconstruction in penetrating carotid injuries. *Arch Surg* 1988;1106-1111
20. Marotta TR, Buller C, Taylor D, Morris C, Zwimpfer T. Autologous vein-covered stent repair of a cervical internal carotid artery aneurysm: Technical Case Report. *Neurosurgery* 1998; 408-413
21. Menawat SS, Dennis JW, Laneve LM, Frykberg ER. Are arteriograms necessary in penetrating zone II neck injuries? *J Vasc Surg* 1992;397-401
22. Sengupta D, Harper M, Jennett B. Effect of carotid ligation on cerebral blood flow in baboons. *J Neurol Neurosurg Psychiatry* 1973;736-741
23. Salar G, Mingrino S. Development of intracranial saccular aneurysms: Report of two cases. *Neurosurgery* 1981;462-465
24. Ledgerwood AM, Mullins RJ, Lucas CE. Primary repair vs. ligation for carotid artery injuries. *Arch Surg* 1980;488-493
25. Sekhar LN, Patel SJ. Permanent occlusion of the internal carotid artery during skull base and vascular surgery: Is it really safe? *Am J Otol* 1993;421
26. Schmiedek P, Piepgras A, Leinsinger G, Krisch C-M, Einhaupl K. Improvement of cerebrovascular reserve capacity by EC-IC bypass in patients with ICA occlusion and hemodynamic cerebral ischemia. *J Neurosurg* 1994;236-244
27. Jebara VA, Tabet GS, Ashoush R, Ghossain M, Harb J, Potorghese M, Uva MS, Saade B. Penetrating carotid injuries—A wartime experience. *J Vasc Surg* 1991;117-120
28. Samson D. Saphenous vein interposition grafts in the microsurgical treatment of cerebral ischemia. *Arch Surg* 1981; 1578-1582
29. Sekhar LN, Schramm VL, Jones NF. Operative exposure and management of the petrous and upper cervical internal carotid artery. *Neurosurgery* 1986;967-982