Craniofacial Resection: Decreased Complication Rate with a Modified Subcranial Approach

ABSTRACT—The authors have successfully utilized a modified subcranial approach to the anterior skull base, based upon the procedure first described by Joram Raveh, as an alternative to standard craniofacial resection. The complication rate of this procedure in 31 consecutive cases (28 tumors, 2 congenital malformations, and 1 mucocoele) has been 19.4% with no permanent complications, no deaths, no new neurological deficits, no brain injuries, no infections, and no seizures. Minor complications without permanent sequelae included two cases of tension pneumocephalus, a subdural hygroma, two transient cerebrospinal fluid leaks, and a case of bacterial meningitis secondary to fecal contamination of a lumbar drain in a child. Average length of hospitalization was 7.1 days (range 2 to 16 days). The overall complication rate is considerably below the complication rate for other reported craniofacial procedures. We describe the technique we have used and the results. The subcranial approach as described herein provides wide exposure of the anterior cranial base without brain retraction, does not require prolonged operating times or hospitalization, and has a potentially lower complication rate than reported for other transfrontal transbasal approaches.

Craniofacial approaches have become the procedures of choice for most tumors, trauma, and congenital anomalies involving the anterior cranial fossa and the orbits, nasal cavity, or paranasal sinuses. However, recent reports continue to document a complication rate of 39–50% and a mortality of 3–5% with these procedures,¹ prompting some authors to state that they are too morbid for routine use.² We have used a modified subcranial approach for a variety of lesions at the anterior cranial base and have achieved a lower complication rate than previously reported. We report our technique and results in 31 consecutive cases.

MATERIALS AND METHODS

This series consists of 31 consecutive patients operated upon over a 4-year period by the authors. Patients were evaluated in a multidisciplinary clinic staffed by a team representing the neurosurgery, otorhinolaryngology, and neuro-otology disciplines. Patients were considered appropriate for a subcranial approach when they had tumors, trauma, or congenital anomalies for which surgery was indicated and that anatomically involved the face, orbit(s), nasal cavity, and/or paranasal sinuses

with extension to the central skull base (cribriform plate, fovea ethmoidalis, planum sphenoidale), orbital roof(s), clivus, or anterior cavernous sinus.

TECHNIQUE

If a dural reconstruction is anticipated, a lumbar drain is placed at the beginning of the procedure. The patient is positioned supine with the head on a horse-shoe headrest unless intradural pathology requiring retractors mandates use of the Mayfield skull clamp. No hair is shaved. The hair is parted along the line of the bicoronal incision and maintained in position with rubber bands. The incision should be at least 13 cm posterior to the glabella if a pericranial flap long enough to reach the sphenoid sinus is needed. The eyelids are sutured shut with fine nylon suture. The head and face are prepped with betadine in standard fashion, as are any anticipated donor sites for fat grafts or free tissue transfers. Prophylactic antibiotic and mannitol are administered.

The incision is made with care to preserve the pericranium and temporalis fascia. If exposure of the zygoma is needed, the incision is extended inferiorly to just below the tragus in the natural crease anterior to the tragus. The pericranium is incised posterior to the incision to maximize the length of the pericranial flap and the entire scalp flap dissected anteriorly in the subperiosteal plane until the fat pad in the temporalis fascia is reached. The temporalis fascia is then incised radially bilaterally and the dissection continued anteriorly deep to the fascia to avoid injury to the frontalis branches of the facial nerve. The subperiosteal dissection is continued forward to expose the nasal bones and orbital rims laterally to the frontozygomatic sutures. The supraorbital neurovascular bundles must be preserved by careful dissection from their canals or notches. The periorbital is dissected from the orbital walls medially and the anterior ethmoid arteries are ligated. During this dissection, it is important to maintain the integrity of the medial canthal ligaments. The scalp flap and its critical pericranium are covered with moist sponges and carefully protected throughout the case.

A small incision is made in the temporalis muscle on one side and a single burhole placed so that it will be covered by subsequent closure of the muscle. The dura is stripped from the inner table of the skull and a small, low unilateral craniotomy sawed from about the lateral canthus to a point 10 to 15 mm short of the midline. The dura and superior sagittal sinus are then stripped contralaterally under direct vision and a second, contralateral craniotomy removed. The frontal sinus is cranialized to enlarge the working space intracranially. The dura is stripped from the orbital roofs and region of the crista galli to provide access for the orbitonasal osteotomy and covered with moist cottonoids. While the otorhinolaryngologist is performing the orbitonasal os-

teotomy, the neurosurgical team is plating the two craniotomy bones back together on a back table using titanium microplates. Residual frontal sinus on the frontal bones is drilled away to prevent mucocele formation.

Osteotomies are then made vertically through the orbital rims (Fig. 1). While protecting the orbital contents with malleable brain retractors, a side cutting bur is used to make an osteotomy across the orbital roof to the contralateral side. This osteotomy traverses the anterior portion of the fovea ethmoidalis and passes anterior to the crista galli, leaving the olfactory apparatus intact. Bilateral osteotomies are then made in the medial walls of the orbits extending inferomedially from the roof of the orbit to a point just anterior to the lacrimal crest and then onto the nasal process of the maxilla. An osteotomy of the nasal bones is performed connecting the two maxillary cuts. Several millimeters of the nasal bones are left in place and attached to the upper lateral cartilages to prevent valvular collapse. An osteotome is then used from the intracranial side to detach the nasal septum from the nasoglabellar complex. The orbitonasal osteotomy is then removed (Fig. 2), drilled free of frontal sinus mucosa, and microplates attached laterally and anteriorly for eventual replacement.

If necessary for en bloc resection, one or both olfactory nerves in their dural sleeves are severed as inferiorly as possible. Careful extradural removal of the crista galli aids in this dissection. Cerebrospinal fluid (CSF) will of necessity leak from these incisions, which are promptly closed. The dura may now be stripped posteriorly as far as the tuberculum sellae, if indicated. No brain retraction is necessary to accomplish this due to the low approach afforded by the removal of the orbitonasal complex. Both olfactory nerves may be spared using the approach described by Spetzler et al.³

If there is an intradural component of the tumor, it is resected at this time using standard neurosurgical techniques, and the dura closed or reconstructed as indicated. Osteotomies of the frontal floor are then performed and the tumor is resected en bloc. Facial degloving may be necessary to free the inferior end of tumors reaching the hard palate. Orbital exenteration is performed if indicated for some malignant tumors.

When complete tumor removal is confirmed by pathological examination of frozen sections from the margins of the resection, the reconstruction is begun. Reconstruction of the orbital walls or frontal bones is done using split calvarium when indicated. We do not routinely use bone in reconstruction of the frontal floor. Drill holes are placed in the remaining bone edges along the medial orbital roof and the roof of sphenoid sinus. Anchoring sutures for the pericranial flap are placed in these holes and the needles left in place. A vascularized pericranial flap is then carefully raised. If no orbital exenteration has been performed, the orbitonasal bar is now plated in place and then the pericranial flap placed in the saw kerf between the orbitonasal bar and the cran-

Figure 1. Anterior (A) and posterior (B) views of a skull model illustrating the saw cuts used for the low bifrontal craniotomy and orbitonasal osteotomy.

iotomy bones. If an orbit has been exenterated, the pericranial flap is placed prior to replacement of the orbitonasal bar so as to provide living tissue coverage for the orbital rim on the exenterated side. The anchoring sutures are then passed through the pericranial flap from the external side about 15 mm from the end. When the flap is then pulled into place, this cuff of extra pericranium will hang down into the sphenoid sinus. The anchoring sutures are tied to draw the flap into place. If a facial degloving incision or orbital exenteration has been performed, the flap may be inspected from below to be certain that no dura can be seen. An endoscope may be used for the same purpose if no facial exposure has been made. If a free tissue transfer is necessary for closure, it is performed at this time. If cranialization of a large frontal sinus or placement of the pericranial flap below the orbitonasal complex has left a large dead space between the pericranial flap and the dura, a free fat graft is used to obliterate this space. The dura is tacked up to the bone edges, the craniotomy is replaced, and the wound closed. If an orbital exenteration has

been performed, the pericranial flap is covered with a split thickness skin graft. Nasal and/or orbital packing is placed as indicated and nasal trumpets are placed to divert the air stream from the reconstruction and lower the risk of pneumocephalus.

Steroids are not used postoperatively unless indicated for intradural disease. Antibiotics are continued until all drains and packing are removed. The lumbar drain is initially set at 3 cc per hr and gradually increased to 6 cc/hr if there is no indication of progressive pneumocephalus. Drainage is continued for 72 hr. The drain is then capped and the patient observed for CSF leak overnight. If no leak is detected, the drain is removed. Patients are advised to avoid nose blowing, straining, Val Salva maneuver, or placing the head in a dependent position for 4–6 weeks. Saline nasal irrigation is recommended to minimize crusting.

RESULTS

In the period from January, 1994 to March, 1998, 31 patients underwent a subcranial operation using the approach described above (Figs. 3 and 4). The final pathological diagnoses are listed in Table 1. The average estimated blood loss for the cases was 802 cc. The average operative time was 6.5 hr. Patients spent an average of 1 day in the intensive care unit and were hospitalized for an average of 7.1 days (range 2 days to 16 days). One patient previously operated and irradiated for a malignant meningioma required a free tissue transfer as part of the primary procedure and was hospitalized for 16 days. Follow-up time ranged from 4 months to 50 months (mean 21 months).

There were 6 complications in 6 patients in this series (19.4%). There were no new neurological deficits, no brain injuries, no seizures, no wound infections, no cases of osteomyelitis, and no unexpected cosmetic deformities. Two patients developed a tension pneumo-

Figure 2. Photograph of the orbitonasal osteotomy illustrating the extent of removal of the nasal bones and orbital rims.

Figure 3. Preoperative sagittal (A) and coronal (B) T1-weighted gadolinium-enhanced MRI scans of a patient with a sinonasal undifferentiated carcinoma.

cephalus during the first postoperative day requiring temporary tracheotomy. There were no unusual features of these procedures to suggest why these developed. One patient had undergone only a unilateral sectioning of the olfactory nerve and primary closure of this very small dural defect. One patient developed a subdural hygroma, which required bur hole drainage. There were two delayed CSF leaks. One patient had been previously irradiated and underwent an orbital exenteration

Figure 4. Postoperative coronal T1-weighted gadolinium-enhanced MRI scan of the patient in Figure 2 showing complete resection of the tumor and the expected enhancement of the vascularized pericranial flap used in the reconstruction.

closed with a vascularized pericranial flap and a split thickness skin graft. This repair broke down 5 weeks postoperatively and required a radial forearm free tissue transfer for repair. The second patient was nursed in the Trendelenberg position in the recovery room at the order of the anesthesiologist, resulting in massive CSF

Table 1. Histologic Diagnoses in 31 Subcranial Resections

Carcinoma, adenocarcinoma	2
Carcinoma, adenoid cystic	2
Carcinoma, carcinoid	1
Carcinoma, Hurthle cell	1
Carcinoma, sinonasal undifferentiated	3
Carcinoma, squamous cell	3
Chondroblastoma	1
Congenital absence of sphenoid wing	1
Encephalocele	1
Esthesioneuroblastoma	4
Hemangiopericytoma	1
Inverted papilloma	3
Melanoma	2
Meningioma, malignant	1
Mucocele	1
Myxoma of orbit	1
Ossifying fibroma	1
Sarcoma, fibrosarcoma	1
Sarcoma, osteogenic	1

egress and requiring immediate removal of the lumbar drain to prevent further brain collapse. He then developed an obvious CSF leak 2 weeks later, which responded to lumbar drainage. One child developed *Streptococcus faecalis meningitis* after defecating on the lumbar drain. This was cured with antibiotics without sequelae.

DISCUSSION

When applied to tumors arising in the anterior skull base and paranasal sinuses, early use of the transfrontal approach usually was combined with a transfacial approach.⁴⁻⁷ These reports described a significant incidence of death, CSF fistula, and infection. The further development of the transfrontal transbasal approach is usually ascribed to Derome.⁸ Experience with this approach led to the elimination of facial incisions in many cases.^{9,10} Removal of the orbital rims to facilitate exposure of the cranial base was reported as early as 1913.¹¹ The modern refinement of the bifrontal craniotomy by the addition of orbital osteotomies have been reported by numerous recent authors.¹²⁻²⁰ These orbitofrontal operations have been given various names including the extended frontal approach,¹⁸ the extensive transbasal approach,¹⁶ and the telecanthal approach.¹²

The subcranial approach was pioneered by Raveh,^{21,22} for repair of trauma, congenital anomalies,²³ and tumors^{24,25} of the anterior skull base. The subcranial approach differs from these other orbitocranial approaches by including more of the nasal bones in the orbitonasal osteotomy and in not involving detachment of the medial canthal ligaments or mobilization of the orbits. Unless the tumor transgresses the dura, we use an extradural approach for resection of the frontal floor, preferring not to take the dura with the specimen unless oncologically indicated.²⁶

The complication rate for craniofacial resections has fallen steadily in the 35 years since Ketcham's report in 1963.^{1,4,18,19,27-30} In one review of recent series, the overall complication rate was estimated at 39–50% and the mortality rate at 3–5%.¹ In looking at recent reports, Sekhar et al¹⁸ reported 19 complications in 49 (39%) patients undergoing an extended frontal approach, although most of these patients had additional procedures as well. Frontal lobe injury, seizures, infections, and neurologic deterioration were seen in 3, 3, 2, and 2 patients, respectively. Deschler et al¹ reported an overall complication rate of 40% in 52 patients undergoing combined transcranial and transfacial approaches, with 10 infections and 3 brain injuries secondary to retraction. Close and Mickey²⁶ reported 2 infections and 1 CSF leak in 11 patients undergoing transcranial resections. McCutcheon et al³¹ reported on

76 patients with 6 postoperative hematomas requiring surgical evacuation, 5 cases of pneumocephalus, 3 CSF leaks, 2 bifrontal cerebral infarctions, and 1 death, with an overall complication rate of 37% (28/76).

The complication rate for this series using the subcranial approach compares favorably with rates for other series of anterior skull base procedures. Our overall complication rate was 19.4%, with no permanent complications, and with two complications not being directly related to the procedure itself, but to technical errors during the postoperative period. We believe that the complication rate will be further lowered by increasing experience with this approach, and that pessimism regarding the utility of this approach is unwarranted.² It is important to emphasize that the subcranial approach, with its aggressive removal of orbital and nasal bone, is performed without the use of brain retraction when there is no intradural pathology, thereby minimizing the risk of parenchymal injury present in other series. Unlike other series, we have not had any epidural or bone flap infections. Although the reasons for this are not certain, careful tissue handling, good hemostasis, and meticulous reconstruction may contribute to minimizing the risk. In addition, nasal packing is used briefly or not at all and this may decrease the incidence of infection.

Pneumocephalus remains a potential complication for which an absolute solution is not yet available and which is potentially caused or exacerbated by lumbar drainage. However, we believe that lumbar drainage is an important part of ensuring an adequate repair. We do not start lumbar drainage until the patient has been examined and found to be awake and alert and then we begin at only 3 cc/hr, gradually increasing the rate every 12 hr if there is no evidence of decreased level of alertness. The use of nasal trumpets to divert exhaled air from the cranial base remains an unproven adjunct in this regard. We do not utilize routine tracheostomy due to the low incidence of pneumocephalus overall.

One patient who had been previously radiated developed a CSF leak 5 weeks after surgery through an enterated orbit. The orbit had been repaired with a vascularized pericranial flap covered with a split thickness skin graft. Although this repair was satisfactory in five other patients in this series, they had not been previously irradiated. Although a single case is not always a valid test, we will likely recommend free tissue transfer should this situation arise again.

The subcranial approach as performed in this series has been safe and effective. The procedure is well tolerated by patients with an average length of stay below the average 10.5 days for craniotomy, age >17, except for trauma according the 1994 Health Care Finance Administration data. The complication rate is below that published for other anterior cranial base procedures. We believe that this approach is preferable to frontal cran-

iotomy alone or to craniotomy combined with transfacial approaches requiring facial incisions.

ACKNOWLEDGMENTS

This paper is dedicated to the memory of Matthew Vogel, who fought his disease courageously and inspired us all.

REFERENCES

1. Deschler DG, Gutin PH, Mamelak AN, McDermott MW, Kaplan MJ. Complications of anterior skull base surgery. *Skull Base Surg* 1996;6:113-118
2. Goel A. Letter to the Editor. *Neurosurgery* 1997;40:218
3. Spetzler RF, Herman JM, Beals S, Joganic E, Milligan J. Preservation of olfaction in anterior craniofacial approaches. *J Neurosurg* 1993;79:48-52
4. Ketcham AS, Wilkins RH, Van Buren JM, Smith RR. A combined intracranial facial approach to the paranasal sinuses. *Am J Surg* 1963;106:698-703
5. Ray BS, McLean JM. Combined intracranial and orbital operation for retinoblastoma. *Arch Ophthalmol* 1943;30:437-445
6. Smith RR, Klopp CT, Williams JM. Surgical treatment of cancer of the frontal sinus and adjacent areas. *Cancer* 1954;7:991-994
7. Van Buren JM, Ommaya AK, Ketcham AS. Ten years' experience with radical combined craniofacial resection of malignant tumors of the paranasal sinuses. *J Neurosurg* 1968;28:341-350
8. Derome P. Les tumeurs spheno-ethmoidales. Possibilites de'ex-ereuse et de reparation chirurgicales. *Neurochirurgie* 1972;18 (Suppl 1):1-164
9. Arita N, Mori S, Sano M, et al. Surgical treatment of tumors in the anterior skull base using the transbasal approach. *Neurosurgery* 1989;24:379-384
10. Blacklock JB, Weber RS, Lee Y-Y, Goepfert H. Transcranial resection of tumors of the paranasal sinuses and nasal cavity. *J Neurosurg* 1989;71:10-15
11. Frazier CH. An approach to the hypophysis through the anterior cranial fossa. *Ann Surg* 1913;57:145-150
12. Fujitsu K, Saijoh M, Aoki F, et al. Telecanthal approach for meningiomas in the ethmoid and sphenoid sinuses. *Neurosurgery* 1991;28:714-720
13. Haines SJ, Marentette LJ, Wirtschafter JD. Extended fronto-orbital approaches to the anterior cranial base: Variations on a theme. *Skull Base Surg* 1992;2:134-141
14. Jackson IT, Marsh WR, Hide TA. Treatment of tumors involving the anterior cranial fossa. *Head Neck Surg* 1984;6:901-913
15. Jane JA, Park TS, Pobereskin LH, Winn HR, Butler AB. The supraorbital approach: Technical note. *Neurosurgery* 1982;11:537-542
16. Kawakami, K, Yamanouchi Y, Kawamura Y, Matsumura H. Operative approach to the frontal skull base: Extensive transbasal approach. *Neurosurgery* 1991;28:720-725
17. Persing JA, Jane JA, Levine PA, Cantrell RW. The versatile frontal sinus approach to the floor of the anterior cranial fossa. Technical note. *J Neurosurg* 1990;72:513-516
18. Sekhar LN, Nanda A, Sen CN, Snyderman CN, Janecka IP. The extended frontal approach to tumors of the anterior, middle, and posterior skull base. *J Neurosurg* 1992;76:198-206
19. Shah JP, Sundaresan N, Galicich J, Strong EW. Craniofacial resection for tumors involving the base of the skull. *Am J Surg* 1987;154:352-358
20. Sundaresan N, Shah JP. Craniofacial resection for anterior skull base tumors. *Head Neck Surg* 1988;10:219-224
21. Raveh J, Vuillemin T. The surgical one stage management of combined craniomaxillofacial and frontobasal fractures: Advantage of the subcranial approach in 374 cases. *J Craniomaxillofac Surg* 1988;16:160-172
22. Raveh J, Vuillemin T. Subcranial management of 395 combined frontobasal-midface fractures. *Arch Otolaryngol Head Neck Surg* 1989;114:1115-1122
23. Raveh J, Vuillemin T. Advantages of an additional subcranial approach in the correction of craniofacial deformities. *J Craniomaxillofac Surg* 1988;16:350-358
24. Raveh J, Vuillemin T. The subcranial-supraorbital temporal approach for tumor resection. *J Craniofac Surg* 1990;1:53-59
25. Raveh J, Laedrach K, Speiser M, et al. The subcranial approach for fronto-orbital and anteroposterior skull-base tumors. *Arch Otolaryngol Head Neck Surg* 1993;119:385-393
26. Close LG, Mickey B. Transcranial resection of ethmoid sinus cancer involving the anterior skull base. *Skull Base Surg* 1992;2:213-219
27. Ketcham AS, Hoye RC, Van Buren JM, Johnson RH, Smith RR. Complications of intracranial facial resection for tumors of the paranasal sinuses. *Am J Surg* 1966;112:591-596
28. Levine PA, Scher RL, Jane JA, et al. The craniofacial resection: Eleven-year experience at the University of Virginia: Problems and solutions. *Otolaryngol Head Neck Surg* 1989;101:665-669
29. Richtsmeier WJ, Briggs RJ, Koch WM, et al. Complications and early outcome of anterior craniofacial resection. *Arch Otolaryngol Head Neck Surg* 1992;118:913-917
30. Terz JJ, Young HF, Lawrence W. Combined craniofacial resection for locally advanced carcinoma of the head and neck: Carcinoma of the paranasal sinuses. *Am J Surg* 1980;140:618-624
31. McCutcheon IE, Blacklock JB, Weber RS, et al. Anterior transcranial (craniofacial) resection of tumors of the paranasal sinuses: Surgical technique and results. *Neurosurgery* 1996;38:471-480

REVIEWER'S COMMENTS

Ross et al. have described nicely the technique of subcranial approach used for a large number of malignant tumors involving the anterior cranial fossa. The authors have described a variety of pathological conditions, a large number of which represent malignant problems. They report a complication rate of 19.4% with no mortality in the postoperative period. There were no complications related to neurological deficit, brain injury, or seizures. In this series they use free-flap reconstruction only in one patient. When craniofacial resection is performed for tumors of the paranasal sinuses involving the anterior skull base, especially when orbital exenteration is performed, the defect is best reconstructed by the use of free flap, especially the rectus abdominis flap. The free-flap reconstruction offers excellent soft tissue coverage and decreases significantly the problems related to CSF leak. The technique of subcranial approach should be familiar to every skull base surgeon practicing craniofacial resections.

Ashoe Shaha, M.D.