10. Selesnick SH, Jackler RK, Pitts LW. The changing clinical presentation of acoustic tumors in the MRI era. *Laryngoscope* 1993;103:431-436
11. Lassaletta L, Fontes L, Melcón E, Sarria MJ, Gavilan J. Hearing preservation with the retrosigmoid approach: Myth or reality? 102nd Annual Meeting of the American Academy of Otolaryngology-Head Neck Surgery; September 8-12, 2001; Denver, CO
12. Fagan PA, Misra SN, Doust B. Facial neuroma of the cerebellopontine angle and the internal auditory canal. *Laryngoscope* 1993;103:442-446
13. Martin N, Sterkers O, Mompoin D, Nahum H. Facial nerve neuromas: MR imaging. Report of four cases. *Neuroradiology* 1992;34:62-67
14. Parnes LS, Lee DH, Peerless SJ. Magnetic resonance imaging of facial nerve neuromas. *Laryngoscope* 1991;101:31-35
15. Latack JT, Gabrielsen TO, Knake JE, et al. Facial nerve neuromas: radiologic evaluation. *Radiology* 1983;149:731-739
16. Fenton JE, Chin RY, Tonkin JP, Fagan PA. Transtemporal facial nerve schwannoma without facial nerve paralysis. *J Laryngol Otol* 2001;115:559-560
17. Liu R, Fagan P. Facial nerve schwannoma: surgical excision versus conservative management. *Ann Otol Rhinol Laryngol* 2001;110:1025-1029

Commentary

The authors present an interesting case of a 56-year-old woman with a right-sided cerebellopontine facial nerve neuroma that extended into

the internal auditory canal. The authors chose the retrosigmoid approach to preserve preoperative hearing. Intraoperatively, they found that the tumor involved the facial nerve as evidenced by the presence of the vestibular and cochlear nerves as shown on intact intraoperative photograph. These nerves were displaced superoposteriorly. The authors indicated that complete resection was achieved. That they could achieve a sural nerve graft is truly surprising, especially given that the tumor extended almost a centimeter into the internal auditory canal. Whether the anastomosis will provide some recovery of facial function remains to be seen. Not surprisingly, the patient had complete facial paralysis.

It is easy to play armchair quarterback; however, if we find what appears to be a facial neuroma during an approach to a cerebellopontine angle tumor, we seldom recommend obtaining a biopsy of these tumors, let alone resecting them. In our experience and that of others, complete facial paralysis would be guaranteed. If these small, benign tumors were to enlarge, radiosurgery or possibly open surgery could be considered.

Randall W. Porter, M.D.¹