of the occipital condyle. *Clin Neurol Neurosurg* 1998;100: 44–45

13. Bridgman SA, McNab W. Traumatic occipital condyle fracture, multiple cranial nerve palsies, and torticollis: a case report and review of the literature. *Surg Neurol* 1992; 38:152–156
14. Legros B, Fournier P, Chiaroni P, Ritz O, Fuscuardi J. Basal fracture of the skull and lower (IX, X, XI, XII) cranial nerves palsy: four case reports including two fractures of the occipital condyle—a literature review. *J Trauma* 2000; 48:342–348
15. Wani MA, Tandon PN, Banerji AK, Bhatia R. Collet–Sicard syndrome resulting from closed head injury: case report. *J Trauma* 1991;31:1437–1439
16. Bozboga M, Unal F, Hepgul K, Izgi N, Turantan I, Turker K. Fracture of the occipital condyle—case report. *Spine* 1992;17:1119–1121
17. Massaro F, Lanotte M. Fracture of the occipital condyle. *Injury* 1993;24:419–420

Commentary

This article provides a straightforward description of two patients suffering from an unusual skull base fracture and useful information for our readership. The discussion helps underscore the three accepted types of occipital fractures and how each type generally dictates prognosis and treatment.

Clarence T. Sasaki, M.D.¹