short-term follow-up. The extent of decompression of the optic canal is likely to be a factor in recurrence because any residual bone is likely to contain fibrous dysplasia.¹ Thus, as more walls are decompressed, the likelihood decreases that continued growth of residual fibrous dysplasia will entrap the optic nerve. The extensive decompression required and the risk associated with intracranial facial surgery may necessitate the transcranial extradural approach.

REFERENCES

- Chen Y-R, Breidahl A, Chang C-N. Optic nerve decompression in fibrous dysplasia: indications, efficacy, and safety. *Plast Reconstr Surg* 1997;99:22-30
- Liakos GM, Walker CB, Carruth JAS. Ocular complications in craniofacial fibrous dysplasia. *Br J Ophthalmol* 1979;63:611-616
- Papadopoulos MC, Casey AT, Powell M. Craniofacial fibrous dysplasia complicated by acute, reversible visual loss: report of two cases. *Br J Neurosurg* 1998;12:159-161
- Edgerton MT, Persing JA, Jane JA. The surgical treatment of fibrous dysplasia with emphasis on recent contributions from cranio-maxillo-facial surgery. *Ann Surg* 1985;202:459-479
- Katz BJ, Nerad JA. Ophthalmic manifestations of fibrous dysplasia: a disease of children and adults. *Ophthalmology* 1998;105:2207-2215
- Sassin JF, Rosenberg RN. Neurological complications of fibrous dysplasia of the skull. *Arch Neurol* 1968;18:363-369
- Harris WH, Dudley HR Jr, Barry RJ. The natural history of fibrous dysplasia. *J Bone Joint Surg* 1962;44A:207-233
- Leeds N, Seaman WB. Fibrous dysplasia of the skull and its differential diagnosis. *Radiology* 1962;78:570-582
- Schwartz DT, Alpert M. The malignant transformation of fibrous dysplasia. *Am J Med* 1962;32:393-403
- Lüdecke DK, Treige W. Pressure-irrigation-suction system. Technical note. *Acta Neurochirurgica (Wien)* 1982;66:123-126
- Papay FA, Morales L Jr, Flaharty P, et al. Optic nerve decompression in cranial base fibrous dysplasia. *J Craniofac Surg* 1995;6:5-10
- Weisman JS, Hepler RS, Vinters HV. Reversible visual loss caused by fibrous dysplasia. *Am J Ophthalmol* 1990;110:244-249
- Kurokawa Y, Sohma T, Tsuchita H, Kitami K, Suzuki S, Sohma K. Hemorrhage into fibrous dysplasia following minor head injury. Effective decompression for the ophthalmic artery and optic nerve. *Surg Neurol* 1989;32:421-426
- Bland LI, Marchese MJ, McDonald JV. Acute monocular blindness secondary to fibrous dysplasia of the skull: a case report. *Ann Ophthalmol* 1992;24:263-266
- Saito K, Suzuki Y, Nehashi K, Sugita K. Unilateral extradural approach for bilateral optic canal release in a patient with fibrous dysplasia. *Surg Neurol* 1990;34:124-128
- Michael CB, Lee AG, Patrinely JR, Stal S, Blacklock JB. Visual loss associated with fibrous dysplasia of the anterior skull base. Case report and review of the literature. *J Neurosurg* 2000;92:350-354
- Osguthorpe JD, Gudeman SK. Orbital complications of fibrous dysplasia. *Otolaryngol Head Neck Surg* 1986;97:403-405
- Chen Y-R, Fairholm D. Fronto-orbital-sphenoidal fibrous dysplasia. *Ann Plast Surg* 1985;15:190-203
- Chen Y-R, Noordhoff MS. Treatment of craniomaxillofacial fibrous dysplasia: how early and how extensive? *Plast Reconstr Surg* 1990;86:835-842
- Finney HL, Roberts TS. Fibrous dysplasia of the skull with progressive cranial nerve involvement. *Surg Neurol* 1976;6:341-343

Commentary

This small series of patients with orbito-frontal fibrous dysplasia who underwent optic nerve decompression describes the well-respected senior surgeon's operative techniques and his argument for prophylactic optic nerve decompression in asymptomatic patients. The authors treated three patients, two of whom were asymptomatic with respect to visual symptoms. The authors illustrate their anatomical results with computed tomography scans.

The authors used an extradural transcranial approach. They rightly emphasize the use of lumbar drainage and continuous irrigation and suction with a micropressure suction-irrigation system previously described by Lüdecke and Triage.¹ The authors state that "because predicting the arrest of pericanal fibrous dysplasia is difficult, the growth of fibrous dysplastic bone should be assumed to continue and optic nerve compression should be assumed to be relatively imminent." Unfortunately, this assumption is not based on natural history data, and the opposite assumption could just as

easily be made. The authors' argument that "decompression of the optic nerve is likely to be of benefit on the theoretical grounds that once vision has been impaired, the chances of being able to restore vision are greatly reduced" is actually undermined by numerous other reports as well as by their own experience as demonstrated by Patient 3 who presented with loss of visual acuity. This patient's vision improved significantly after decompression of the optic nerve. Furthermore, the postoperative nasal field cut, fortunately transient, incurred by Patient 2 exemplifies the clear danger of intervention in previously visually asymptomatic patients.

Until further prospective data about progression of symptoms in patients with initially asymp-

tomatic frontal orbital fibrous dysplasia are available, it will be impossible to gauge the true risk-benefit ratio for optic nerve decompression in asymptomatic patients. Until such natural history data are available, I advocate early intervention in symptomatic patients only.

REFERENCE

1. Lüdecke DK, Treige W. Pressure-irrigation-suction system. Technical note. *Acta Neurochirurgica (Wien)* 1982;66:123-126

B. Gregory Thompson, M.D.¹