been developing such systems for the management of a variety of medical conditions,² of which a system for the diagnosis, physical examination, and treatment of hypercholesterolaemia is currently undergoing field trials with general practitioners in the west midlands. The system is supplied for the use of a Psion Organiser and incorporates regional knowledge in a particular subject as appropriate for use in primary care or by junior hospital medical staff. The medical knowledge supplied facilitates patient management up to a level considered appropriate for the general practitioner or non-specialist physician and contains locally determined referral criteria for specialist attention. Preliminary results suggest that such a format is acceptable to both general practitioners and practice nurses and that programmable pocket computers are an appropriate vehicle for the devolution of medical expertise to non-specialists or non-medical staff such as practice nurses. Inappropriate referrals to hospital clinics may also be reduced by providing clear guidelines to referring general practitioners with this medium.

We also envisage using such systems for providing useful advice on foreign travel prophylaxis, poisoning, laboratory sampling requirements, protocols for standard medical tests, etc.

The potential for devolution of medical expertise is just beginning to be realised. Pocket sized computers can retain and instantaneously recall a large amount of medically relevant data and, unlike a text book, can be readily updated, are portable, and have a variety of other functions.

M F RYAN N CORBETT
I CLARK M PETERS

Department of Medical Computing,
Wolfson Research Laboratories,
Queen Elizabeth Medical Centre,
Birmingham B15 2TH

1 Wyatt J. Construction of clinical scoring systems. *Br Med J* 1990;300:538-9. (24 February.)

2 Ryan MF, Clark I, Peters M, Cramb R. A low cost indicator for rationalisation of lipid clinic referrals. In: Halloran SP, ed. *Focus '89*. London: Association of Clinical Biochemists, 1989:68.

Streptokinase in acute aortic dissection

SIR,—The report by Mr J Butler and colleagues emphasises the risk of leaving major medical management decisions to inexperienced junior medical staff.¹ In each of the cases described the house officer was apparently the first medical practitioner to see patients admitted as emergencies. This is an illogical method for assessing patients. The administration of potentially life saving but also potentially life threatening treatments surely requires a higher level of skill and experience than should be expected from a house officer. It is noteworthy that in all of the patients described in this report aortic dissection should have been considered on the basis of the chest radiograph, which was available to the admitting house officer. There is insufficient data in the report by Dr Nicholas P Curzen and colleagues² to judge whether the same circumstances occurred in their patients, although in one case the chest radiograph was suggestive of aortic dissection.

In the study by Mr Butler and colleagues it is not surprising that the diagnosis of aortic dissection was not considered by the admitting house officer, both in view of the limited experience in cardiovascular diseases that most medical students receive during their training and also the limited exposure to training in radiology in most undergraduate medical training.

The problems concerned in distinguishing acute aortic dissection from other causes of severe acute chest pain should not cause the uses of thrombolytic agents to be reduced. Rather it is

an important reason for changing the way in which patients admitted with acute conditions are assessed. Surely it is no longer satisfactory to delegate assessment of acutely ill patients to the most junior members of the medical team; this should now be the responsibility of a more senior (that is, registrar grade) admitting physician. This will improve the accuracy of the initial management of these patients and will help to prevent the potential catastrophes described by Mr Butler and colleagues. It may also help to prevent an epidemic of unnecessary requests for investigation for possible aortic dissection before doctors administer thrombolytic treatment.

G HARTNELL

Department of Radiodiagnosis,
Bristol Royal Infirmary,
Bristol BS2 8HW

1 Butler J, Davies AH, Westaby S. Streptokinase in acute aortic dissection. *Br Med J* 1990;300:517-9. (24 February.)

2 Curzen NP, Clarke B, Gray HH. Intravenous thrombolysis for suspected myocardial infarction: a cautionary note. *Br Med J* 1990;300:513. (24 February.)

Medical problems of adults who were sexually abused in childhood

SIR,—Assessing the possible effects of adversity during childhood on health in adulthood requires studies that either include appropriate control subjects for comparison or eliminate the effects of confounding variables statistically. In their study on the medical problems of adults who were sexually abused in childhood Dr R P Arnold and colleagues used neither of these techniques.¹ This leads to most basic problems in interpreting their results. For example, we would surely expect the total number of medical specialties contacted by any person (whether sexually abused in childhood or not) to increase with age. The data presented (tables I and III) show exactly this expected correlation between age and the number of specialties contacted for the seven abused women (Spearman's test, $\rho=0.88$; $z=2.16$; $p<0.05$). This renders the interpretation of the total specialties contacted extremely hazardous without comparison with appropriate matched controls.

Dr Arnold and colleagues suggest that childhood sexual abuse may be an aetiological factor in the development of physical symptoms in adulthood. Their data support this hypothesis weakly. The duration of abuse correlates positively with the number of non-psychiatric admissions (derived from table II) ($\rho=0.81$; $z=1.98$; $p<0.05$). The interpretation of this result, however, is unclear as the study confounds sexual abuse with severe family disturbances, owing to biased sampling. The authors themselves point out that they ascertained that most of the patients "had unambiguous histories of severe family problems in childhood," so that the group was "not representative of all cases of childhood sexual abuse." Clearly, comparison with carefully selected control subjects (for example, people exposed as children to comparable durations of severe marital discord or violence but not to sexual abuse) is needed before we can attribute any excess ill health in adulthood specifically to the sexual abuse component of family disturbance during childhood.

The null hypothesis that sexual abuse during childhood does not alter demand for medical services in adulthood cannot be rejected without much more careful work. The topic of childhood sexual abuse is so sensitive that studies to assess its possible long term effects should meet the highest scientific standards if their results are to inform rather than inflame public debate.

JONATHAN H WILLIAMS

Banbury, Oxfordshire

1 Arnold RP, Rogers D, Cook DAG. Medical problems of adults who were sexually abused in childhood. *Br Med J* 1990;300:705-8. (17 March.)

2 Meddis R. *Statistics using ranks: a unified approach*. Oxford: Basil Blackwell, 1984.

SIR,—In their paper Dr R P Arnold and colleagues draw attention to the possible long term physical effects of childhood sexual abuse.¹ Research on this subject is less extensive than that on the psychological effects, in fact it is paltry: to our knowledge there have been only three studies in non-clinical populations.^{2,4} There are problems, however, in drawing any conclusions from the type of model employed in the paper by Dr Arnold and colleagues. It is debatable whether adult psychiatric patients are a useful sample to study as the prevalence of childhood sexual abuse in such patients is consistently higher than expected.⁵ In addition, psychiatric patients have a greater incidence of physical illness than non-psychiatric patients and may well be referred more and have a higher rate of interventions as a group.^{6,7} Indeed, any samples comprising patients with clinical complaints are problematical—some surveys performed at normal outpatient gynaecological clinics, for example, suggest that the prevalence of childhood sexual abuse in such patients may be extremely high, perhaps in the region of 65%.⁸

The authors' method of finding patients was also interesting; they seem to have been found by chance and not elicited by a structured interview or questionnaire, as advocated by some authorities.⁹ Therefore, they may well not have been representative of patients who were sexually abused in childhood—for example, one patient had made 60 attempts at suicide.

One particularly important point is the statement, which is often cited, that victims of childhood sexual abuse seem to suffer abdominal pain, headache, and backache in adulthood. There is no evidence of this from studies in non-clinical samples, and though Bachmann *et al*¹⁰ state this in their review, the original sources do not say this.¹¹ Of course, there have been studies in patients with chronic pelvic pain that have found a high prevalence of childhood sexual abuse,^{12,14} but this is not the same thing at all. This is quite apart from the issue of whether the high prevalence of childhood sexual abuse found in these populations is any greater than the prevalence in the general population, in which estimates in women have varied from 6% to 62%¹⁵; in this light findings of prevalences of 35%¹⁴ or 64%¹⁵ have far less clear implications. It suits many of us to believe that there is a clear somatic link between childhood sexual abuse and later physical symptoms, but the evidence simply does not yet exist.

The study by Dr Arnold and colleagues is based on a small, highly selected population in which case definition is unclear, and there are no control data. To apply the results from these seven psychiatric patients to the general population and suggest that they provide evidence of useful predictive factors of relevance to clinical practice is misleading. Childhood sexual abuse is undoubtedly important; it is also topical. It may well be a marker for the much wider issue of childhood neglect, and a broader perspective needs to be adopted to enable us to begin to tackle the difficult issues (medical, social, political, and economic to name but four) of what to do about this. Nor is it yet clear what the best methods of treatment in adults are should childhood sexual abuse be uncovered. Is it, therefore, enough to say that the general practitioner has a key part to play in deflecting multiple referrals when it is far less clear what the general practitioner (and the physician, gynaecologist, surgeon, and psychiatrist) should do? The time for descriptive, poorly focused studies is past and there is an urgent need to embark on proper community surveys of childhood sexual abuse, its incidence, and effects as exemplified by some^{15,16} before any

further half truths grow up around this fraught subject.

RICHARD FRY

Department of Mental Health Sciences,
St George's Hospital,
London

RUTH WALLIS

Department of Public Health Medicine
and Epidemiology,
King's College Hospital,
London

- 1 Arnold RP, Rogers D, Cook DAG. Medical problems of adults who were sexually abused in childhood. *Br Med J* 1990;300:705-8. (17 March.)
- 2 Sedney M, Brooks B. Factors associated with a history of childhood sexual experience in a non-clinical female population. *Journal of the American Academy of Child Psychiatry* 1984;23:215-8.
- 3 Briere J, Runtz M. Symptomatology associated with childhood sexual victimisation in a non-clinical adult sample. *Child Abuse and Neglect* 1988;12:51-9.
- 4 Bagley C, Ramsay R. Sexual abuse in childhood: psychosocial outcomes and implications for social work practice. *Journal of Social Work and Human Sexuality* 1985;4:33-47.
- 5 Mullen PE, Romans-Clarkson SE, Walton VA, Herbison GP. Impact of sexual and physical abuse on women's mental health. *Lancet* 1988;341:5.
- 6 Shepherd M, Cooper B, Brown AC, Kalton G. *Psychiatric illness in general practice*. Oxford: Oxford University Press, 1966.
- 7 Croft-Jeffreys C, Wilkinson G. Estimated costs of neurotic disorder in UK general practice 1985. *Psychol Med* 1990;19:549-58.
- 8 Ribberink ML, Slurink GE, Everaerd W, Hanewald G. The incidence of sexual violence among women with gynecological complaints. *Journal of Psychosomatic Obstetrics and Gynaecology* 1989;10(suppl 1):25.
- 9 Finkelhor D, ed. *A sourcebook on child sexual abuse*. Beverly Hills, California: Sage, 1986.
- 10 Bachmann GA, Moeller TP, Benett J. Childhood sexual abuse and the consequences in adult women. *Obstet Gynecol* 1988;71:631-42.
- 11 Lindberg FH, Distad LJ. Post-traumatic stress disorders in women who experienced childhood incest. *Child Abuse and Neglect* 1985;9:329-34.
- 12 Gross RJ, Doerr H, Caldirola D, Guzinski G, Ripley H. Borderline syndrome and incest in chronic pelvic pain patients. *Int J Psychiatry Med* 1981;10:79-96.
- 13 Walker E, Katon W, Harrop-Griffiths J, Holm L, Russo J, Hickock LR. Relationship of chronic pelvic pain to psychiatric diagnoses and childhood sexual abuse. *Am J Psychiatry* 1988;145:75-80.
- 14 Raskin DE. Diagnosis in patients with chronic pelvic pain. *Am J Psychiatry* 1984;141:824.
- 15 Russell DEH. *The secret trauma: incest in the lives of girls and women*. New York: Basic Books, 1986.

SIR,—Dr R P Arnold and colleagues have described the incidence of medical problems in adults who were sexually abused during childhood.¹ They emphasise the increased incidence of somatisation in these adults. This concept is crucial for those who care for patients with epilepsy as illustrated by our recent experience of adolescent or young adult females presenting with ostensibly poorly controlled "epilepsy." We have now seen five such patients, who were eventually found to be suffering from psychogenic pseudoseizures rather than true epilepsy. Subsequent inquiry disclosed a history of sexual abuse during childhood in each patient. This was particularly clear in the two cases described below.

A 17 year old girl had been diagnosed as having epilepsy at the age of 13 years. She was admitted four years later as it had not been possible to achieve adequate control of the attacks as an out-patient. The attacks were of two types. The first consisted of sudden loss of consciousness with urinary incontinence but without a clonic phase. The second was characterised by a generalised clonic phase and urinary incontinence without complete loss of consciousness. She had stopped bedwetting only at the age of 5 years and had developed secondary enuresis at the age of 10. Full investigation including computed tomography showed no abnormality. Electroencephalography showed some bitemporal slow wave activity with occasional sharp wave transients. Ambulatory electroencephalography yielded normal results even during an "attack" associated with urinary incontinence. It therefore seemed that the attacks were predominantly psychogenic pseudoseizures,

and inquiries were made into her social background. These interviews disclosed that she had been sexually abused by her brother from the age of 8 and also in later years by her grandfather. After these disclosures and appropriate counselling the attacks largely resolved.

A 25 year old woman presented with a 12 year history of poorly controlled epilepsy. The attacks of altered consciousness with urinary incontinence had resolved at the age of 18 only to recur at the age of 20. They once again remitted but came back three years later, remaining poorly controlled until she was seen. The computed tomogram was normal. She reported attacks during 24 hours of ambulatory electroencephalography, but there were no corresponding correlates in the electroencephalogram. She complained of continuing attacks while in the ward, none of which were ever apparent to the nursing staff. We thought that these attacks were pseudoseizures. Detailed inquiry subsequently disclosed that she had been sexually abused at the age of 10 by an elderly male neighbour, who undressed her and lay on top of her simulating intercourse. At the age of 20 she met her fiancé and soon began having intercourse. Three years later she was assaulted by an uncle who penetrated her vagina digitally. The number of attacks fell abruptly after the disclosure of these facts.

These two cases illustrate the common clinical problem of differentiating non-organic pseudoseizures from true epilepsy. Detailed investigation and extensive manipulation of treatment was performed in both cases without obtaining adequate control of the attacks. Although the urinary incontinence was initially thought to favour an organic origin for the seizures, it was in fact enuretic rather than epileptic. Enuresis is common in sexually abused children.² Several interviews were needed to elicit the full background. The second was the most productive of factual information, the first tending to prepare the ground and build confidence to share distressing material. This experience shows the importance of a team approach, the physician, psychologist, and medical social worker combining to devise a suitable line of treatment.

Although we have reported on only two patients in detail, we suspect that this might be a more widespread problem than previously realised. Our cases show further aspects of the delayed sequelae of childhood sexual abuse during adult life and emphasise the importance of detecting such abuse at an early stage and depriving the abusers of access to the child at the earliest possible opportunity.

J D MITCHELL

H N GIBSON

Royal Preston Hospital,
Preston PR2 4HT

- 1 Arnold RP, Rogers D, Cook DAG. Medical problems of adults who were sexually abused in childhood. *Br Med J* 1990;300:705-8. (17 March.)
- 2 Skuse DH. Emotional abuse and neglect. *Br Med J* 1989;298:1692-4.

Attitudes to viability of preterm infants

SIR,—The study from the Trent region by Dr A C Fenton and colleagues of variability from unit to unit in attitudes to the viability of preterm infants showed that this variability had an impact on the reporting of perinatal deaths.¹ The authors concluded that there would be advantages in introducing the World Health Organisation's recommendation that all births and deaths of fetuses weighing ≥ 500 g should be registered.²

The Office of Population Censuses and Surveys discussed possible changes in criteria for registration of stillbirths in its white paper *Registration:*

Proposals for Change.³ This was published the previous month and reported on the results of a consultation for changes in the law about the registration of births, marriages, and deaths that were set out in an earlier green paper *Registration: a Modern Service*.⁴ The green paper asked for views about whether separate legislation should be introduced to change the current legal definition of stillbirth by reducing the current gestational age limit of 28 weeks.

The white paper reported that "There were 102 responses on this issue, both from individuals and from a wide variety of organisations in the medical field. Virtually all favoured gestational age as a means of defining stillbirths, with no significant support for birthweight."² This is hardly surprising as the possibility of using birth weight was not mentioned in the green paper. As a consequence it is unlikely that anyone without prior knowledge of World Health Organisation definitions would have put them forward as a basis for future legislation.

The differences reported in the Trent region exist to a much larger extent internationally as countries differ in the criteria used in their legislation about the registration of live births, stillbirths, and infant deaths. The World Health Organisation's recommendations were drawn up with the aim of encouraging countries to change their legislation so that comparable statistics can be produced. If England and Wales retain gestational age without birth weight as the sole criterion for stillbirth registration we will lose the opportunity to do so.

ALISON MACFARLANE

National Perinatal Epidemiology Unit,
Radcliffe Infirmary,
Oxford OX2 6HE

- 1 Fenton AC, Field DJ, Mason E, Clarke M. Attitudes to viability of preterm infants and their effect on figures for perinatal mortality. *Br Med J* 1990;300:434-6. (17 February.)
- 2 World Health Organisation. *International classification of diseases, injuries and causes of death*. Ninth revision. WHO: Geneva, 1978.
- 3 Office of Population Censuses and Surveys. *Registration: proposals for change*. London: HMSO, 1990. (Cmd 939.)
- 4 Office of Population Censuses and Surveys. *Registration: a modern service*. London: HMSO, 1988.

Community care

SIR,—Further to Dr Michael Ashley-Miller's editorial,¹ I should like to describe the experience of Leicestershire District Health Authority in one aspect of providing community care. For three years we have been offering a medical examination to handicapped adults. Long serving clinical medical officers, who are members of community mental health teams and have experience of screening and surveillance of the developing child, do the medicals. Liaison meetings are held with the family practitioner committee and with representatives of the relevant paramedical professions.

The known population of handicapped adults in Leicestershire is 2100. With 320 sessions a year we offer a three yearly medical, with earlier review if necessary. These take place countywide, in day centres, technical colleges, and long stay hospitals.

A letter of invitation is sent to the patients,² and carers are invited to attend. The patient's general practitioner is notified of all baseline data and any abnormal findings. (Inpatients of more than two years' duration have no general practitioner—this is seen as a disadvantage.) Medical advice is offered to staff when this is in the interest of the patient. All data are entered on to the computer and form part of the mental handicap register, which is to be used for long term planning.

An evaluation of the service—in terms of referrals—was carried out on 10% of 1340 casenotes. In 134 patients, 59 referrals were made: a 44% referral rate. Some patients were referred to more than one