Type of coronary event	No of events (treated/control)	Sum of observed-expected events*	Overall odds ratio (95% confidence interval)	p Value
Non-fatal	368/495	-63.5	0.73 (0.64 to 0.84)	0.00
Fatal	169/197	-14.0	0.85 (0.69 to 1.05)	0.06
All	523/670	-73.5	0.77 (0.68 to 0.86)	0.00

*The sum of observed-expected events in the combined trials, as used in the meta-analysis, is half of the actual difference in events (treated-control) when the treated and control groups are of equal size.

disease events.¹ This is illustrated by a further meta-analysis, by identical methods, of the same six trials (table). Even if overall mortality was not significantly reduced a reduced morbidity from cardiac disease remains as a net gain and is a sufficient goal.

The debate about mortality should not distract attention from the fact that both primary and secondary trials of cholesterol lowering have successfully shown reductions in their primary end point—namely, all coronary heart disease events.² Primary prevention trials have been more effective in reducing non-fatal than fatal coronary heart disease events. Participants experienced many more non-fatal than fatal events, hence the number of non-fatal events avoided is correspondingly larger.

Apart from the evident health benefits to individual patients of reducing morbidity from coronary heart disease, the savings in the direct costs of health care are potentially large and exceed those associated with reducing mortality from coronary heart disease. In 1985 the estimated direct cost of coronary heart disease to the NHS (England and Wales) was £389.9m, and in 1984 there were 178 870 inpatient cases of coronary heart disease in England alone (Coronary Prevention Group and British Heart Foundation database). The estimated 7.2 million Americans with coronary heart disease account for over 27 million office visits and over two million admissions to hospital a year, and America's direct costs amounted to an estimated \$15.3 billion in 1987.³

Substantial further gains in life expectancy are unlikely in developed societies, and the primary purpose of risk reduction then becomes to compress morbidity into the last few years of life and to improve the quality and vigour of life.⁴ That a decrease in overall mortality has not thus far been shown in clinical trials does not argue against pursuing a reduction in morbidity from coronary heart disease.

JACQUES E ROSSOUW
BASIL M RIFKIND

National Institutes of Health,
Bethesda,
Maryland 20892,
United States

- Muldoon MF, Manuck SB, Matthews KA. Lowering cholesterol concentrations and mortality: a quantitative review of primary prevention trials. *Br Med J* 1990;301:309-14. (11 August.)
- Yusuf S, Wittes J, Friedman L. Overview of results of randomized clinical trials in heart disease. 2. Unstable angina, heart failure, primary prevention with aspirin, and risk factor modification. *JAMA* 1988;260:2259-63.
- National Heart, Lung, and Blood Institute. *Morbidity from coronary heart disease in the United States*. Bethesda: NHLBI Education Program Information Center, 1990.
- Fries JF, Green LW, Levine S. Health promotion and the compression of morbidity. *Lancet* 1989;i:481-3.

AUTHORS' REPLY.—Dr Canter points out that several of the trials that we analysed included some subjects with coronary heart disease and suggests that these investigations therefore should not have been grouped with the other primary prevention trials. All six studies, however, met our definition of a primary prevention trial, which specified that "... criteria for participants' eligibility did not include a history of coronary heart disease."

Although some of the trials excluded people with clinical evidence of heart disease, it is erroneous

to conclude that subjects in these studies were therefore free of coronary atherosclerosis; the prevalence of silent or undiagnosed coronary heart disease can be conservatively estimated at 20% in men typical of these trials—50 years old with an average cholesterol concentration of 6.6 mmol/l. Hence no "genuine" primary prevention trials of cholesterol reduction exist, and though some subject populations were more disease free than others, each of the six studies approximates a primary prevention trial.

Dr Canter also points out, as did we, that the fairly short duration of these trials leaves many questions unanswered regarding the potential benefits (or hazards) of more protracted treatment. Though understandably constrained by the high cost of longer trials, this is perhaps the most important limitation of the data available. It is unclear, however, that a doubling or tripling of the duration of treatment would have been sufficient. No trend toward reduced total mortality was seen in the primary prevention trials reviewed. Moreover, in the coronary drug project, a secondary intervention trial with substantially higher mortality from coronary heart disease than primary prevention studies, a reduction in total mortality was observed at only the 15 year follow up.¹

Dr Ravnskov notes that in two trials random assignment of subjects to treatment and control conditions failed to create perfectly comparable groups. He also believes that the inclusion of these studies led us to overestimate the benefits of cholesterol reduction on mortality from coronary heart disease. In practice randomisation cannot completely ensure equal distribution of all potentially confounding variables, but neither can relatively minor inequalities in group characteristics be cited to invalidate study findings.

Finally, we agree completely with Drs Rossouw and Rifkind regarding the great importance of the reduction in morbidity from coronary heart disease. Indeed, the well documented prevention of heart attacks is the principal reason for widespread acceptance and popularity of cholesterol lowering regimens. Statements regarding "compression of morbidity," on the other hand, may be premature. The trials did not assess morbidity (from, for example, psychiatric illness) that might be related to the increased mortality from suicides and violence; similarly, the incidence of cancer was rarely reported. Thus whether primary prevention through cholesterol reduction results in a "net gain," in terms of either morbidity or mortality, remains to be determined.

MATTHEW F MULDOON
STEPHEN B MANUCK
KAREN A MATTHEWS

University of Pittsburgh School of Medicine,
Pittsburgh, PA 15260,
United States

- Canner PL, Berge KG, Wenger NK, et al. Fifteen year mortality in coronary drug project patients: long-term benefit with niacin. *J Am Coll Cardiol* 1986;8:1245-55.

Hereditary (primary) haemochromatosis

SIR.—As early venesection rescues genetically susceptible people from the effects of iron overload, potentially returning their life expectancy to that of

the general population, Dr Niall D C Finlayson rightly advises screening of their asymptomatic close relations.¹

Blood tests provide useful indications, but measurement of the iron concentration in the liver gives the most unequivocal evidence. This does not necessarily require liver biopsy: dual energy computed tomography can be used.² The same tissue slice is scanned at two energy levels and the difference in the computed tomographic number calculated: the dual energy differential. Iron, with a higher atomic number than most other elements in the liver, will absorb relatively more at lower energy levels whereas the absorption by other substances in the liver is little changed. The difference in absorption at the two energy levels is therefore largely attributable to the presence of iron and correlates closely with the iron concentration in biopsy specimens (correlation coefficient 0.993), thus obviating the need for biopsy in the relatives of patients with hereditary haemochromatosis.

Chapman *et al* found that computed tomography correlated more closely with the chemical estimation of iron concentrations in liver than serum ferritin concentration.³ Studies have shown that serum ferritin concentration may be normal in patients with precirrhotic haemochromatosis³ and may be increased by alcohol abuse; these factors should not affect estimation by computed tomography.

Unfortunately, this potentially useful technique is not generally available despite the wide availability of computed tomography as most scanners do not have the dual energy facility. Slight modifications of scanners would provide a non-invasive accurate method of estimating the hepatic iron concentration when screening for hereditary (primary) haemochromatosis.

IAN KELLY
ROBERT DICK

Royal Free Hospital, London NW3 2QG

- Finlayson NDC. Hereditary (primary) haemochromatosis. *Br Med J* 1990;301:350-1. (18-25 August.)
- Chapman RWG, Wilham G, Bydder G, Dick R, Sherlock S. Computed tomography for determining liver iron content in primary haemochromatosis. *Br Med J* 1980;280:440-2.
- Wands JR, Rowe JA, Mezey SE, et al. Normal serum ferritin concentration in precirrhotic hemochromatosis. *N Engl J Med* 1976;294:302-5.

Symptoms of low blood pressure

SIR.—Dr Simon Wessely and colleagues were surprised to find a relation between low blood pressure and tiredness¹ and perhaps it is the nature of their training that urged them to argue against accepting their results on the basis that there was no clear cut pathophysiological explanation. I believe that there are reasons for disputing that conclusion.

Blood pressure reflects the degree of peripheral resistance against which the heart has to work. In hypotension due to autonomic failure pooling of blood in the veins of the lower limbs seems to be corrected when venous tone is increased by drugs such as inhaled ergotamine.² When venous pooling occurs the reduced venous return would be associated with a lowered intracardiac packed cell volume, which would not only have an effect on peripheral resistance but result in suboptimal delivery of oxygen to the tissues. Therefore the tiredness of hypotension could be the consequence of inadequate energy generation, a proposal that is in accord with physiological concepts of physical fatigue. Griffiths stated that "Fatigue is probably caused by both an oxygen deficiency and an accumulation of metabolites."³ Schmidt and Thews wrote, "Physical fatigue results from changes in skeletal muscle in longterm performance, the depletion of the energy stores and accumulation of

lactic acid (the fatigue substance) which reduce performance capacity.^{7,8} Incubation with physiological concentrations of lactic acid reduced the filterability of erythrocytes,⁹ and more than 40 years ago high blood concentrations of lactate after moderate exercise in subjects with neurocirculatory asthenia were considered to suggest deficient aerobic metabolism.⁶

In New Zealand members of the Dunedin Myalgic Encephalomyelitis Support Group were found to have reduced blood filterability when they were acutely unwell,⁷ and analysis of red cell shape has shown that subjects with the symptoms of myalgic encephalomyelitis have appreciably increased percentages of non-discocytic red cells.^{8,9} As such cells will impair capillary blood flow the chronic tiredness of subjects with the symptoms of myalgic encephalomyelitis and a reduced percentage of normal discocytes is probably explicable in terms of inadequate energy generation.¹⁰

Observations made on red cell shape in other disorders that have chronic tiredness as a common symptom provide support for the statement by Ffrench that "there is no doubt that oxygen lack is the first cause of tissue cell exhaustion which is manifested early by clinical tiredness."¹¹ Without disputing that central fatigue exists in some patients with chronic tiredness it is suggested that studies of those factors that influence capillary blood flow and oxygen delivery could be of value in separating those whose problem is an affective disorder.

L O SIMPSON

University of Otago,
Dunedin, New Zealand

- 1 Wessely S, Nickson J, Cox B. Symptoms of low blood pressure: a population study. *Br Med J* 1990;301:362-6. (18-25 August.)
- 2 Biaggioni I, Zygmunt D, Haile V, Robertson D. Pressor effect of inhaled ergotamine in orthostatic hypotension. *Am J Cardiol* 1990;65:89-92.
- 3 Griffiths M. *Introduction to human physiology*. New York: McMillan, 1981: 73-4.
- 4 Schmidt RF, Thews G. *Human physiology*. 2nd ed. Berlin: Springer-Verlag, 1989:656-9.
- 5 Shand BI. Changes in blood rheology induced by lactic acid. *Proceedings of the University of Otago Medical School* 1986;64: 71-2.
- 6 Cohen ME, Consolazio F, Johnson RE. Blood lactate response during moderate exercise in neurocirculatory asthenia, anxiety neurosis or effort syndrome. *J Clin Invest* 1947;26: 339-42.
- 7 Simpson LO, Shand BI, Olds RJ. Blood rheology and myalgic encephalomyelitis: a pilot study. *Pathology* 1986;18:190-2.
- 8 Simpson LO. Blood from healthy animals and humans contains nondiscocytic erythrocytes. *Br J Haematol* 1989;73:561-4.
- 9 Simpson LO. Nondiscocytic erythrocytes in myalgic encephalomyelitis. *N Z Med J* 1989;102:126-7.
- 10 Wiles CM, Jones DA, Edwards RHT. Fatigue in human metabolic myopathy. *Ciba Found Symp* 1981;82:264-82.
- 11 Ffrench G. The clinical significance of tiredness. *Can Med Assoc J* 1960;82:665-71.

Choice cuts for patients with AIDS?

SIR,—We share the concern of Dr Chris Bentley and Professor Michael W Adler that the NHS review may threaten the quality and quantity of provision of services for genitourinary medicine.¹

Countries, such as the United States, that cut their services for sexually transmitted diseases under the guise of cost effectiveness when these diseases were decreasing in the mid-1950s have later had a serious increase in cases. In Britain a case was fought for maintaining the services for sexually transmitted diseases at that time.² This has been reflected in the decreasing prevalence of bacterial diseases, in particular syphilis and gonorrhoea, and at the same time we are fortunate that the infrastructure for the prevention and treatment of the increasingly prevalent viral diseases, such as herpes, warts, and HIV, remains intact. This illustrates the problem of subjecting services to market forces. Success of preventive measures is difficult to quantify, and a decreasing incidence of sexually transmitted diseases may be

construed as a signal to cut resources when it should be further consolidated.

The uncertainty of the future funding of the services for sexually transmitted diseases in the light of the provider-purchaser role is already having serious consequences in our health authority. Services that have revenue consequences, although funded by the current AIDS monies, may not be allowed to proceed because of this uncertainty. This has caused alarm among us and has implications for services throughout Britain. We hope that urgent action is taken centrally to resolve this issue and to ensure that funding for our confidential and open access service is maintained to prevent and treat sexually transmitted diseases, including AIDS.

B T GOH
G E FORSTER

Royal London Hospital, London E1 1BB

- 1 Bentley C, Adler MW. Choice cuts for patients with AIDS? *Br Med J* 1990;301:501-2. (15 September.)
- 2 King AJ. "These dying diseases." *Venerology in decline? Lancet* 1958;i:651-7.

SIR,—Some changes have occurred since Dr C Bentley and I submitted our editorial on the possible effects of the NHS and Community Care Act 1990 on services for genitourinary medicine and for patients with AIDS or infected with HIV.¹

Although the government was not willing to amend the act to reflect the special circumstances of self referral services, it did clarify the position with a series of statements in the House of Lords. The key speech was made by Baroness Blatch in response to the amendment moved by Lord Kilmarnock, chairman of the all party parliamentary group on AIDS.²

The first important point concerns funding. It was stated that the government expects regional health authorities' funding of district health authorities to reflect usage of self referral services. Therefore "funding . . . will have to take into account . . . self-referral services for all-comers, and not just for residents." As far as we are aware, this is the first time that this has been stated.

The second point refers to the issue of confidentiality. Here the baroness stated that "treatment will begin and end in one local authority. There will not be a cross-boundary flow of information back to the home authority of the patient who is receiving services."

These important statements are very welcome and reassuring and should be given wider circulation. But many questions still remain. Do these statements, for example, equally apply to patients with AIDS admitted for treatment or, as with the accident and emergency model, will these episodes be separately billed? If they are billed separately how will confidentiality be maintained?

Clarification on such issues is urgently needed as 1 April draws ever closer, but we are pleased that the government has started to appreciate the problems that the act presents and remain hopeful that our concerns will be tackled.

MICHAEL ADLER

University College and Middlesex School of Medicine,
London W1N 8AA

- 1 Bentley C, Adler MW. Choice cuts for patients with AIDS? *Br Med J* 1990;301:501-2. (15 September.)
- 2 Blatch, Baroness. Third reading of the NHS and Community Care Bill. *House of Lords Official Report (Hansard)* 1990 June 25;520:cols 1356-7. (No 107.)

Osteoporosis after 60

SIR,—Dr Roger Smith is correct to stress that the treatment of postmenopausal osteoporosis must be tailored to the individual but he is, in our view, incorrect to suggest that oestrogens are no longer first choice treatment for most women over the age of 60.¹

There are three problems to resolve. Firstly, is oestrogen therapy effective in preventing or reversing bone loss in this age group? Secondly, are the side effects of oestrogen therapy unacceptable in the over 60s? Thirdly, are the non-skeletal benefits of oestrogen replacement such that hormonal therapy is greatly to be preferred to sodium fluoride or diphosphonates?

Civitelli *et al*² have shown a 6% increase in vertebral bone density after oral therapy in osteoporotic women, and we have recently shown a similar increase in vertebral bone density after one year of oestrogen implants regardless of age, menopausal age, and initial bone density.³ There was, however, a significant correlation between the increase in vertebral bone density and the serum oestradiol concentrations achieved. Older women or those with established osteoporosis will have a substantial increase in bone density, and presumably bone strength, if adequate plasma oestradiol concentrations are achieved.

The side effects of regular uterine bleeding, which may be heavy or painful or coexist with progestogenic symptoms or symptoms of the postmenopausal syndrome,⁴ may be an obstacle in older women, but with counselling most will accept these in return for the benefits of increased bone density. Currently research is directed towards developing regimens of continuous combined oestrogen and progestogen preparations that do not produce vaginal bleeding.^{5,6} Breast discomfort is another occasional complication, but the return of sexuality after a quiet 10 years is a more common and more disturbing problem, which is often not expressed to the doctor.

The third question reflects the eccentric view that women are not merely skeletons and that the beneficial effects of oestrogens on general well-being must be considered. Long term oestrogen therapy is associated with a decreased overall mortality and decreased incidence of heart attacks and strokes.⁷ The fear of breast cancer among both patients and doctors is real enough, but published reports concerning the incidence of breast cancer in oestrogen users are confused and inconclusive. The only studies that consider the mortality from breast cancer in these patients show a reduction.^{8,9} This may be seen as encouraging when assessing the risk-benefit of long term oestrogen use in women aged over 60.

JOHN STUDD
TIMOTHY GARNETT
MICHAEL SAVVAS

King's College Hospital,
London SE5 9RS

- 1 Smith R. Osteoporosis after 60. *Br Med J* 1990;301:452-3. (8 September.)
- 2 Civitelli R, Agnusdei D, Nardi P, Zacchei F, Avioli LV, Gennari C. Effects of one-year treatment with estrogens on bone mass, intestinal calcium absorption, and 25-hydroxy-vitamin D-1α-hydroxylase reserve in post-menopausal osteoporosis. *Calcif Tissue Int* 1988;42:77-86.
- 3 Studd JWW, Savvas M, Fogelman I, Garnett T, Watson NR, Cooper D. The relationship between plasma estradiol and the increase in bone density in postmenopausal women following treatment with subcutaneous hormone implants. *Am J Obstet Gynecol* (in press).
- 4 Magos AL, Brewster E, Singh R, O'Dowd T, Brincat M, Studd JWW. The effects of norethisterone in post-menopausal women on oestrogen replacement therapy, a model for pre-menstrual syndrome. *Br J Obstet Gynaecol* 1986;93: 1290-9.
- 5 Magos AL, Brincat M, Studd JWW. Endometrial and bleeding response to continuous estrogen and progestogen therapy in the postmenopausal woman. *Obstet Gynecol* 1985;66:836-7.
- 6 Leather A, Studd JWW. Commentary: can the withdrawal bleed following oestrogen replacement therapy be avoided? *Br J Obstet Gynaecol* (in press).
- 7 Ross R, Pyke MC, Mack TM, Henderson BE. Oestrogen replacement therapy and cardiovascular disease. In: Drife JO, Studd JWW, eds. *HRT and osteoporosis*. Berlin: Springer-Verlag, 1990:209-22.
- 8 Hunt K, Vessey M, MacPherson K, Coleman M. Long-term surveillance of mortality and cancer incidence in women receiving hormone replacement therapy. *Br J Obstet Gynaecol* 1987;94:620-35.
- 9 Bergkvist L, Adami H, Persson I, Berstrom R, Krusemo U. Prognosis after breast cancer diagnosis in women exposed to estrogen and estrogen-progestogen replacement therapy. *Am J Epidemiol* 1989;130:221-8.