- All letters must be typed with double spacing and signed by all authors.
- No letter should be more than 400 words.
- For letters on scientific subjects we normally reserve our correspondence columns for those relating to issues discussed recently (within six weeks) in the *BMJ*.
- We do not routinely acknowledge letters. Please send a stamped addressed envelope if you would like an acknowledgment.
- Because we receive many more letters than we can publish we may shorten those we do print, particularly when we receive several on the same subject.

What's in a name?

SIR,—As medical treatment has become more sophisticated, there has been an increasing need to assess both its effectiveness and its cost. Many techniques have evolved for this purpose, including the medical staff round, the surgical “death and complications” meeting, and the pathologist’s quality control review. Most of these approaches have lacked an effective mechanism for translating the lessons that are learnt into procedures for better management. It is to the credit of the most recent techniques of medical audit that they apply themselves firmly to this follow through problem.

Prompted by the medical profession in this country, the government has now asked all doctors to undertake audit and has provided funds for this purpose. Despite the strong support of the royal colleges, however, this approach has had a mixed reception. An often expressed fear is that if current American audit systems are adopted uncritically the quality of care will suffer because of an undue emphasis on cost cutting and fault finding.

Many of us would argue that medical audit, as it has developed in this country, has set new, clinically oriented goals that may help to satisfy the needs of management but are not a part of it. To the uninitiated the word audit seems to belie this claim. Audit is “an official examination of accounts with verification by reference to witnesses and vouchers” (*Shorter Oxford English Dictionary*). The processes that we seek to encourage are more properly seen as clinical review (review is “the act of looking over something (again) with a view to correction or improvement”).

Since I have started talking at audit meetings about clinical review instead of medical audit I have found it much easier to make new friends.

M H LESSOF

Guy's Hospital,
London SE1 9RT

Guidelines for management of asthma

SIR,—I agree with Dr P John Rees¹ that the guidelines issued by the British Thoracic Society and others^{2,3} deserve careful consideration from all those who care for patients with asthma but am uneasy about several of the directions in the section on acute severe asthma.

In the introduction it is stated that the directions are designed to help doctors “in the home,” as well as in hospital, but except for the panel entitled “Criteria for emergency referral to hospital” most of the directions are geared to hospital practice. It would have been preferable to give separate directions for general practitioners, which could

have featured a list of the drugs and equipment (including a portable oxygen cylinder and a nebuliser) that a general practitioner should carry in his or her car and detailed directions on how best to treat an acute attack of asthma with the drugs and equipment specified on the list.

I cannot see any point in keeping a resuscitation box in the home of a patient liable to catastrophic sudden severe asthma as such a box would normally be carried in the doctor’s car to deal with cardiac arrest, and it would be unrealistic to believe that relatives, even if provided with a box, could resuscitate an asthmatic patient in that condition. Indeed, hypoxic cardiac arrest in asystole, which is the most common terminal event in acute severe asthma, can be treated effectively only in hospital as external cardiac compression cannot maintain a circulation in patients with severely overdistended lungs. It is therefore essential that patients in this high risk category should have an oxygen cylinder and a nebuliser in their homes and are able to get into hospital as quickly as possible if their asthma worsens. Although the guidelines mention “a management plan that is mutually agreed on by the patient, the general practitioner, and the consultant,” presumably to expedite admission, no reference is made to the emergency self admission scheme⁴ that has proved so successful in Edinburgh and elsewhere. The advice to go to the nearest hospital could be a recipe for disaster unless that hospital knows the patient well, is prepared to admit him or her without demur, and has facilities immediately available for intubation and ventilation.

The statement that “antibiotics are not indicated unless there is evidence of a bacterial infection” must be challenged. The most reliable clinical indicator of bacterial infection is purulent sputum, but that is hardly a useful criterion as patients with acute severe asthma can seldom produce any sputum at all. Since bacterial infection is always a potential complication, what harm can it do to prescribe an antibiotic in these circumstances?

“Do not attempt intubation until the most expert available doctor (ideally an anaesthetist) is present” implies that a junior doctor in the accident and emergency department may be allowed to attempt intubation after an appropriate expert has arrived on the scene even if that expert is not an anaesthetist. Tracheal intubation in a patient with acute severe asthma is always difficult and hazardous and causes great distress to an un-anaesthetised subject. It would have been safer, and less ambiguous, to have stated that intubation should be performed only by an anaesthetist except when it is desperately required and no anaesthetist is available.

IAN W B GRANT

Kirknewton,
West Lothian EH27 8EA

1 Rees PJ. Guidelines for the management of asthma. *BMJ* 1990;301:771-2. (6 October.)

2 British Thoracic Society, Research Unit of the Royal College of Physicians of London, King's Fund Centre, National Asthma Campaign. Guidelines for management of asthma in adults. 1. Chronic asthma. *BMJ* 1990;301:651-3. (29 September.)
3 British Thoracic Society, Research Unit of the Royal College of Physicians of London, King's Fund Centre, National Asthma Campaign. Guidelines for management of asthma in adults. 2. Acute severe asthma. *BMJ* 1990;301:797-800. (6 October.)
4 Crompton GK, Grant IWB, Bloomfield P. Edinburgh Asthma Admission Service: report on 10 years' experience. *BMJ* 1979;iii:1199-201.

AUTHOR'S REPLY.—The guidelines need to be concise to get their important messages across clearly and to the widest possible audience. Everyone with an interest in asthma is likely to find some aspects that they disagree with or that they consider have the wrong emphasis, and Dr Grant has criticised several aspects of the acute management plan. Catastrophic, sudden severe asthma attacks can develop over minutes or hours, and the means to begin treatment need to be immediately available to the patient not in the general practitioner’s car. Patient and relatives should be trained to begin treatment with oxygen, nebulised or injected β agonist, and steroids. Such immediate intervention can be life saving in this condition whereas waiting for the most prompt general practitioner might prove fatal.

The emergency self admission scheme has indeed proved successful in Edinburgh and elsewhere, and its wider application would certainly be helpful. Dr Grant, however, quotes rather selectively in choosing “go to the nearest hospital.” The guidelines actually state “go to the nearest hospital as previously agreed with the general practitioner,” emphasising the importance of choosing an adequately equipped casualty department with which the patient has a prior arrangement, as noted by Dr Grant.

Most infections that precipitate asthma attacks are viral. The diagnosis of bacterial infection may be difficult in patients with acute asthma, but it seems correct to try to target antibiotics to the small group most likely to benefit. This will still mean that they are prescribed more often than they are needed.

Dr Grant is right to stress the difficulties of intubation in acute asthma. The statement on intubation comes in a series of direct instructions on management of the acute attack. In this context it seems unlikely that “do not attempt intubation until the most expert available doctor (ideally an anaesthetist) is present” could really be interpreted as letting anybody have a go as long as the anaesthetist was in the room.

JOHN REES

United Medical and Dental Schools,
London SE1 9RT

SIR.—As a consultant in accident and emergency I think that there is one cause for concern in the