meretricious publication, which overemphasises throughput and gives scant consideration to quality (the priority needed for NHS patients) and the wider clinical obligations of radiologists, oncologists, and their departments.

D J STOKER
C H PAINE

Royal College of Radiologists,
London W1N 3DG

1 Timperley WR. Selling the pathology service. *BMJ* 1990;301:828. (13 October.)

Hirsutism

SIR,—Dr Gerard S Conway and Professor Howard S Jacobs emphasise the use of cyproterone acetate in combination with oestrogen for treating hirsutism.¹ In our experience Dianette, which contains 35 µg of ethinyloestradiol and only 2 mg of cyproterone acetate, is relatively ineffective, and it is in any case not licensed for long term maintenance therapy. Marvelon, a licensed oral contraceptive containing 30 µg ethinyloestradiol and 150 µg desogestrel is at least as effective a treatment for hirsutism as Dianette.²

Higher dose combinations of cyproterone acetate, which are more effective than Dianette and probably than Marvelon, are not licensed for use in hirsutism. The major disadvantage of cyproterone acetate is its long half life (two to five days in the deep compartment).³ This means that to produce regular menstruation it must be given in a regimen of 10 days out of 28, with ethinyloestradiol for 21 days.⁴ This so called reverse sequential regimen leads to wide fluctuations in both circulating concentrations of the antiandrogen and gonadotrophin suppression. Cyproterone acetate is also a weak glucocorticoid and leads to adrenocortical suppression at high doses.⁵ The rather complicated regimen leads to poor compliance. Furthermore, cyproterone acetate has been shown to cause liver tumours in rats, and a warning to this effect is given in the Dianette data sheet.

Spironolactone is presently the only serious alternative antiandrogen for severe hirsutism. Its efficacy in a dose of 50 to 200 mg daily has been shown in several studies.^{6,7} Because it has no progestational activity it can be given continuously in combination with an oral contraceptive such as Marvelon, with a likely additive effect as Marvelon probably acts principally by suppressing endogenous ovarian androgen production.

Dr Conway and Professor Jacobs dismissed spironolactone on the basis of a circular letter from the Committee on Safety of Medicines, which limited licensed indications notably for hypertension.⁸ Doubt over the long term safety of spironolactone was raised because of the development of mammary tumours, liver tumours, and myelocytic leukaemia in rats given potassium canrenoate, an aldosterone antagonist to which spironolactone is chemically related and with which it shares a common metabolite (canrenone). In long term animal studies there was no evidence of a tumourigenic or carcinogenic effect in rats given spironolactone.⁹

Unfortunately, unless unlicensed drugs are used there is no effective treatment for women with severe and often disabling hirsutism. The drug companies are unlikely to seek to extend the licences for drugs long out of patent. Meanwhile those of us concerned with treating such women will continue to use these drugs, while conducting the much needed long term efficacy and safety studies and fully discussing the safety issues with our patients.

JOHN O'DRISCOLL

The Skin Hospital,
Salford M60 9EP

DAVID C ANDERSON

Hope Hospital,
Salford M6 8HD

- Conway GS, Jacobs HS. Hirsutism. *BMJ* 1990;301:619-20. (29 September.)
- Dewis P, Petros P, Newman M, Anderson DC. The treatment of hirsutism with a combination of desogestrel and ethinyl oestradiol. *Clin Endocrinol* 1985;22:22-36.
- Gerhards E, Ropke H, Schulze PE. Biodynamics of cyproterone in man following oral and intravenous administration. *Acta Endocrinol* 1970;64:28-52.
- Hammerstein J, Meckies J, Leo-Rossberg I, Molty L, Zielske F. Use of cyproterone acetate in the treatment of acne hirsutism and virilism. *J Steroid Biochem* 1975;6:827-36.
- Hague WM, Munro DS, Sawers RS, Duncan SLB, Honour JW. Long-term effects of cyproterone acetate on the pituitary adrenal axis in adult women. *Br J Obstet Gynaecol* 1982;89:981-4.
- Boiselle A, Tremblay RR. New therapeutic approach to the hirsute patient. *Fertil Steril* 1979;32:276-9.
- Barth JH, Cherry CA, Wojnarowska F, Dawber PRP. Spironolactone is an effective and well tolerated systemic antiandrogen therapy for hirsute women. *J Clin Endocrinol Metab* 1989;68:966-70.
- Committee on Safety of Medicines. Spironolactone. *Current Problems* 1988;21(Jan).
- Wagner BM. Long-term toxicology studies of Spironolactone in animals and comparison with potassium canrenoate. *Journal of Drug Development* 1987;1(suppl 2):7-11.

SIR,—We agree with Dr Gerard S Conway and Professor Howard S Jacobs that current treatments for hirsutism are associated with considerable complications or theoretical risks.¹ Weight loss is not an option for normal weight women with the polycystic ovary syndrome, and though cosmetic measures are of value in some women, they are expensive for prolonged treatment and their provision by the NHS is inconsistent and of low priority.

The authors suggest that Dianette may not be sufficient for reversing excess hair growth. We have recently assessed Dianette as a first line treatment of hirsute women with the polycystic ovary syndrome. Full results will be reported elsewhere. Briefly, 28 oligomenorrhoeic hirsute women with polycystic ovaries, as shown by ultrasonography, and with a median score on the Ferriman-Gallwey index of 13 (range 5-32) were treated for up to one year with Dianette.

We found significant improvements in facial and chest hirsutism by 24 weeks' treatment and in abdominal and leg hirsutism by 36 weeks, with a corresponding decrease in the median Ferriman-Gallwey index.² Two women withdrew from the study prematurely because of lack of response (initial Ferriman-Gallwey scores 15 and 22), and four others showed no overall improvement. There were no changes in glucose tolerance or mean insulin concentrations in response to an oral glucose load after six months' treatment with Dianette despite an appreciable decrease in the mean concentration of free testosterone. Dianette was well tolerated overall.

The disassociation between rapid improvement of biochemical hyperandrogenism and slower clinical response to antiandrogen treatment is well recognised and does not seem to be influenced by the drug used. Dianette is of value in the initial treatment for hirsutism in women with the polycystic ovary syndrome, particularly for those women with lesser degrees of hirsutism, in whom avoidance of higher dose cyproterone acetate may lessen side effects.

I GOLLAND

St Mary's Hospital,
Portsmouth PO3 6AD

M ELSTEIN

University Hospital of South Manchester,
Manchester M20 8LR

- Conway GS, Jacobs HS. Hirsutism. *BMJ* 1990;301:619-20. (29 September.)
- Ferriman D, Gallwey J. Clinical assessment of body hair growth in women. *J Clin Endocrinol Metab* 1961;21:1440-7.

AUTHORS' REPLY,—Dr O'Driscoll and Professor Anderson seem to have misunderstood our reference to drug licensing. Our aim was to note that although several preparations are used to treat hirsutism, only Dianette is licensed. We did not intend to suggest that other drugs ought not to

be used merely because they were not licensed for the indication. This is in fact a common misunderstanding of the role of the Committee on Safety of Medicines, which is to advise the Licensing Authority on the Safety of Medicines, not to determine the practice of medicine.

To our knowledge, there have been no randomised trials comparing the effectiveness of spironolactone with that of combined antiandrogen drugs. We have found spironolactone less effective than cyproterone acetate, and we cannot help noticing that most of the published reports proposing its use originate from countries where cyproterone acetate is not yet available.

In terms of efficacy we consider the reversed sequential regimen of cyproterone acetate and ethinyloestradiol the appropriate first line treatment for moderate to severe hirsutism. As Dr Golland and Professor Elstein show, patients with milder disease may benefit from milder treatment. Whether, in terms of safety, this approach is optimal remains to be established, but newer drugs (such as flutamide) and newer strategies (such as treatment with inhibitors of 5 α -reductase) may eventually offer safer treatment.

G S CONWAY

H S JACOBS

University College and Middlesex School of Medicine,
London W1N 8AA

SIR,—Although Dr Gerard S Conway and Professor Howard S Jacobs allude to defects in adrenal steroid synthesis as a cause of hirsutism,¹ perhaps more emphasis should have been given to the non-classic forms of congenital adrenal hyperplasia, which are present in an important subgroup of young women presenting with hirsutism and other clinical manifestations of hyperandrogenism.

Five enzymes are necessary for the conversion of cholesterol to cortisol, and partial deficiencies of three of these (21-hydroxylase, 3 β -hydroxysteroid dehydrogenase, 11 β -hydroxylase) are well recognised.² Partial 21-hydroxylase deficiency is the most widely studied and may be the commonest, occurring in approximately 0.3% of the general caucasian population; there is considerable ethnic variation such that about 3% of east European Jews are affected.³

The condition most commonly presents after the menarche with symptoms of hyperandrogenism with and without menstrual disturbance,^{4,5} and may be responsible for up to 6% of unselected hyperandrogenic women presenting to hospital clinics.⁶ It is clinically indistinguishable from the polycystic ovary syndrome^{6,7} and, indeed, the appearances of polycystic ovaries on ultrasonography are often seen in non-classic 21-hydroxylase deficiency.^{8,9} Standard measurements of serum concentrations of androgens such as testosterone, androstenedione, and dihydroepiandrosterone sulphate cannot discriminate between the two conditions,^{6,9} but non-classic 21-hydroxylase deficiency can be readily distinguished by an exaggerated serum 17-hydroxyprogesterone response (measured at 60 minutes) to 0.25 mg soluble tetracosactrin administered intravenously.¹⁰ Unstimulated morning measurements of serum 17-hydroxyprogesterone concentrations may also be diagnostic, but values in the normal range cannot be used to exclude the condition.¹¹

The importance of diagnosing non-classic 21-hydroxylase deficiency lies in the excellent prognosis on glucocorticoid replacement therapy. A small dose of dexamethasone (0.25-0.5 mg) taken at bedtime leads to suppression of adrenal androgens, resolution of clinical signs of hyperandrogenism, and resumption of normal menses and fertility.^{10,12} At present, the incidence of non-classic 21-hydroxylase deficiency among women presenting with symptoms of hyperandrogenism to endocrine and gynaecology clinics in the United Kingdom is unknown. The cost effectiveness of screening such patients for the condition

is therefore unclear, but screening certainly merits consideration given the cost of long term antiandrogen therapy for hirsutism compared with that of dexamethasone. A diagnosis of 21-hydroxylase deficiency would prevent the patient from being exposed to the potential adverse effects of long term treatment and may also avoid the necessity for infertility treatment.

GRAHAM C TOMS
JOHN P MONSON

Royal London Hospital,
London E1 1BB

- 1 Conway GS, Jacobs HS. Hirsutism. *BMJ* 1990;301:619-20. (29 September.)
- 2 New MI, White PC, Pang S, Dupont B, Speiser PW. The adrenal hyperplasias. In: Scriver CR, Beaudet AL, Sly WS, Valle D, eds. *The metabolic basis of inherited disease*. 6th ed. Vol 2. New York: McGraw-Hill, 1989:1881-917.
- 3 Speiser PW, Dupont B, Rubinstein P, Piazza A, Kastelan A, New MI. High frequency of nonclassical steroid 21-hydroxylase deficiency. *Am J Hum Genet* 1985;37:650-67.
- 4 Kohn B, Levine LS, Pollack MS, et al. Late onset steroid 21-hydroxylase deficiency: a variant of classical congenital adrenal hyperplasia. *J Clin Endocrinol Metab* 1982;55:817-27.
- 5 Kuttann F, Coullin P, Girard F, et al. Late-onset adrenal hyperplasia in hirsutism. *N Engl J Med* 1985;313:224-31.
- 6 Chrousos GP, Loriaux DL, Mann DL, Cutler GB. Late onset 21-hydroxylase deficiency mimicking idiopathic hirsutism or polycystic ovarian disease: an allelic variant of virilizing congenital adrenal hyperplasia with a milder enzymatic deficiency. *Ann Intern Med* 1982;96:143-8.
- 7 Lobo RA, Goebelsman U. Adult manifestation of congenital adrenal hyperplasia due to incomplete 21-hydroxylase deficiency mimicking polycystic ovarian disease. *Am J Obstet Gynecol* 1980;138:720-6.
- 8 Conway GS, Honour JW, Jacobs HS. Heterogeneity of the polycystic ovary syndrome: clinical, endocrine and ultrasound features in 556 patients. *Clin Endocrinol* 1989;30:459-70.
- 9 Eldar-Geva T, Hurwitz A, Vecsei P, Palti Z, Milwidsky A, Rosler A. Secondary biosynthetic defects in women with late-onset congenital adrenal hyperplasia. *N Engl J Med* 1990;323:855-63.
- 10 New MI, Lorenzen F, Lerner AJ, et al. Genotyping steroid 21-hydroxylase deficiency: hormonal reference data. *J Clin Endocrinol Metab* 1983;57:320-6.
- 11 Azziz R, Zacur HA. 21-Hydroxylase deficiency in female hyperandrogenism: screening and diagnosis. *J Clin Endocrinol Metab* 1989;69:577-84.

SIR,—Dr Gerard S Conway and Professor Howard S Jacobs fail to mention the importance of the responsiveness of hair follicles to androgen stimulation.¹

Though there is evidence that most women with hirsutism have excess androgen concentrations, some women with excess androgen concentrations do not have hirsutism, and the correlation between the degree of androgen excess and the degree of hirsutism is poor.² A widely accepted explanation for these findings is appreciable interindividual variation in response of hair follicles to circulating androgens. One possible mechanism is an increased activity of 5 α -reductase in the skin of women with hirsutism. This enzyme converts testosterone to the more biologically active dihydrotestosterone and its activity has been shown to be increased in genital skin of women with hirsutism when compared with that of controls.³

It is important to appreciate the role of hair follicle responsiveness to circulating androgens in hirsutism. Antiandrogen treatments directed more specifically at the skin may provide the best future approach to treating hirsutism.

R A SHEEHAN-DARE

Leeds General Infirmary, Leeds LS1 3EX

- 1 Conway GS, Jacobs HS. Hirsutism. *BMJ* 1990;301:619-20. (29 September.)
- 2 Reingold SB, Rosenfield RL. The relationship of mild hirsutism or acne to androgens. *Arch Dermatol* 1987;123:209-12.
- 3 Serafini P, Lobo RA. Increased 5 α -reductase activity in idiopathic hirsutism. *Fertil Steril* 1985;43:74-8.

Catheterisation

SIR,—The results of Mr R Carter and colleagues¹ have some interesting corollaries with our studies on nurses and catheterisation.² In our survey of

294 patients with catheters in five district general hospitals only 1% of catheterisations in women performed on the ward were done by a doctor, and, indeed, 52% of these procedures had been proposed by nursing staff.² Female length catheters were used for only five of the 165 women. The choice of catheter gauge size was generally appropriate, sizes above 16 Charrière being used only in patients who had had a prostatectomy, but 72% of catheters used had a balloon size of 30 ml, usually only necessary after urological surgery.³ A simple audit of equipment can rapidly pinpoint deficiencies in both the stocks of catheter available and their storage conditions.⁴

The confusing array of catheters available requires that both the selection of stock held on a ward and the choice of catheter for each patient be undertaken by expert staff—for example, the urologist or nurse continence adviser.

Finally, nurses providing equipment for catheterisations will probably have received formal instruction in this aspect of patient care.

ANNE MULHALL

University of Surrey,
Guildford GU2 5XH

- 1 Carter R, Aitchison M, Mufti GR, Scott R. Catheterisation: your urethra in their hands. *BMJ* 1990;307:905. (20 October.)
- 2 Crow RA, Chapman RG, Mulhall AB. Indwelling catheterisation and related nursing practice. *Journal of Advanced Nursing* 1988;13:489-95.
- 3 Slade N, Gillespie WA. *The urinary tract and the catheter. Infection and other problems*. Chichester: John Wiley, 1985.
- 4 Mulhall AB, Lee K. The provision of urethral catheter. An equipment audit. *Quality Assurance in Health Care* (in press).

Influenza vaccination and the elderly

SIR,—Dr K G Nicholson raises interesting points about immunisation against influenza.¹ I wish to report the results of an attempt to improve immunisation uptake.

The study was based in a West Lothian health centre over the winters of 1986-7 and 1987-8. We invited all registered patients who were thought to be appropriate for immunisation as determined from computerised medical records by disease categories and age over 65 years. The non-responders were sent another letter to explore the reasons for non-attendance. A more comprehensive letter (explaining reasons for immunisation and possible side effects) was sent in the next autumn and the same follow up letter was sent to non-responders. In both these programmes the patients were sent a letter with an appointment time for immunisation and any immobile or housebound patient was offered a house visit. We thus tried to ensure that patients' refusal to be immunised was not due to us not putting over the message properly.

The table shows the results. Overall, the uptake was 70.4% (629/894) in 1986-7 and 60.6% (419/691) in 1987-8 (only those patients registered with the practice for both winters were included in the second year). Very few patients attended the health centre because of our follow up letter about non-attendance.

Reasons for non-uptake of influenza vaccination in patients considered appropriate for immunisation over winters of 1986-7 and 1987-8. Values are numbers of patients

	1987-8							
	Not offered, died, or moved away	No chest problems	Fear of needles	Protection not important	Previous bad reaction	No reply	Accepted	Administrative reasons
1986-7								
No chest problems	6	5	1			6	8	1
Fear of needles	2		2				2	
Protection not important	6	1		1	1	13	3	
Previous bad reaction	5			1	17	9	9	
No reply						51	14	
Accepted	157	1		1	10	86	352	6
Administrative reasons	13	1			1	8	12	1
Minor reasons	9					13	19	1
1987-8								
No chest problems								
Fear of needles								
Protection not important								
Previous bad reaction								
No reply								
Accepted								
Administrative reasons								
Minor reasons								

The results are surprising. There were cohorts of patients who thought that it was unimportant to be protected against influenza or who had had adverse reactions, and these patients did not alter their response to invitation to immunisation despite a signed explanatory request from their doctor. Nevertheless, an increase in uptake was seen in those patients who had previously regarded themselves as not having a chest problem. This phenomenon has been noted in part before.² We were disappointed that the level of uptake dropped appreciably despite a comprehensive explanatory letter, and we also note a lack of consistency in reasons for refusal for immunisation.

Nicholson *et al* have suggested that providing educational material about the risks and benefits of influenza vaccine and reappraising practice strategies may increase the number of elderly people who are immunised.¹ Unfortunately, our experience in patients of all ages does not confirm this.

I R WAKEFIELD

Edinburgh EH4 1PU

- 1 Nicholson KG. Influenza vaccination and the elderly. *BMJ* 1990;301:617-8. (29 September.)
- 2 Frank JW, Henderson M, McMurray L. Influenza vaccination in the elderly. 1. Determinants of acceptance. *Can Med Assoc J* 1985;132:371-5.
- 3 Nicholson KG, Wiselka MJ, May A. Influenza vaccination of the elderly: perceptions and policies of general practitioners and outcome of the 1985-86 immunization programme in Trent, UK. *Vaccine* 1985;5:302-6.

HIV and surgeons

SIR,—Dr B G Gazzard and Professor C Wastell wrote about the risk of HIV infection among surgeons.¹ Orthopaedic surgeons believe that they are at greater risk during operations than most other surgeons because of their extensive concern in trauma, the sharpness of skeletal fragments, and the nature of orthopaedic instrumentation. The British Orthopaedic Association over the past year has been formulating a policy to help minimise the risk of transmission of HIV and other bloodborne viruses during orthopaedic operations and procedures.

Though we accept that the risk at present in the United Kingdom is small, we do not accept that it can be ignored. The authors comment that nurses seem to be most at risk but fail to recognise that, in the United Kingdom at least, there are more than 10 nurses for each doctor in the health service, and therefore, statistically, more cases would be expected among nurses than among doctors (statement by the chief medical officer to the council of the Royal College of Surgeons, October 1990).

The rather dismissive attitude towards universal precautions is alarming. Though there is considerable disagreement over the need for routine testing for HIV, there does seem to be complete agreement on using universal precautions.^{2,3} This policy is also the mainstay of the United Kingdom health departments' *Guidance for Clinical Health Care Workers*.⁴

As AIDS is spreading these universal precautions should become normal practice throughout