SIR,—Dr Stephen G Spiro is being slightly pessimistic about the overall survival rate after resection of non-small cell lung cancer.¹ He quotes a figure of “around one quarter after five years” and this certainly was the figure 25 years ago, but with improved screening techniques it has been possible to be more selective in choosing patients for resection, with consequent better results.

In 1976 Stanford *et al* reported an overall five year survival rate of 48.8% in patients having definitive surgery.² This was a series of over 1000 patients in the American armed forces. More recently Kadri and I reported a British series with an overall five year survival rate of 38.6%.³ These results are not confined to the young either, as Ishida *et al* recently reported a five year survival of 48% in patients over 70 and 41% in those under 70.⁴

The tragedy is that with improved screening techniques the population of patients suitable for operation has diminished even more. In the south east region nearly 3000 patients die each year of lung cancer, yet the number of patients suitable for surgery is in the region of 200 a year.

J E DUSSEK

Cardiothoracic Unit,
Guy's Hospital,
London SE1

- 1 Spiro SG. Management of lung cancer. *BMJ* 1990;301:1287-8. (8 December.)
- 2 Stanford W, Spirey C, Larsen G, *et al*. Results of treatment of primary carcinoma of the lung. *J Thorac Cardiovasc Surg* 1976;72:441-9.
- 3 Kadri MA, Dussek JE. Survival and prognosis following resection for bronchogenic carcinoma. *Thorax* 1990;45:306-7P.
- 4 Ishida T, Yokoyama H, Kaneko S, *et al*. Long term results of operation for non small cell lung cancer in the elderly. *Ann Thorac Surg* 1990;50:919-22.

Milk for babies and children

SIR,—Follow up milks were created by the baby food industry deliberately to avoid the advertising restrictions of the World Health Organisation's international code of marketing of breast milk substitutes (1981), and since then companies have fought hard to establish credibility for them. Professor B Wharton defends these milks yet provides no adequate rationale for their use or explanation why mothers might “wish to move on from an infant formula.”^{1,2} Perhaps an answer can be found in British parents' magazines, in which follow up milks are promoted in competitions with prizes of bedroom suites and weekend breaks.

As Dr Talukder pointed out in his letter,³ British publications are regularly exported to developing countries and have a profound effect on feeding practices. In many countries these expensive milks are also aggressively promoted on television and on billboards and are confused with infant formulas by illiterate parents. As a consequence, despite the fact that the World Health Assembly of 1986 described follow up milks as unnecessary, their sales are rising steadily. In Niger, for example, 88 000 tins of Nestlé infant formula and 64 000 tins of Nestlé follow up formula are now used every year.

Professor Wharton regards his inaccuracies over the extra cost of powdered milks in the United Kingdom as a mere detail and assumes that parents would use the extra £1.88 per week (£4.55 if the mother continues to breast feed) to buy beer, cigarettes, or disposable nappies rather than the cheap, wholesome foods that a baby might need. His suggestion that public policies should be changed to raise the income of poor families is laudable, but until such time should we not investigate thoroughly whether the additional expense of proprietary milks carries any significant benefit? Professor Wharton is worried about promoting “substandard feeding customs” for the sake of cost but does not consider the damaging effect that promotion of follow up milks has had on infant feeding standards world wide.

Following widespread concern about the confusion and misuse of these milks when they were first introduced to the United Kingdom in 1985 (shown by the volume of letters to the editor published in *Health Visitor* from March to September) the Committee on the Medical Aspects of Food Policy, of which Professor Wharton was a member, rightly recommended that these milks should not be marketed for babies under 6 months. On the advice of the Scientific Committee for Food in 1988 the European Commission is now proposing a directive allowing unlimited promotion of follow up milks for babies of up to only 4 months—not even meeting Codex standards.

This may seem a small matter to paediatricians, but it is vital to the baby food industry. When the 1985 decision was taken health professionals expressed relief, but sales of Wyeth's follow up milk fell noticeably. Since 1985, however, promotion of follow up milks in the United Kingdom has continued to confuse parents and health workers by overemphasising their supposed advantages over “doorstep milk” and neglecting to mention the government's main message, that breast feeding can be continued with advantage throughout the first year. We now have reports of chemists recommending follow up milks for babies who cannot sleep at night or who are allergic to cows' milk and of breast feeding mothers who have changed to follow up milks because they are worried about their babies' diet. If a European Commission directive is adopted with the 4 month age limit the situation will be made much worse and more babies will be put at risk.

Unlike the Committee on Safety of Medicines, neither the Committee on the Medical Aspects of Food Policy nor the Scientific Committee on Food makes public the links that members may have with the food industry, and committee members declare their interests only to fellow members. Inquirers are asked to accept on trust that committee members are impartial. What is there to be gained from such secrecy? Before we allow companies the right to advertise these products as they wish should we not examine independent evidence to see if these milks can have a significant beneficial effect on children? The study by Fomon *et al*, for example, concludes that there is no scientific reason for changing from a normal infant formula to a follow up milk.⁴ If their only known benefit is to company profits isn't it time that doctors addressed the wider implications of their promotion?

PATTI RUNDALL

Baby Milk Action,
Cambridge CB2 1AA

- 1 Wharton B. Milk for babies and children. *BMJ* 1990;301:774-5. (6 October.)
- 2 Wharton B. Milk for babies and children. *BMJ* 1990;301:1392-3. (15 December.)
- 3 Talukder MQ-K. Milk for babies and children. *BMJ* 1990;301:1392. (15 December.)
- 4 Fomon S, Saunders KD, Zeigler EE. Formulas for older infants. *J Pediatr* 1990;118:690-8.

Unexplained haematuria

SIR,—We have done two studies in this department that may help to resolve some of the questions raised by Drs Mark P de Caestecker and Francis W Ballardie.¹ We screened the staff of the Royal Melbourne Hospital and found that some 6% had evidence of haematuria, this being almost invariably glomerular in type. They were referred to their local doctors, and, interestingly, very few were subsequently referred to nephrologists for further evaluation of the documented microscopic haematuria.

We have over a period of 20 years had a policy of performing biopsies of all patients referred with documented glomerular microscopic haematuria. Of 70 patients with isolated glomerular haematuria

seen in the past 12 months, most had either mesangial IgA glomerulonephritis (21) or thin basement membrane disease (37). Drs de Caestecker and Ballardie seem more reluctant to carry out biopsies on the basis of isolated microscopic haematuria, but we have shown that a continuing high urinary erythrocyte count is the best guide to activity and progression in mesangial IgA glomerulonephritis,^{2,3} and it is important to distinguish this condition—which is progressive in 20% of cases—from thin basement disease, which is usually benign. There are probably non-invasive methods of making the distinction between these two common lesions. We presented data at the 1990 American Society of Nephrology meeting to show that concentrations of platelet factor 4 are raised in mesangial IgA glomerulonephritis but not in thin basement membrane disease, and this may provide a reliable non-invasive method of distinguishing these two conditions.

PRISCILLA KINCAID-SMITH
JULIE OWEN
TIM HEWITSON

Department of Nephrology,
Royal Melbourne Hospital,
Melbourne,
Victoria 3050,
Australia

- 1 de Caestecker MP, Ballardie FW. Unexplained haematuria. *BMJ* 1990;301:1171-2. (24 November.)
- 2 Kincaid-Smith P, Nicholls K. Mesangial IgA nephropathy. *Am J Kidney Dis* 1983;3:90-4.
- 3 Bennett WM, Kincaid-Smith P. Macroscopic hematuria in mesangial IgA nephropathy: clinical pathologic correlations. *Kidney Int* 1983;23:393-400.
- 4 Nicholls KM, Fairley KF, Dowling JP, Kincaid-Smith P. The clinical course of mesangial IgA nephropathy. *Q J Med* 1984;53:227-50.
- 5 Kincaid-Smith P, Nicholls K, Birchall I. Polymorphs participate in glomerular injury in mesangial IgA glomerulonephritis. *Kidney Int* 1989;36:1108-11.

SIR,—The editorial by Drs Mark P de Caestecker and Francis W Ballardie¹ was marred by two peculiarities: the authors made no attempt to explain how or why red cells appear in the urine either in good health or during the course of renal disease, and there was no comment on the reported occurrence of red cells in the urine of healthy adults² and healthy children.³ As those authors proposed “normal” limits for urinary erythrocytes it seems clear that renal pathology is not a necessary prerequisite for the passage of red cells through vessel walls. Gibbs and Lynn reported an increased number of urinary blood cells after strenuous exercise⁴ so it seems possible that the mechanism of haematuria in healthy subjects may be directly comparable to that of exercise proteinuria.

Ohsaka *et al* showed by electron microscopy how red cells were extruded through capillary walls at interendothelial cell junctions when intracapillary pressure was raised by massive vasoconstriction caused by habu snake venom.⁵ Their observations, which could provide a basis for understanding the mechanism of haematuria, together with high molecular weight proteinuria, are explicable in terms of basement membrane thixotropy.^{6,7} The pressure dependent permeability of basement membranes was used by Warfel and Hull to explain how white cells migrated through the cutaneous basement membrane.⁸

Although Drs de Caestecker and Ballardie seem to accept the proposal that thin membrane nephropathy might provide a basis for understanding haematuria, this view was not supported by the writer of a recent editorial in the *Lancet*.⁹ That article, titled “Thin-membrane nephropathy—how thin is thin?” noted that it was “intuitively appealing” to imagine that red cells could leak through the basement membrane of the glomerular filter because of the thinness of the membrane, but concluded that “it may be unhelpful to talk with great conviction of thin-membrane nephro-