into a short report forced us to omit these details. The trials quoted by Dr Crossman were also not mentioned for this reason. His recommendation, to stop treatment only in those who had mild hypertension before treatment, is not consistent with the evidence that he invokes.

We think it probable that as a result of our large scale trial and work already published on "white coat hypertension," the accepted criteria for starting drug treatment will have to be modified. Neither the criteria of the MRC trials nor those of the British Hypertension Society, nor indeed Professor Sever's recommendations of "two or more sets of readings over a period of days or weeks," will, we believe, stand the test of time.

The practices around St Bartholomew's Hospital treat far more patients with mild hypertension than does Dr Lim's cardiology department. We do not consider his comments about his general practitioner colleagues to be constructive.

MALCOLM AYLETT
STUART KETCHIN

Glendale Surgery,
Wooler,
Northumberland,
NE71 6DN

1 Lim R. Stopping treatment in patients with hypertension. *BMJ* 1991;303:469. (24 August.)

SIR,—We were not surprised that Drs Malcolm Aylett and Stuart Ketchin were able to withdraw antihypertensive treatment successfully from a small number of highly selected patients,¹ but we wish to echo the note of caution sounded by Dr Richard Lim.² These patients were quite simply the product of an era when antihypertensive treatment was introduced too readily. Treatment was started if the diastolic blood pressure was higher than 100 mm Hg on as few as three occasions, and the Australian and Medical Research Council mild hypertension trials showed clearly that many subjects satisfying this criterion eventually become normotensive without treatment.

The main message from Drs Aylett and Ketchin's study is not that patients starting treatment now should be advised that it need not be lifelong—it is that treatment should not be started until it is certain that it is necessary. The accompanying editorial recognised this³ and cited the British Hypertension Society recommendation that mild hypertension (initial diastolic pressure 90–109 mm Hg)⁴ should be followed for three months before deciding on the need for treatment. In our view, observation for three months is still far too short. In the Australian trial it took four months for the blood pressure to plateau at its final level in mild hypertension treated with placebo.⁵ The implication is that the treatment decision should be reached on the average blood pressure beyond four months—for example, by using measurements taken at four to six months. Observation should be extended even further when non-pharmacological measures such as weight reduction, alcohol reduction, and moderate salt restriction are being used. Prolonged observation is, however, inappropriate when end organ damage is already present, and should be curtailed if the diastolic pressure exceeds 110 mm Hg repeatedly. If antihypertensive drugs are introduced only when diastolic pressure averages 100 mm Hg or higher after a full six months of observation and after a thorough trial of non-pharmacological measures then we doubt that many subjects will subsequently become normotensive if treatment is stopped. When these criteria are followed it would be unkind and potentially dangerous to raise expectations that drug treatment is likely to be only temporary.

There are doubtless many patients now taking antihypertensive drugs who should not have started treatment by today's criteria. It is therefore good practice to attempt to "step down" treatment in

well controlled patients, as Dr Aylett and Ketchin did, with the proviso that follow up should be continued for a full year after treatment has been stopped before declaring the patient normotensive. Even then continuing annual checks of blood pressure would be advisable.

P R JACKSON
W W YEO
L E RAMSAY

Sheffield Hypertension Clinic,
University Department of Medicine and Pharmacology,
Royal Hallamshire Hospital,
Sheffield S10 2JF

- 1 Aylett M, Ketchin S. Stopping treatment in patients with hypertension. *BMJ* 1991;303:345. (10 August.)
- 2 Lim R. Stopping treatment in patients with hypertension. *BMJ* 1991;303:469. (24 August.)
- 3 Burton R. Withdrawing antihypertensive treatment. *BMJ* 1991;303:324–5. (10 August.)
- 4 Swales JD, Ramsay LE, Coope JR. Treating mild hypertension. *BMJ* 1989;298:694–8.
- 5 Ramsay LE. Mild hypertension: treat patients, not populations. *J Hypertens* 1985;3:449–55.

SIR,—British general practice is the best primary care service in the world¹ and does not merit the criticism made by Dr Richard Lim in his letter.² He implies that some general practitioners unjustifiably discontinue treatment as they do not appreciate "that blood pressure may have been normal because of the treatment." He also alleges that general practitioners are irresponsible in not adequately monitoring such patients after stopping their treatment. It is a sad reflection of the standard of care provided by general practitioners in his locality. I can assure Dr Lim that such irresponsible and irrational practice is not a typical representation of care provided in general practice in Britain.

It is important for Dr Lim to appreciate that there are over seven million patients in the British Isles who suffer from hypertension³ and that over 95% of them are managed by their general practitioners. General practitioners have done much to promote understanding of the disease and research. The value of community screening was highlighted by a general practitioner.⁴ British general practitioners worked with the Medical Research Council to dispel the misconception that treatment of mild to moderate hypertension prevents myocardial infarction in all patients.⁵ The need to incorporate patients' perception of treatment and the effect of treatment on their quality of life was suggested by a general practitioner.⁶

It has also been shown that blood pressure can be adequately controlled without therapeutic intervention. I commend Drs Malcolm Aylett and Stuart Ketchin, who are innovative enough to point out that it is necessary to review our practice of controlling blood pressure and even to consider stopping treatment in some of the patients who do not warrant it.⁷ This is to preserve their quality of life, prevent adverse reactions to the drugs, and economise on use of resources. It is regrettable that Dr Lim did not appreciate the main theme of the article. He is justified, however, in sounding such caution if he is not supported by effective and efficient primary care.

S J JACHUCK

Medical Research Centre,
Newcastle upon Tyne NE5 2LH

- 1 Bosanquet N. Family doctors and payment systems: the local options. *BMJ* 1991;303:233–4. (27 July.)
- 2 Lim R. Stopping treatment in patients with hypertension. *BMJ* 1991;303:469. (24 August.)
- 3 Beevers DG, Messerli FH. Hypertension—moving towards the year 2000. *Human Hypertension* 1987;1:1–2.
- 4 Hart JT. *Hypertension: community control of high blood pressure*. 2nd ed. London: Churchill Livingstone, 1987.
- 5 Medical Research Council Working Party. MRC trial of mild hypertension: principal results. *BMJ* 1985;291:97–104.
- 6 Jachuck SJ, Brierly H, Jachuck S, Willcox PM. The effect of hypotensive drugs on the quality of life. *J R Coll Gen Pract* 1982;32:103–5.
- 7 Aylett M, Ketchin S. Stopping treatment in patients with hypertension. *BMJ* 1991;303:345. (10 August.)

SIR,—Drs Malcolm Aylett and Stuart Ketchin are optimistic that many patients currently receiving drug treatment for hypertension could have their treatment stopped, which would improve the quality of their lives and save resources for the NHS. Dr Ralph Burton is more cautious, advocating prolonged observation before treatment and advising new patients with hypertension that their treatment may not be lifelong.¹ He does not discuss non-drug treatment of hypertension.

Evidence suggests that regular physical activity can be as effective as drug treatment,² and reducing weight,³ restricting salt intake,⁴ a vegetarian diet,⁵ relaxation and stress management,⁶ and reducing alcohol intake cause worthwhile reductions in blood pressure. The advantage of non-drug treatments is that many have other healthy "side effects" and that patients are motivated to change their lifestyle. The disadvantage of taking drugs daily with its risk of side effects and cost may be avoided, and patients feel that they can affect their wellbeing and are not victims of disease.

ALISON J WATSON

Department of Public Health Medicine,
Sunderland District General Hospital,
Sunderland SR4 7TP

- 1 Aylett M, Ketchin S. Stopping treatment in patients with hypertension. *BMJ* 1991;303:345. (10 August.)
- 2 Burton R. Withdrawing antihypertensive treatment. *BMJ* 1991;303:324–5. (10 August.)
- 3 Royal College of Physicians of London. *Medical aspects of exercise: benefits and risks*. London: RCP, 1991.
- 4 Croft PR, Brigg D, Smith S, Harrison CB, Branthwaite A, Collins MF. How useful is weight reduction in the management of hypertension? *J R Coll Gen Pract* 1986;36:445–8.
- 5 Australian National Health and Medical Research Council Dietary Salt Study Management Committee. Fall in blood pressure with modest reduction in dietary salt intake in mild hypertension. *Lancet* 1989;i:399–402.
- 6 Law MR, Frost CD, Wald NJ. By how much does dietary salt reduction lower blood pressure? III. Analysis of data from trials of salt reduction. *BMJ* 1991;302:819–24.
- 7 Margetts B, Beilin LJ, Vandongen R, Armstrong BK. Vegetarian diet in mild hypertension: a randomised controlled trial. *BMJ* 1986;293:1468–71.
- 8 Patel C, Marmott M. Can general practitioners use training in relaxation and management of stress to reduce mild hypertension? *BMJ* 1988;296:21–4.

SIR,—The suggestion by Drs Malcolm Aylett and Stuart Ketchin that treatment for hypertension can be stopped in some patients is feasible,¹ and we accept that the savings in drug bills and benefits for patients no longer needing drug treatment are clearly to be welcomed. We are, however, concerned that the principle could be embraced inappropriately.

From our work in cardiology and hypertension clinics it sometimes seems that not all doctors are yet convinced of the value of treating hypertension at all. It is lamentable how often we see patients with end stage hypertensive heart disease, with electrocardiographic changes and radiographic evidence of cardiomegaly, who were first told that their blood pressure was "a bit up" 10 or even 20 years ago and yet have either never received treatment or, worse, have had their treatment stopped by somebody who has never followed that decision up. Such patients are at high risk of sudden cardiac death,² and yet modern treatments for hypertension are quite able to prevent the development of left ventricular damage.³

It is true that there is still much that we do not know about hypertension. New techniques such as ambulatory blood pressure monitoring are teaching us more. It is clear that some people with apparently high pressures are merely "labile" or have "white coat hypertension" whereas others with apparently modest hypertension have a more consistently raised ventricular load. Though the brief study of Drs Aylett and Ketchin and the larger Australian and Medical Research Council trials⁴ have shown us that patients diagnosed as having hypertension at a particular time may not remain hypertensive later, neither of these trials explains the reason for