CORRESPONDENCE

Drugs for Rare Diseases J. M. Walshe, F.R.C.P. 701	Cerebral Oedema in Diabetic Ketoacidosis B. M. Frier, M.R.C.P. 704	¹²⁵I-fibrinogen Standardization R. J. Hawker, M.Sc., and others. 706
Identification of High-risk Labours J. W. W. Studd, M.R.C.O.G. 702	Certification of Death A. R. Griew, M.B. 704	Private Practice in the N.H.S. M. Wallace, F.R.C.S.; M. J. Illingworth, M.B. 706
G.M.C. and Overseas Doctors Lieutenant-Colonel R. J. Henderson, M.D. 702	Services Available for Total Joint Replacement R. T. Austin, F.R.C.S.ED. 704	Junior Hospital Staff Contract M. S. Amos, M.D., and others; J. L. Mitchell, M.R.C.P., and others; M. R. Lowe, M.B. 707
Carrier Solutions for Low-level Intravenous Insulin M. Mc.B. Page, M.R.C.P., and others. 702	Estimation of Glomerular Filtration Rate W. D. Morgan, M.B., and others. 704	Consultants' Work-to-contract J. W. Dickson, F.R.C.S. 707
Are Our Barbiturates Really Necessary? G. D. Pirrie, M.B. 702	Adjustment to Low-dose Heparin to Prevent Thrombosis J. R. O'Brien, F.R.C.PATH. 705	Revenue Allocation in the Reorganized Health Service P. J. N. Howorth, M.D.; H. M. Buckland, M.B. 707
Prescribing Habits G. R. Fowler, B.PHARM. 703	Urinary NAG Activity in Renal Disease P. R. Beck, M.B. 705	Mortgage Problems C. Nemeth, M.R.C.S. 708
Drugs for Addicts A. H. de C. Freed, M.R.C.PSYCH. 703	Myocardial Scanning E. J. Wakley, M.R.C.P., and others. 705	Points from Letters Side Effects of Amitriptyline (M. A. M. Khan); Chloroquine Prophylaxis in East Africa (A. R. Shaw); Colour Disturbance as a Symptom (G. R. Davies); Post-herpetic Pruritus (K. Liddell); Hang-glider Injuries (D. L. Hadley) 708
Aetiology of Optic Neuritis F. C. Rose, F.R.C.P.; A. L. Crombie, F.R.C.S.ED., and others. 703	Malaria in Scotland Neena Stewart, M.B., D.T.M. & H. 705	

Correspondents are urged to write briefly so that readers may be offered as wide a selection of letters as possible. So many are being received that the omission of some is inevitable. Letters should be signed personally by all their authors.

Drugs for Rare Diseases

SIR,—It is almost exactly 20 years since I first started to investigate the possibility of using penicillamine in the management of Wilson's disease.¹ I do not wish to launch into a detailed account of the various and not inconsiderable obstacles which had to be overcome before this new drug was accepted into the pharmacopoeia. Suffice it to say that it underwent a minimum of toxicity testing. The fact that penicillamine had been detected in the urine of patients receiving parenteral penicillin² suggested that toxicity would be no great problem, at least as far as the D-isomer was concerned.³ I took a single dose and, at my request, Merck, Sharpe, and Dohme conducted some acute toxicity tests on mice. Thereafter the compound was brought into clinical use. That it has proved a major breakthrough in the management of Wilson's disease can hardly be denied⁴ and it is of interest that this drug has been found to be of value in other heavy-metal intoxications and in a number of other, unrelated diseases—namely, cystinuria, rheumatoid arthritis, certain diseases of collagen, and perhaps the macroglobulinaemias. It would seem highly improbable that, under present-day conditions, such a puny infant would have blossomed into such a robust member of the therapeutic community. The regulations for the testing and introduction of new drugs would surely have stifled it at birth. Indeed I have evidence that such may be the case.

Penicillamine, like all other active drugs, has certain undesirable side reactions. To some extent it has proved to be a veritable Pandora's box and in a recent leading article (19 July, p. 120) you listed the widespread

and varied toxic reactions which must be set against penicillamine. I will not detail them again here except to note that three patients with Wilson's disease taking this drug have died of Goodpasture's syndrome. Fortunately most of these complications are uncommon in patients with Wilson's disease and all will remit if treatment is stopped in time. For those patients whose disease can be managed by alternative methods this is no great disaster, but for patients with Wilson's disease withdrawal of treatment is tantamount to a death sentence.

In 1966 I was confronted with this very problem when a youth of 15 years, who had been taking penicillamine for Wilson's disease for six years, developed the nephrotic syndrome. Two attempts at desensitization proved unsuccessful, but eventually the situation was saved by the introduction of a new orally active chelating agent, triethylene tetramine dihydrochloride (trien 2HC1).⁵ Now, six years later, he is maintained in good health on this drug. Subsequently it has been necessary to transfer three more patients to this form of treatment, one because of severe thrombocytopenia and one because of the development of a rheumatoid-like arthropathy, while the third developed severe buccal ulceration. Treatment of these patients over the years has been continued simply because the purified trien 2HC1 has been prepared in my laboratory⁶ and encapsulated in the Addenbrooke's Hospital pharmacy. The compound has never been subjected to formal toxicity testing nor has any pharmaceutical company seen fit to undertake its production. This is hardly surprising in view of the thalidomide disaster

and its unfortunate repercussions and because of the enormous expense of testing and clearing a new drug through the Medicines Commission. Such considerations will probably apply to any new drug designed for the management of a rare disease; the exercise will simply not be commercially viable and, whether we like to admit it or not, drug firms, like the rest of us, must make money to remain in business. For a private individual to prepare and supply a drug for clinical use must be legally hazardous whatever the indications may be. Fortunately the situation has not yet been tested in the courts, but to quote a statement I have made elsewhere, "The mind boggles at the thought of what a learned judge might say should such an experiment prove a failure."⁷

Recently the hospital pharmacy agreed to take over the purification of trien 2HC1 as well as the encapsulation, but the situation remains unclear as they will not be able to do so without the issue of a product licence. In my view this do-it-yourself exercise has continued quite long enough and should be placed on a sound commercial basis. Trien should be available on the market, to any one who needs it. Its production should not depend upon the drive of a single individual and cease with his retirement or demise; the drug was designed for and has been proved by clinical use.⁷ It might be argued that it is the duty of the Minister in charge of the Health Service, as laid down by Parliament, that such treatment, if it does not put an unreasonable load on the country's resources, be made available to those patients requiring it.

Meantime the question arises as to what will happen to these patients should I retire from the scene or should a product licence not be issued. Are they to be allowed to die of a readily treatable disease because no one is prepared to supply, or worse still is per-

mitted to produce, the necessary medication? Such a refusal would be little better than an act of legislative homicide. I shall appreciate enlightenment in this dilemma.—I am, etc.,

J. M. WALSH

University Department of Medicine,
Addenbrooke's Hospital,
Cambridge

- 1 Walshe, J. M., *Lancet*, 1956, 1, 25.
- 2 Walshe, J. M., *Quarterly Journal of Medicine*, 1953, 22, 483.
- 3 Walshe, J. M., *American Journal of Medicine*, 1956, 21, 487.
- 4 Walshe, J. M., in *Treatment of Inherited Metabolic Disease*, ed. D. N. Raine. Lancaster, M.T.P., 1975.
- 5 Walshe, J. M., *Lancet*, 1969, 2, 1401.
- 6 Dixon, H. B. F., Gibbs, K., and Walshe, J. M., *Lancet*, 1972, 1, 853.
- 7 Walshe, J. M., *Quarterly Journal of Medicine*, 1973, 42, 441.

Identification of High-risk Labours

SIR,—In commenting on our paper on this subject (7 June, p. 545) Mr. D. D. Matthews (19 July, p. 157) correctly draws attention to the special dangers of delayed multigravid labour and Professor M. K. O'Driscoll (p. 157) points out that the multigravid labours not stimulated according to the protocol appear to have had a slightly longer first stage than a similarly selected group of dysfunctional primigravid labours. Though such difference is not statistically significant, the point is taken that these multigravid labours are not shorter. It was stated that this group were either unstimulated or received oxytocins more than four hours past the nomogram. The long multigravid labours were the direct result of unwillingness to use oxytocin in these cases.

Primigravid labour is, in general, longer than multigravid labour, but after exclusion of complicating or iatrogenic factors the length of the first stage from a given admission cervical dilatation is identical for both groups.¹ Abnormal multigravid labour occurs less frequently, but as it is more treacherous the need for careful selection of patients for stimulation based upon projected normal labour patterns was stressed, with only 13% of multigravid labours requiring stimulation compared with 36% of primigravidae.

The data presented showed a longer first stage, higher instrumentation rate, and lower Apgar scores for babies born to those patients in the unstimulated or belatedly stimulated group, and I express surprise that these differences "do not appear to be great enough to substantiate the claim," particularly as this group had a greater mean admission cervical dilatation and were mistakenly adjudged to be progressing well enough not to require stimulation.

Professor O'Driscoll does uncover an omission by his call for elaboration of the zero perinatal loss of 741 spontaneous labours in that this continuing study of the dynamics of labour excluded fetal abnormalities incompatible with life, low-birth-weight babies (<2.5 kg), substantial antepartum haemorrhages, cord accidents, breech deliveries, and multiple births.¹ The overall caesarean section rate of 1.4% compares with a 2.3% caesarean section rate in a similar group of spontaneous labours of a large statisticographical labour study yet to be published. The different cervimetric progress of unstimulated spontaneous labours and those augmented clearly shows that dysfunctional labour may be detected by the assessment of cervical dilatation on admission in labour and by another

Progress of cervical dilatation in spontaneous labour without epidural anaesthesia. (M=multigravidae; P=primigravidae.)

vaginal examination after four hours of contractions (see figure). I do not believe that this can be regarded as a retrospective diagnosis of labour.

No doubt different criteria for the commencement of labour will remain, but it is my belief that the conclusion that (with the exception of an intrapartum haemorrhage) normal babies of mature birth weight should not die during spontaneous labour if delay is recognized and corrected, if labour is monitored, and if difficult instrumental procedures are prohibited is one which is not discrepant with the view from the National Maternity Hospital, Dublin.

The labour stencil identifies the high-risk labours which may require intensive care in hospital² or referral to a specialized unit from general practitioner supervision, ensuring that acceleration of labour may be introduced at the correct time in patients with delay in progress.—I am, etc.,

JOHN STUDD

King's College Hospital,
London S.E.5

- 1 Duignam, N. M., Studd, J. W. W., and Hughes, A. O., *British Journal of Obstetrics and Gynaecology*, 82, 593.
- 2 Studd, J. W. W., *British Medical Journal*, 1973, 4, 451.

G.M.C. and Overseas Doctors

SIR,—Dr. N. Ahmed (6 September, p. 595) raises so many points that a very long letter would be required to reply to them. I will take up one only, his proposal that better facilities should be provided in Britain for training overseas doctors. I can assure Dr. Ahmed that many of us who know India and Pakistan well, have served long years in these countries, and have affection for their inhabitants, would be only too glad to extend a welcome to their doctors and help train them: if we could be sure that when their postgraduate training was over they would return and use their acquired knowledge and skill in their own lands instead of slipping off into general practice in the U.K. or some equally lucrative work elsewhere in the west—for lucrative either of these would be compared with what they would earn at home. As a former officer in the Indian Medical Service and a recent worker for the World Health Organization I have been dismayed by the example of doctors from the Indian subcontinent or Ceylon taking up permanent posts in Britain when their postgraduate work is over, and indeed breaking the bond given to return, instead of going

back and working in their own sadly under-doctored lands.

Much outcry is being made by Indian and Pakistani doctors who fail the General Medical Council examination that they have been failed on account of their poor knowledge of English. I have even heard it said that London language colleges are preparing crash courses to enable them to pass. I do not think this is at all necessary. As a rule Indian and Pakistani doctors speak fairly good English and it does not take them long to become familiar with the way the average British patient speaks. Some, it is true, are weak in English, but this is only a minor factor; the true weakness of many Indian and Pakistani doctors is that their basic medical knowledge and technical ability are poor. While an immigrant from the subcontinent who passes the G.M.C. examination could be entrusted straight away with a house officer post, he who fails would in my view require further training, under supervision, for at least two years—that is, in four house posts of six months each, but supervised, assisted, and trained all the time. The G.M.C. could surmount the difficulty of the 60% or so who fail the examination by employing them in a new grade of trainee house officer at a salary lower than the standard house officer rate until the examiners were satisfied.

But I would like to see a rule introduced by the British Government that on completion of postgraduate training every Indian and Pakistani doctor must return to his own country, because however much the British National Health Service may be tottering, however much it may require to be propped up by imported labour, its need for medical staff is minimal compared with that of India and Pakistan.—I am, etc.,

R. J. HENDERSON

Public Health Laboratory,
Royal Infirmary,
Worcester

Carrier Solutions for Low-level Intravenous Insulin

SIR,—Dr. E. W. Kraegen and his colleagues (23 August, p. 464), advocate the use of polygeline solution to prevent adsorption of insulin to plastic or glass. This is clearly an important consideration when high dilutions of insulin are used, as in their technique when 20 units are diluted in 500 ml in the infusion container. However, if insulin is diluted in a syringe to 1-4 U/ml and given at a rate of 4-10 U/h, as is the case in the technique which we and our colleagues have described (29 June 1974, p. 687) and found very satisfactory in a series of over 60 cases, there is no clinically significant loss of insulin and complexity and expense are avoided.—We are, etc.,

M.M.C.B. PAGE
P. J. WATKINS
D. A. PYKE

King's College Hospital,
London S.E.5

Are Our Barbiturates Really Necessary?

SIR,—In the 1930s the Plaza-Torian leaders of our profession were successful in stopping a move to take heroin out of the pharmacopoeia, so that it was only after three decades culminating in the heroin "epidemic" with its crime, misery, and deaths, that the