Diagnosis using search engines

Probably heralds a much more sophisticated web resource

Research p 1143

Thousands of computer systems targeted at medical diagnosis (often described as expert systems) have been developed during the past 50 years. Most have had relatively little impact on day to day clinical practice; for example, because they are not easily accessible at the point of care; have a complex interface; can deal with only a narrow focus (one symptom or clinical problem); are not integrated with clinical information systems; depend on particular software or hardware platforms; or are expensive to maintain and extend.

In this week's *BMJ* a study by Tang and Ng assesses the effectiveness of a web search engine (Google) as a diagnostic aid.¹ Using general purpose web search engines as a diagnostic aid is new, although using computers to aid diagnosis is not. The article, published online on 10 November 2006, has already provoked a widespread response, with one daily newspaper concluding that "Researchers have found that a simple Google search can solve diagnostic problems which mystify even the best general practitioners."²

This conclusion is misleading in at least three respects. Firstly, the Google searches reported in the study were not simple. Although the search required typing only a few words and clicking a button, doctors used extensive knowledge and experience to choose the search terms effectively. Secondly, Google did not solve the diagnostic problems. It returned a set of documents ranked according to their relation to the search terms, using metrics essentially based on counting the occurrence of word stems. Inferring a diagnosis from the returned documents depended on the skill of the doctors. Thirdly, the implication that diagnostic problems can be ordered on a linear scale of difficulty, against which human and machine performance can be compared, is unwarranted. The study shows that in a certain class of diagnostic challenges experts may gain an advantage by including a web search in their deliberations.

Current search engines offer unprecedented accessibility at no direct cost to the enquirer, an all purpose user interface, and an immediate return of information on almost any topic. Most impressive of all is the volume of new material that is indexed automatically every day. However, the current web is based on documents in HTML (hypertext markup language) format, and thus has two limitations. Firstly, search engines cannot automatically index resources in other formats; for example, information in relational databases. Secondly, HTML indicates only how a document should be rendered, not what it means. Search engines

cannot identify which words in a document represent symptoms, diagnoses, drug names, or parts of the body, let alone reason about these concepts.

In fewer than 10 years the web has already had a considerable impact on clinical medicine. Some argue, however, that current web technologies are merely a stepping stone towards a much more powerful infrastructure—the semantic web—which aims to create a universal medium for information exchange by putting documents with computer processable meaning on the worldwide web. The semantic web can overcome current limitations and offer much stronger support for tasks such as medical diagnosis.³

The development of the semantic web is already under way. Two key technologies are RDF (resource description framework) and OWL (web ontology language).⁴ RDF allows future web search engines to access data and resources in addition to HTML documents. OWL is an international standard language for building web enabled ontologies—that is, domain knowledge that can be downloaded freely and used by computer systems such as search engines.

Web enabled medical ontologies and the semantic web will be important resources for clinical medicine in the next decade. Special interest groups have been established (for example, W3C Semantic Web Health Care and Life Sciences Interest Group; www.w3.org/2001/sw/hcls/) and several targeted ontologies have been constructed (for example, the National Cancer Institute thesaurus; www.mindswap.org/2003/CancerOntology/), but a much wider coordinated effort is needed.

No clinician diagnosing patients should feel insecure about the findings reported by Tang and Ng. Their study should encourage us, however, to think about strategies for maximising the benefits in daily clinical practice of the evolving semantic web.

Martin Gardner *research fellow*

(martin@dcs.gla.ac.uk)

Department of Computing Science, University of Glasgow, Glasgow G12 8QQ

Competing interests: None declared.

1 Tang H, Ng JHK. Googling for a diagnosis—use of Google as a diagnostic aid: internet based study. *BMJ* 2006 doi: 10.1136/bmj.39003.640567.AE.

2 Researchers impressed by Google's medical diagnoses. *Independent* 10 November 2006.

3 Berners-Lee T, Hendler J, Lassila O. The semantic web. *Sci Am* 2001;May:34-43.

4 Shadbolt N, Hall W, Berners-Lee T. The semantic web revisited. *IEEE Intelligent Syst* 2006;21:96-101.

doi: 10.1136/bmj.39044.369745.BE

Sexual relationships between doctors and former patients

New guidance gets the balance right in stopping short of a complete ban

In new guidance, the General Medical Council (GMC) has warned doctors to think long and hard before embarking on a sexual relationship with a former patient. It has not introduced a blanket ban, which might have been vulnerable to a human rights challenge, but it is far from permissive. The guidance says such a relationship “may be inappropriate irrespective of the length of time elapsed.”¹ The guidance will not please everyone.

Consider the general practitioner in a remote rural practice. The edict could cast the shadow of inappropriate behaviour across any future partner he or she may meet. Surely the medical oath did not include a vow of chastity? Previously the GMC prohibited only relationships with current patients. So what of those relationships already under way? Are these now subject to suspicion? Should doctors in such relationships, as the guidance infers, discuss their relationships with a member of the GMC standards and ethics team?

Clearly the focus is on vulnerable patients. Behind the guidance can be felt the pressure of the inquiries into Clifford Ayling, William Kerr, and Michael Haslam. In these cases, predatory doctors sexually exploited vulnerable patients entrusted to their care.^{2,3} A balancing act is needed then, which acknowledges the freedoms of competent adults to form relationships but also protects the interests of vulnerable patients. Has the GMC got the balance right?

Traditionally the doctor-patient relationship has been seen as characterised by an imbalance of power, with the doctor in the position of authority. “A relationship between a doctor and a patient is never really equal” argues the GMC’s president, Graeme Catto.⁴ By prohibiting behaviour that is not in the best interests of patients, codes of ethics have enabled doctors to concentrate on doing the best for their patients. In this way, the power of the doctor is harnessed to the good, engendering trust and maximising therapeutic outcomes. Professionalism demands that both parties are protected by clear and mutually recognised boundaries. The ethics department of the British Medical Association has advised doctors on how to manage intrusive patients—how to deal with stalkers and those who shower doctors with declarations of undying love.

But isn’t such a take on the doctor-patient relationship out of date, even paternalistic? Online resources have reduced the information gap between doctors and patients, patient autonomy is greater than ever before, and not everyone who walks through a consulting room door is a shrinking violet. Some doctors feel that the shoe is now firmly on the other foot. The classic paradigm, with the single handed (usually male) practitioner exploiting vulnerable female patients is less likely today because doctors work in multidisciplinary teams, where such behaviour would be exposed.

It is also easy to think of examples that verge on the absurd; should junior doctors treating fellow medics in accident and emergency be forever precluded from dating them? As the GMC now explicitly prohibits relationships with former patients who were “vulnerable” at the time—itsself an interpretative nightmare—is anything additional required? As the patient as consumer comes of age, and the power and status of professionals wanes, is Graeme Catto right or is the doctor-patient relationship now a meeting of equals?

As with so many questions that arise in ethics, the honest answer must be, it depends. Recent history is sadly strewn with sobering examples of doctors abusing their power—Rodney Ledward, Harold Shipman, Peter Green. If so many doctors had not abused their power in the past, calls for restraint would not be as loud. Irrespective of the pressure that the softening of traditional hierarchies of authority is bringing to bear, the doctor-patient relationship remains a kind of exemplar. Patients are often vulnerable when they visit doctors. They can be sick, distressed, and disorientated, even if they express their need in an aggressive or overconfident way.

Patients may need to reveal the most private information. Without the expectation that boundaries will be observed and trust respected, they may be less forthcoming, and patient care will suffer.

Information trawled from the internet will never be a substitute for informed professional judgment. It is for the interpretation of such information as much as for its provision that we rely on professionals. So it is a special kind of relationship, rooted in trust as much as contract, governed by professional boundaries that protect doctors as well as patients, and subject in complex ways to the ebb and flow of power.

The GMC has a difficult job to do. In falling short of a blanket ban, in recognising that there will always be exceptions, it has made a wise choice. In case you think this is a recent problem, consider the words of Hippocrates from the fourth century BC, “Into whatever house I enter, I will go into them for the benefit of the sick and will abstain from mischief and corruption and from the seduction of females or males, of freemen or slaves.”

Julian Sheather *senior ethics advisor*

(JSheather@bma.org.uk)

Medical Ethics Department, British Medical Association, BMA House, London WC1 9JR

Competing interests: None declared.

1 General Medical Council. *Maintaining boundaries*. London: GMC, 2006. www.gmc-uk.org/guidance/current/library/maintaining_boundaries.asp.

2 Department of Health Committee of Inquiry. *Independent investigation into how the NHS handled allegation about the conduct of Clifford Ayling*. London: DOH, 2004. www.dh.gov.uk/assetRoot/04/08/90/65/04089065.pdf.

3 HM Government. *The Kerr/Haslam inquiry*. Norwich: Stationery Office, 2005. www.official-documents.co.uk/document/cm66/6640/6640.pdf.

4 Eaton L. Doctors are warned against sex with former patients. *BMJ* 2006;333:876.

doi 10.1136/bmj.39042.529641.BE

BMJ 2006;333:1132

Donor breast milk banking

Unregulated expansion requires evidence of benefit

Mother's milk is recommended for all babies, but mothers of preterm babies and other babies in intensive care are often unable or too ill to provide enough milk for their baby's needs. Mother's milk protects against infection and enhances neurocognitive development; it may also improve enteral tolerance and long term metabolic health as well as reduce asthma and other atopic disorders. Donor breast milk from a human milk bank and formula are options to make up the shortfall. The extent to which pasteurised donor breast milk retains the biological properties of mother's milk is uncertain, and its place in present day neonatal intensive care is unclear.

Donor human milk is not the optimal food for preterm growth.¹ This is especially a problem because preterm infants also have limited tolerance for large volumes of milk and because infants who receive milk by gavage or spoon, unlike those who suckle at the breast, cannot regulate energy intake.² The protein and mineral content of donor milk can be increased by breast milk fortifier or formula, but this is an imprecise science as the composition of human milk is very variable. Composition alters with maternal diet; the energy content of milk is lower at the beginning of a feed than at the end and in drip breast milk than in expressed breast milk. Donors have also

usually delivered at term or have been lactating for some time, both of which result in lower nutritional content. Milk banks in the United Kingdom do not determine the nutritional content of donor breast milk. For these and other reasons it can be difficult to achieve an adequate nutritional intake in preterm infants.³

Human milk can transmit infection. HIV is the most well recognised viral pathogen, but transmission of cytomegalovirus is not uncommon, and raw breast milk can cause serious and even fatal illness in preterm babies.⁴ Pasteurisation (62.5°C for 30 minutes) is necessary but this destroys or reduces many of the unique biological properties for which human milk is valued.

In contrast to blood banks, which are tightly regulated, human milk banks in the UK vary in size and operating procedures and are unregulated, except for the voluntary adoption of standards of practice laid down by the UK Association of Milk Banks (www.ukamb.org). In Canada, Mexico, and the United States the Human Milk Banking Association of North America (www.hmbana.org) has different standards. For example, in the US milk donations are not accepted from women who were in the UK for more than three months, or in Europe for more than five years, between 1980 and 1996, presumably because of

MAURO FERRARIO/SPL

the dangers of Creutzfeldt-Jakob disease. In the UK, unlike the rest of the world, donor breast milk is not pooled, but is prepared in aliquots from individual donors.

What evidence is there to support the use of donor breast milk? A meta-analysis of trials in preterm newborns, all conducted in the early 1980s, showed that donor breast milk reduced the risk of necrotising enterocolitis—a serious inflammatory condition of the bowel—when compared with formula.⁵ Infant growth was slower, however, and benefit was seen only when breast milk or formula was the sole source of nutrition. Current practice aims to use donor milk as a supplement to mother's milk and not as sole diet. Only two studies have compared outcomes when either donor milk or formula was used as a supplement to mother's milk.^{6,7} Neither found a significant difference in the rate of necrotising enterocolitis. For other less frequent indications, and for long term outcomes, justification for using donor milk is anecdotal. Thus, the role of donor breast milk in current neonatal practice remains to be established.

Where does this leave us? Human milk banks around the world have arisen through the voluntary efforts of committed individuals. Their altruism is undeniable, but unregulated expansion of human milk banking requires evidence of benefit. Donor breast milk is expensive, and although donors are unpaid the cost of UK human milk banking (£30–150 per litre) is probably an underestimate of the full cost to the National Health Service.⁸

These uncertainties are reflected in the divergent attitudes of clinicians towards the use of donor milk and compounded by variations in acceptability by different communities.⁹ Research is needed in four key areas: to determine whether donor milk, when used as a supplement to mother's milk, has short term or long term benefits (or both) for preterm and growth restricted newborns, who are at particular risk of necrotising enterocolitis and adverse neurodevelopmental outcome, and for other groups such as babies who have undergone surgery, who are at high risk of milk intolerance; to conduct an economic assessment of the cost of human milk banking to the NHS; to

evaluate counselling and practical support to enhance successful lactation in mothers who deliver preterm¹⁰; and to explore cultural and social attitudes to donor breast milk.

The UK has 17 human milk banks and some 255 neonatal units. If donor breast milk is beneficial, clinical guidelines should reflect objective evidence, access should be equitable, and milk banking procedures should be consistent. If it is not beneficial NHS resources might be better directed towards supporting mothers' own lactation.

Neena Modi *professor of neonatal medicine*

(n.modi@imperial.ac.uk)

Division of Medicine, Chelsea and Westminster Campus, Imperial College London, London SW10 9NH

Competing interests: The author has received research funding in the past five years from Action Medical Research, the Wellcome Trust, Numico Research, London Specialised Commissioners, Westminster Children's Research Fund, Hospital Infection Society, Portland Hospital, Bailey Thomas Charitable Fund, Institute of Obstetrics and Gynaecology Trust, and March of Dimes. She is chief investigator for planned and current trials involving newborn feeding.

- 1 Heiman H, Schanler RJ. Benefits of maternal and donor human milk for premature infants. *Early Hum Dev* 2006 Oct 19 [Epub ahead of print].
- 2 Dewey KG, Heinig MJ, Nommsen LA, Lonnerdal B. Maternal versus infant factors related to breast milk intake and residual milk volume: the DARLING study. *Pediatrics* 1991;87:829–37.
- 3 Cooke RJ, Ainsworth SB, Fenton AC. Postnatal growth retardation: a universal problem in preterm infants. *Arch Dis Child Fetal Neonatal Ed* 2004;89:F428–30.
- 4 Hamprecht K, Maschmann J, Vochem M, Dietz K, Speer CP, Jahn G. Epidemiology of transmission of cytomegalovirus from mother to preterm infant by breastfeeding. *Lancet* 2001;357:513–8.
- 5 Boyd CA, Quigley MA, Brocklehurst P. Donor breast milk versus infant formula for preterm infants: a systematic review and meta-analysis. *Arch Dis Child Fetal Neonatal Ed* 2006 Apr 5 [Epub ahead of print].
- 6 Schanler RJ, Lau C, Hurst NM, Smith EO. Randomized trial of donor human milk versus preterm formula as substitutes for mothers' own milk in the feeding of extremely premature infants. *Pediatrics* 2005;116:400–6.
- 7 Lucas A, Gore SM, Cole TJ, Bamford MF, Dossetor JF, Barr I, et al. Multi-centre trial on feeding low birthweight infants: effects of diet on early growth. *Arch Dis Child* 1984;59:722–30.
- 8 Tully MR. Cost of establishing and operating a donor human milk bank. *J Hum Lact* 2000;16:57–9.
- 9 Ighogboja IS, Olarewaju RS, Odumodu CU, Okuonghae HO. Mothers' attitudes towards donated breastmilk in Jos, Nigeria. *J Hum Lact* 1995;11:93–6.
- 10 Sisk PM, Lovelady CA, Dillard RG, Gruber KJ. Lactation counseling for mothers of very low birth weight infants: effect on maternal anxiety and infant intake of human milk. *Pediatrics* 2006;117:e67–75.

doi 10.1136/bmj.39034.651563.80

Uninsured Americans and the new Democratic Congress

Is universal coverage now more likely?

At any point in time, around 45 million Americans have no health insurance, although only about half of them will have been in this situation for a year or more. About another 20 million formally insured Americans have limited coverage, with many subtle exclusions and coverage caps. About a quarter of these people could probably afford private coverage if it were available on reasonable terms. Most of them, however, are in families headed by one or two low income workers, and they could not afford the \$13 000 (£7 000; €10 000) or so that it costs to provide

an average family of four with good health care in the United States (and much more if anyone in the family is chronically ill).

If uninsured Americans fall critically ill they usually receive the necessary health care at the nearest hospital. Few of them, however, receive the timely primary care available to insured Americans. Furthermore, many of them are subsequently hounded by doctors,

This article was posted on *bmj.com* on 23 November 2006: <http://bmj.com/cgi/doi/10.1136/bmj.39042.375544.BE>

BMJ 2006;333:1134–5

hospitals, or bill collectors over unpaid hospital bills, which pushes a good number of them towards or into bankruptcy. After a thorough study, the prestigious Institute of Medicine of the National Academy of Sciences estimates that around 18 000 Americans die prematurely each year for want of health insurance, in a country whose Congress rushed back into session on a weekend to mandate continued life support for a brain dead patient.¹

In August 2003 I published a paper entitled "Is there hope for the uninsured?"² My answer then was "most probably not." As economic forecasts go, this one has been accurate so far. The question now is whether the Democrats' ascendancy to power in Congress requires a recalibration of that dire forecast, especially in light of the recent unveiling of a proposal for universal coverage by America's Health Insurance Plans, the national association of private health insurers in the US.³ My revised answer is "probably not," although I would love to be proved wrong by subsequent events.

It is not for want of attention that so many uninsured Americans find themselves in their predicament. Their plight has been explored in a decades' old series of conferences, town hall meetings, workshops, public hearings before Congress and state legislatures, television talk shows, and even, in recent years, exuberant nationwide observances of "cover the uninsured week."⁴ An entire industry has sprung up in the health services research community, dedicated annually to counting and categorising the number of uninsured people, arguing over the accuracy of the various counts, and debating whether uninsured people are worse off for their lack of coverage.

An equally energetic research enterprise has sprung up to create health reform proposals designed to bring universal coverage to the US. Although some of these proposals have been crafted with scholarly detachment, they mostly embody distinct political ideologies or the pecuniary interests of groups who profit from health care. The previously mentioned health insurance proposal unveiled by the health insurance industry on 13 November 2006 is a classic example of this genre. Although America's Health Insurance Plans would tolerate a modest expansion of government insurance programmes for poor Americans, for the most part the plan calls for federal subsidies towards the purchase of the private health insurance policies the association's members sell. It is, like all such plans, designed to allow the originator to do well by doing good.

It may be asked why, during more than a century with so many health reform proposals on the table at

any time, not one has become the law of the land, if only by accident. The answer seems to be twofold.

Firstly, unlike Europeans, Americans tend to believe that poverty is the product of free choice rather than lineage and bad luck. The persistence of that myth, despite empirical evidence to the contrary, has made it difficult to raise taxes for universal health insurance, even though the US has the lowest ratio of taxes to gross domestic product in the Organisation for Economic Cooperation and Development.

However, a more important obstacle to universal health insurance in America is that, although all politically powerful interest groups have their own preferred strategy for universal coverage, these groups always prefer the status quo as the alternative—a phenomenon known as Altman's law (after Brandeis University economist Stuart H Altman). Because no single plan has garnered the support of a political majority, we always revert to the current non-system mix of employer, government, and self coverage or no coverage. And Altman's law holds.

The Democrats' ascendancy to congressional power will probably trigger another national conversation on the plight of uninsured people. But even if the Democrats mustered the courage and the votes to raise taxes or enlarge the nation's already huge deficit for the sake of universal coverage, President Bush's veto pen would undoubtedly nip that idea in the bud, ironically appealing to the very budget deficit for which he is largely responsible.

Thus, it is a safe bet that any flurry of activity on uninsurance will be just another instalment in the never ending series of America's national conversations on the topic—a conversation that resembles nothing so much as the rambling of a drunken lover at a bar—big talk, little action. In health care it has become "the American way."

Uwe E Reinhardt *James Madison professor of political economy*

(reinhard@princeton.edu)

Princeton University, Princeton, NJ 08544, USA

Competing interests: None declared.

1 Institute of Medicine. *Hidden cost, value lost: uninsurance in America*. Washington, DC: National Academies Press, 2003.

2 Reinhardt UE. Is there hope for the uninsured? *Health Affairs* 2003;22:w3-390.

3 AHIP announces proposal to expand access to health insurance coverage to every American, 13 November 2006. Washington, DC: America's Health Insurance Plans. www.ahipbelieves.com/Default.aspx?tabid=65.

4 Cover the uninsured week 2006 provides platform to urge nation's leaders to take federal action. Washington, DC: Robert Wood Johnson Foundation. www.rwjf.org/newsroom/newsreleasesdetail.jsp?id=10404&gsa=1.

doi 10.1136/bmj.39042.375544.BE

Reconfiguring acute hospitals in England

Changes need to be planned, not left to the market

News p 1142

BMJ 2006;333:1135-6

The financial pressures facing the national health service (NHS) in England, coupled with changes in medical practice, are raising fundamental questions about the current configuration of acute hos-

pitals. With some services being provided in people's homes and in primary care, and others being concentrated in specialist centres, the viability of the current network of district general hospitals is in doubt.

If the closure of entire hospitals is likely to be the exception rather than the rule, many hospitals will have to reduce the range of services they provide to adjust to changes in their environment. This was acknowledged by David Nicholson, the newly appointed NHS chief executive, in a recent letter to MPs in which he emphasised, "Our aim should not ... be to preserve the status quo, but to think imaginatively about how we can unlock the resources, both in terms of money and people, to re-direct them where most benefit can be achieved for patients."

As Nicholson's letter makes clear, hospitals have been affected by long term changes in how health care is provided to patients. These changes centre on increases in day case procedures, reductions in lengths of stay in hospital, and an expansion of the work undertaken outside hospitals; for example, in the management of chronic disease. The net effect has been to reduce the need for beds in acute hospitals by enabling care to be delivered in other settings, thereby raising questions about the current organisation of services.¹

The challenge this creates is how to persuade MPs and the public that changes in hospital configuration are desirable when they entail reduced access to some specialist services. This is illustrated by events at Kidderminster Hospital in the late 1990s when its accident and emergency service was closed and it was used mainly as an ambulatory surgical centre.

Opposition to these changes at Kidderminster was led by a local general practitioner, and the local Labour MP lost the 2001 general election as a result of the campaign. The loss of this parliamentary seat by the Labour party exemplifies the political risks of promoting reforms that affect the public's access to health care.

This lesson has not been lost on the government, which now handles changes to the role of NHS hospitals more carefully. An independent reconfiguration panel has been established to advise on such changes. Moreover, Sir Ian Carruthers, until recently acting NHS chief executive, will work with health authorities on proposals for service reconfiguration.

Much of Sir Ian's time is likely to be spent ensuring that the NHS takes a strategic approach to change and follows rigorous and fair decision making processes. This includes engaging stakeholders such as local authorities and the public as early as possible, and allowing sufficient time for other options to be considered.

Experience has shown that it is essential to involve clinicians in hospitals and primary care in work on reconfiguration and to secure their commitment to change. The NHS should also make it clear why change is needed and articulate a persuasive and reasoned case to support proposals that are bound to be controversial. Failure to do so may mean that changes designed to improve the quality of care may be blocked.

In many cases reconfiguration will probably require hospitals to collaborate with each other to ensure that the public has access to a full range of specialist services on different sites. Tension exists between the need for collaboration to agree on the appropriate location of specialist services and the expectation that hospitals will compete for patients in the evolving programme of health reform in England.

In this respect, government policies have accentuated the challenges presented by long term changes in service provision.² Where hospitals are run by separate organisations competing for income in the healthcare market, duplication of services may occur on adjacent sites, and quality of care may suffer.

The government may need to review its own policies to enable necessary reconfigurations to occur. Bringing about improvements in care for patients and the public will not be achieved simply through the invisible hand of the market, despite the dominance of market based thinking in government.

In this context, the virtues of service planning should not be forgotten, particularly in dealing with hospitals that get into difficulty and in ensuring that services are concentrated in fewer centres if this improves outcomes. Recognising the need for a planned market in health care, if this is not an oxymoron, should loom large in the next iteration of health reform.

Chris Ham *professor of health policy and management*

(C.J.Ham@bham.ac.uk)

Health Services Management Centre, University of Birmingham, Birmingham B15 2RT

Competing interests: None declared.

1 National Health Service National Leadership Network Local Hospitals Project. *Strengthening local services: the future of the acute hospital*. London: Department of Health, 2006. www.nhsconfed.org/docs/strengthening_local_services_reference.pdf.

2 Ham C. Does the district general hospital have a future? *BMJ* 2005;311:1331-3.

doi 10.1136/bmj.39044.592662.BE