in any advertisement for any other article or service or in relation to any person. Hence this advertisement would be illegal if printed in any Irish publication.

I would never dare to suggest that the *BMJ* should consider Irish law (or, indeed, the laws of other countries) in deciding how and what to publish. But perhaps the journal might look closer at its own advertising practices. An editorial last year called for a total ban on tobacco advertising,¹ and though I congratulate the journal on having published this, might it not be a good idea for it to practise what it preaches?

FENTON HOWELL

European Medical Association Smoking or Health,
Laytown,
County Meath,
Republic of Ireland

1 Vickers A. Why cigarette advertising should be banned. *BMJ* 1992;304:1195-6.

Advertiser's response

EDITOR,—We believe that our advertisement reflects the fact that Zestril can be effective for a broad range of patients. It is not intended to condone smoking or to suggest that the potential benefits of Zestril are enhanced by smoking.

We believe that, when faced with hypertensive patients, most doctors would combine advice on factors such as smoking, diet, and exercise with drug treatment. We also believe that most doctors recognise that patients are sometimes slow or unwilling to comply with such advice. Indeed, for many years this company has provided, as a service to the medical profession, a range of educational materials aimed at helping patients modify their lifestyle.

Before this company releases any new advertisements we ensure that they are ethically correct, relevant, and appropriate. To assist us in this we also carry out market research, in which the advertisement is shown to medical practitioners and the possibility of any negative implications should be determined. None of the inferences mentioned by Fenton Howell were highlighted in this research.

Since the advertisement first appeared in January we have had no other comments of this or a similar nature. We endorse Howell's commitment to eradicating smoking and do not believe that this advertisement in any way compromises his goals. We hope that this letter will help allay his concerns.

MARK S JONES

Zeneca Pharma,
Wilmslow,
Cheshire SK9 5AZ

Treatment at Broadmoor Hospital

EDITOR,—We wish to comment on some of the issues raised by the report of the inquiry into the death of Orville Blackwood at Broadmoor Hospital in August 1991 and Luisa Dillner's news article.¹ We find the report's subtitle, *Big, Black and Dangerous?*, invidious, and slang such as "turbo jetting" is unacceptable in a serious report. We disagree fundamentally with the inquiry team's view that "medication is given for the pattern of behaviour, not the diagnosis." We believe that the diagnosis is paramount.

We question Dillner's uncritical acceptance that "the ethnic minority patients within the hospital are terrified that the authorities are 'killing them off.'"

This hospital regrets not only the death but also the fact that the reason for death is not yet understood despite two coroner's inquests. Nevertheless, the coroner said: "I have been impressed

by the competence of the nurses, by their devotion to duty, and their desire to act in the best interest of their patient, and it is my opinion that society owes them a debt of gratitude, that they are prepared to devote their lives to helping the least well equipped, difficult and often dangerous members of our society." Of the doctor, the coroner said: "He acted with probity, propriety and with professional rectitude, and no legitimate criticism is justified."

In our view, some of the quoted recommendations on drug treatment are simplistic. We have written elsewhere that guidelines given in the *British National Formulary* on dosages are unsatisfactory.² Monitoring of plasma concentrations of drugs is certainly not routine in clinical practice, and there can be a 100-fold variation in the plasma concentrations of most neuroleptic drugs in patients taking the same dose.³ When more than one drug is prescribed the assays interfere with each other. The reported therapeutic plasma concentrations for some antipsychotic drugs cannot by any means be regarded as established.

Professor Malcolm Lader's statement that "these drugs probably kill one patient every two weeks" needs urgent investigation. Major questions about drug treatment remain unanswered, and action is required. But psychiatric patients have a considerably increased mortality as a result of untreated illness. We trust that the title of Dillner's article—"Inquiry says depot injections can kill"—will not spread unnecessary fears among patients in the community and induce them to stop their drug treatment.

ANN BARKER
DILYS JONES
ALAN FRANEY

Broadmoor Hospital,
Crowthorne,
Berkshire RG11 7EG

- 1 Dillner L. Inquiry says depot injections can kill. *BMJ* 1993;307:641. (11 September.)
- 2 Barker A. An open letter to the executive editor of the *British National Formulary*, *Psychiatric Bulletin* 1993;17:557.
- 3 Van Putten T. Neuroleptic plasma levels in treatment-resistant schizophrenic patients. In: Angrist B, Schulz SC, eds. *The neuroleptic nonresponsive patient: characterization and treatment*. Washington, DC: American Psychiatric Press, 1990.

Psychotherapy

Is wanted, needed, and valued in the NHS

EDITOR,—We are surprised that Tom Fahy and Simon Wessely seem so poorly informed about psychotherapy services in the NHS and purchasers' views of them.¹ We take issue with several of the points they make.

The authors assume that purchasers and managers will have difficulty in finding evidence for the effectiveness of dynamic psychotherapy and therefore will be unwilling to pay for such services. In fact, the experience of psychotherapy services nationally since the reform of the health service has been that purchasers, managers, and patients seem to regard dynamic psychotherapy as a valued and wanted component of mental health services, as has been shown in surveys of general practice.² Though the outcome of such psychotherapy for complex problems is not easily measured in controlled trials looking at simple outcome variables, the authors' reference to Andrews's assertions³ rather dismisses the large body of evidence to which they refer earlier,⁴ evidence that does support the effectiveness of dynamic psychotherapy.

Fahy and Wessely then promulgate several misconceptions about the nature of psychotherapy services in the NHS. Firstly, the most common model of dynamic psychotherapy in the NHS entails once weekly treatment for nine to 12 months (that is, 30 to 40 hours). It would be unusual for patients in the NHS to be seen for as

long as 330 hours outside a specialist training centre. Secondly, the patients seen by dynamic psychotherapists working in the NHS do not conform to the caricature implied by the authors' phrase "the traditional clients of their colleagues in private practice."⁵ Many of the patients have been subject to deprivation and abuse in childhood and present with severe personality disorders or other serious long term difficulties. Such patients often make heavy demands on general practices and other NHS services, but few have access to the specialist understanding they require. Thirdly, there is a long record of dynamic psychotherapists treating the mental health problems encountered in general hospitals.⁶ The authors' place of work, King's College Hospital, has a history of such work, with dynamic psychotherapists helping to manage abnormal illness behaviour.

The scope for such psychodynamic approaches to treatment is hindered by the shortage of psychotherapists working in the NHS. At present there are only about 100 whole time equivalent medical consultant psychotherapists in Britain, who serve less than half of the districts. An adequate response by providers to the "new rigours of the purchaser-provider split" might thus be to rectify this inadequate provision.

NOT TEMPLE
MPH PATRICK

Tavistock Clinic,
London NW3 5BA

- 1 Fahy T, Wessely S. Should purchasers pay for psychotherapy? *BMJ* 1993;307:576-7. (4 September.)
- 2 Morton MJ, Staines J. GP use of psychotherapy services. *Psychiatric Bulletin of the Royal College of Psychiatrists* 1993;17:526-7.
- 3 Andrews G. The essential psychotherapies. *Br J Psychiatry* 1993;162:447-51.
- 4 Milton J. *Presenting the case of psychoanalytic psychotherapy services. An annotated bibliography. Discussion document.* London: Association for Psychoanalytic Psychotherapy in the NHS and Tavistock Clinic, 1992.
- 5 Schoenberg, P. Psychosomatic disorders. In: Holmes J, ed. *Textbook of psychotherapy in psychiatric practice*. London: Churchill Livingstone, 1991.

Short term therapies not always adequate

EDITOR,—Tom Fahy and Simon Wessely emphasise the importance of rigorous evaluation of psychotherapies available as NHS treatments.¹ Some approaches, however, are more amenable to scientific evaluation than others. In general, psychotherapies that have a brief focus and a rationalist-reductionist approach use methods that are easy to describe to deal with objectives that can be specified and measured. If life was always this simple there would be no further problem and no place for more complex or sophisticated psychotherapeutic approaches. Indeed, for most patients requiring NHS counselling or psychotherapy these short term, clearly focused approaches are best and should be readily available from primary care and community mental health teams.

An appreciable number of patients, however, have longstanding problems, causing major functional impairment and giving rise to repeated presentation with intractable symptoms in general practice as well as in mental health and general hospital services. Examples are people with chronic neurotic problems, intractable eating disorders, and borderline personality disturbances and victims of prolonged abuse in childhood. Though some of these people may benefit from short term intervention, many need long term therapy that is flexible and receptive to their needs and free of the implicit impatience of short term approaches.

Psychotherapists in Britain generally accept the need for a range of approaches to be made available in each district to meet the differing needs of patients. There is, however, a possibility that the present overcommercial culture of the NHS will lead to a "pile 'em high, sell 'em quick" approach