breathing air—a considerable reduction from normal and one that should correspond with an arterial oxygen pressure of about 8.7 kPa. The boy's thermal injury from steam would probably have been limited to the airway above the larynx by reflex closure of the vocal cords, so any alveolar-arterial oxygen gradient would have been small. Thus the hypoxaemia must have been caused by hypoventilation. The alveolar gas equation, although a crude estimate when a respiratory steady state is improbable, indicates a carbon dioxide pressure of over 8 kPa and therefore appreciable hypoventilation.

This shows how a pulse oximeter can help assess hypoventilation in such a patient. Endotracheal intubation of patients with upper airway obstruction is seldom easy or free from hazard, and non-invasive pulse oximetry is valuable to confirm the need for such action.

A J BALL
M B HOUGH
N J H DAVIES

Shackleton Department of Anaesthesia,
Southampton General Hospital,
Southampton SO9 4XY

- 1 Hutton P, Clutton-Brock T. The benefits and pitfalls of oximetry. *BMJ* 1993;307:457-8. (21 August.)
- 2 Williams EGN, Dymock M. Acute airway obstruction after aspiration of boiling tea from teapot spout. *BMJ* 1993;307:923. (9 October.)

Sanctions delay aid to the former Yugoslavia

EDITOR,—The Norfolk and Norwich Novi Sad Association was set up to further the twinning links that have existed for about 30 years between Norwich and Novi Sad in what is now the Federal Republic of Yugoslavia. For the past 18 months the association has been trying to supply help to the refugees in Novi Sad. Our next load will be 2 tonnes of medicines (including medicines to treat tuberculosis), medical supplies, baby food, and so on.

The present sanctions have made obtaining the necessary permissions much slower and more difficult than previously. Everything that we take, from purchased medicines to back copies of the *BMJ*, has to be cleared by the United Nations in New York. To get this clearance we have to send in detailed applications to the Department of Trade and Industry in London. The department, having checked the application, forwards it to the UN. The clearance procedure at the UN currently takes six to 10 weeks. Once clearance has been obtained the Department of Trade and Industry issues the appropriate documents to allow the goods into the former Yugoslavia.

In addition, we have to get letters of safe conduct from the Hungarian embassy and the embassy of the Federal Republic of Yugoslavia and entry visas. The original application has to state the date on which we propose to enter the Federal Republic of Yugoslavia (though letters of clearance are valid for three months). We therefore have to estimate three months in advance what we are likely to collect. This also means that some of the baby food we are given has almost passed its "best before" date on delivery.

All the goods we take out are delivered to the Red Cross of Vojvodina in Novi Sad. We have established a good relationship with the officials of the Red Cross and are satisfied that everything we take is allocated to a suitable recipient. Having experienced generous hospitality and kindness in Novi Sad in the past, we are happy to try to help our friends to care for the refugees and the disadvantaged in society.

The present situation in the former Yugoslavia undoubtedly has several causes, but, as Mary E Black says,¹ the sanctions are making things worse

because of the delays they cause. People will die from lack of food and proper health care before this winter is over.

PETER BECKLEY
DIANA BECKLEY

Norfolk and Norwich Novi Sad Association,
Rockland St Mary,
Norwich NR14 7BX

- 1 Black ME. Collapsing health care in Serbia and Montenegro. *BMJ* 1993;307:1135-7. (30 October.)

Giftedness

EDITOR,—It is unfortunate that in a recent editorial all the 20 references are to American publications and five are by the editorial's author, John F Feldhusen.¹ Yet a great deal of good, current work on giftedness in Europe and elsewhere could have been mentioned. The editorial's content shows the same restriction: Feldhusen states, for example, that gifted children are often found to be bored at school and that accelerated teaching is valuable, but this has been shown only in parts of the American educational system.

Feldhusen wonders what happens to gifted children when they grow up and then responds by citing just one study, by Terman and Oden, which was begun in the 1920s in California and is methodologically unsound. There are now, however, several longitudinal studies in many parts of the world that have reached different conclusions.^{2,3} We know, for example, that giftedness in children is limited: no child has ever effected the reorganisation of a domain of knowledge, and many people who have made such changes as adults did not seem to be gifted as children. We also know from studies of separated twins that there is a considerable genetic influence on abilities⁴ and that how children move from potential to achievement depends to a large extent on the educational and emotional support they receive from the people who care for them.

Studies of the development of gifted children are valuable because they throw light on the development of all children. Adequate education is of primary importance, both for gifted children and for society, and the problems of providing this in the normal school system are of concern to many education authorities. Sound and practical answers are emerging from research, but they need a more up to date and open minded reading of the literature than Feldhusen's editorial presents.

JOAN FREEMAN

European Council for High Ability,
London W1H 1RE

- 1 Feldhusen JF. Giftedness. *BMJ* 1993;307:1088-9. (30 October.)
- 2 Freeman J. *Gifted children growing up*. London: Cassell, 1991.
- 3 Heller KA. The nature and development of giftedness: a longitudinal study. *European Journal for High Ability* 1991;2:174-8.
- 4 Thompson LA, Plomin R. Genetic influences on cognitive ability. In: Heller KA, Monks FJ, Passow AH. *International handbook of research and development of giftedness and talent*. Oxford: Pergamon, 1993;103-13.

Antenatal screening for genetic disease

EDITOR,—Jennifer Dennis and colleagues' letter¹ referring to the paper by Statham and Green on women's experiences after serum screening for Down's syndrome² and Marteau's accompanying editorial³ describes problems that have also been experienced by the Child Growth Foundation.

We are receiving increasing numbers of calls from parents distressed by a possible or definitive prenatal diagnosis of Turner's syndrome. After identification of associated abnormalities by trials or screening procedures or confirmation by amnio-

centesis, these parents are confused and distressed. In most cases the parents' distress seems to have been exacerbated by a lack of informed support. The information they have received on Turner's syndrome has mostly been based on out of date textbooks and can be both incorrect and overwhelming. It should always be explained that, as with any syndrome, there can be a great variation in the degree to which people are affected. Sometimes the parents are given the diagnosis by telephone and are then left waiting for an appointment for the implications of the diagnosis to be fully explained. This delay leads worried parents to turn to relevant support groups and places an increasing strain on the voluntary sector when the responsibility should be shouldered by primary care or hospital based antenatal care and genetic services.

We realise that some health districts have implemented good informed support services, but not all have done so. Services to provide informed caring support should develop in parallel with the technology that makes it possible to diagnose disorders prenatally. Learning of a prenatal abnormality is a horrifying experience for a pregnant woman; it is essential that both parents have sufficient knowledge to make an informed decision on whether a pregnancy should be terminated and that they are supported whatever their decision may be. We must recognise the need for informed support services that keep up with the advances in technology.

VRELI FRY
JILL RUSTAD

Child Growth Foundation,
London W4 1PW

- 1 Dennis J, Sawtell M, Rutter S. Counselling needed after screening for Down's syndrome. *BMJ* 1993;307:1005. (16 October.)
- 2 Statham H, Green J. Serum screening for Down's syndrome: some women's experiences. *BMJ* 1993;307:174-6.
- 3 Marteau TM. Psychological consequences of screening for Down's syndrome. *BMJ* 1993;307:146-7.

Metered dose inhalers

EDITOR,—Are there any compelling reasons why metered dose inhalers should not be transparent, like disposable cigarette lighters? The contents could then be easily assessed with regard to both the quantity remaining and the degree of mixing.¹

AUSTEN MOORE

Walsall WS1 2DG

- 1 Williams DJ, Williams AC, Krucke DG. Problems in assessing contents of metered dose inhalers. *BMJ* 1993;307:771-2. (25 September.)

Social deprivation and health

Money must go where it is most needed

EDITOR,—Margaret Eames and colleagues indicate that the government's targets for reducing mortality from coronary heart disease and other diseases, set out in *The Health of the Nation*, could be achieved if mortality in the country's poorest areas could be reduced to the rates seen in the more affluent areas.¹

The revised strategy for promoting health and managing chronic disease in general practice ignores the need to direct resources into areas of greatest social deprivation.² Instead, the strategy is based on capitation and list size.³ Thus many deprived populations will once again lose out and be marginalised. Marsh and Channing also clearly showed that disadvantaged populations need carefully planned targeting of resources by the whole primary health care team to provide care equal to that of a more affluent population served by the same practice.⁴

In deprived areas general practitioners need to work in the community with community health workers, benefits rights workers, health promotion officers, and housing officers to ameliorate the multifactorial problems caused by socioeconomic, educational, and health deprivation. Social policies aimed at a more equitable distribution of wealth—by empowering deprived people through educational and employment opportunities—are needed to reverse the dramatic widening of the health-wealth gap.⁵

The message is clear. To reverse the marginalisation of disadvantaged areas additional resources must be targeted in health, education, and employment if the targets of *The Health of the Nation* are to have any chance of being met. Disproportionately greater resources are needed to give deprived people an average chance of health.

JOY A MAIN
PAUL MAIN

Bristol BS13 0JP

- 1 Eames M, Ben-Shlomo Y, Marmot MG. Social deprivation and premature mortality: regional comparison across England. *BMJ* 1993;307:1097-102. (30 October.)
- 2 Main JA, Main PGN. Allocating resources to doctors in deprived areas. *BMJ* 1990;299:1528.
- 3 Main JA, Main PGN. *The health divide*. *BMJ* 1993;306:273.
- 4 Marsh GH, Channing DM. Narrowing of the health gap between a deprived and an endowed community. *BMJ* 1988;296:173-6.
- 5 Smith GD, Egger M. Socioeconomic differentials in wealth and health. *BMJ* 1993;307:1085-6.

Wealth redistribution has its price

EDITOR,—Margaret Eames and colleagues have added weight to widespread observations of a positive and progressive association between higher socioeconomic position and improved health.¹ It seems that more income, wealth, and education and higher status occupations are all correlated with less illness and a lower mortality, and such improvements continue right to the top of the social scale.² So far, there seems to be no limit to the ability of social advantage to add to health and life expectancy.²

It is time that we changed our way of conceptualizing socioeconomic inequalities in health. Instead of continually harping on deprivation as a cause of sickness—where deprivation is defined as a “comparative” concept with “no specific threshold”—perhaps it would be more reasonable to stop using the word deprivation when talking about some of the most privileged and prosperous people in the history of the world. Otherwise we are put into the absurd position of asserting that (to misquote Orwell) everybody is deprived, but some of us are more deprived than others.

We might consider the reciprocal view that, instead of deprivation causing sickness, advantage causes health. This is the salutogenic model of health as outlined by Antonovsky and elaborated by M P Kelly and myself.^{3,4} The salutogenic perspective highlights several points not mentioned anywhere in the 20 October issue of the *BMJ*.

For instance, one consequence of the finding that “premature mortality increases linearly with increasing deprivation”¹ is that a policy to produce “a reversal in the dramatic upwards redistribution in wealth”⁵ would surely damage the health of the higher social classes even as it improved the health of the lower social classes. This may, indeed, be a price worth paying; but we should not ignore the fact that there is a price to pay. Redistribution removes advantage at the same time as it addresses deprivation. Wealth redistribution would not give us something for nothing: there would be health losers as well as health gainers—a situation which would not promote the Utopian goal of “allowing everyone the opportunity of achieving the maximum health.”³ Neither can we assume that “one way of achieving national targets would be to reduce geographical and social variations.”³

It is unhelpful to become fixated on “differen-

tials” and “inequalities” while downplaying the massive improvements in public health over the past decades. A more practical policy would be to concentrate on improving health wherever possible. “Everyone” will never achieve maximum health, and we should be careful not to throw away actual achievements in pursuit of unattainable ideals.

BRUCE G CHARLTON

Department of Epidemiology and Public Health,
University of Newcastle upon Tyne,
Newcastle upon Tyne NE2 4HH

- 1 Eames M, Ben-Shlomo Y, Marmot M. Social deprivation and premature mortality: regional comparison across England. *BMJ* 1993;307:1097-102. (30 October.)
- 2 Charlton BG. Is inequality bad for the national health? A reinterpretation of the evidence. *Lancet* (in press).
- 3 Antonovsky A. *Health, stress and coping*. San Francisco: Jossey Bass, 1985.
- 4 Kelly MP, Charlton BG. A scientific basis for health promotion: time for a new philosophy? *Br J Gen Pract* 1992;42:223-4.
- 5 Davey Smith G, Eggar M. Socioeconomic differentials in wealth and health. *BMJ* 1993;307:1085-6. (30 October.)

Disease registers valuable if geographically defined

EDITOR,—William F Kelly and colleagues’ analysis of data held on a hospital diabetes register illustrates the potential for using a chronic disease register to explore associations such as that between ill health and deprivation.¹ The authors analysed data relating to patients who attended a hospital diabetes centre during one year. The study, however, is flawed. The sample studied made up approximately 0.5% of the district’s population, with half of the sample being treated with insulin. Comparison with population based studies shows that the data refer to only a subset of people with diabetes in the district.²

Only with a geographically defined population register might it be possible to investigate the associations with deprivation that the authors describe. Results suggesting that diabetic patients from deprived wards were older and less likely to use insulin probably reflect nothing more than different patterns of care in general practice, with general practitioners in the least deprived wards caring for more of their older, non-insulin treated patients, who will consequently not be recorded as attending hospital. The presence of complications may be a factor in the decision to refer to hospital care, and this referral pattern may also vary among general practitioners in different areas.

The potential for research using disease registers is considerable but may be fully realised only when comprehensive, valid registers based on geographically defined populations are developed.

DAN KILLALEA
ALWYN DAVIS
GAIL HANSTOCK
JANE HANSON

Department of Public Health Medicine,
Dudley Health Authority,
Dudley DY1 2DD

- 1 Kelly WF, Mahmood R, Kelly MJ, Turner S, Elliot K. Influence of social deprivation on illness in diabetic patients. *BMJ* 1993;307:1115-6. (30 October.)
- 2 Neil AW, Gatling W, Mather HM, Thompson AV, Thorogood M, Fowler GH, et al. The Oxford community diabetes study: evidence for an increase in the prevalence of known diabetes in Great Britain. *Diabetic Med* 1987;4:539-43.

Clinical research in trust hospitals

Threatened by liability for negligence

EDITOR,—For many years our local ethics committee has accepted research proposals from NHS staff within the district on the grounds that, although the NHS had no formal indemnity for its staff who might cause injury through negligence

while undertaking research, an ex gratia payment would be most likely in the event of a claim being made.

We were aware that responsibility for such indemnity had been devolved, first to regions and then districts. We were also aware that Crown immunity had been withdrawn from hospitals. But the consequences had escaped us and they became clear only when our local trust refused to sign any agreements with external manufacturers, who were willing to cover non-negligent injury. Legal opinion had been sought and was to the effect that a trust, under its existing funding and rules of constitution, might not accept so great a liability for research, similar though it is to what happens in ordinary clinical practice.

Trusts therefore seem to have responsibility for something that they are neither funded for nor allowed to undertake—the provision of cover for negligent injury arising from research, whether alongside or outside ordinary clinical practice.

The entailments of this are now clear. As local research ethics committees cannot accept protocols without cover for both negligent and non-negligent injury, this means an effective end to clinical research of any kind which may incur risks of negligence within an NHS trust. And, when all the world’s a trust, that implies within the NHS as a whole.

The Department of Health was asked some time ago for guidance about this, but none has emerged in writing. We have been advised to buy insurance privately. The situation contrasts oddly with the brochure entitled *Research for Health*, published recently by the Department of Health.

D W VERE

Department of Pharmacology and Therapeutics,
London Hospital Medical College,
London E1 2AD

**We sent the above letter to the Department of Health for their response.

Pharmaceutical companies should indemnify trusts

EDITOR,—NHS bodies are not allowed to offer advance indemnity to subjects in research projects. This applies equally to directly managed units and NHS trusts. Any liability of the NHS is carried by the district health authority for directly managed units or by the NHS trust for patients participating in clinical studies in the same way as those NHS bodies carry liability for negligence arising out of normal clinical activity.

The Association of the British Pharmaceutical Industry and this department have agreed the terms of a standard form of indemnity for use in clinical trials sponsored by the pharmaceutical industry. This provides for the sponsoring company to indemnify the district health authority or trust against claims arising out of the administration of the product under investigation and associated procedures. The indemnity extends to employees of the health authority or trust but does not include an indemnity for their negligent actions. The department will be issuing guidance on the use of a clinical trial indemnity form in due course.

We are of course aware of more general concerns about the way risks arising from clinical negligence should be dealt with by NHS trusts. We have for that reason set up a joint departmental-NHS group (which includes representation from the royal colleges) to look at the possibility of establishing a mutual insurance scheme for clinical negligence claims and to consider how to manage risks and claims. We will certainly ensure that the specific concern raised by D W Vere is examined by the group.

J S METTERS

Department of Health,
London SW1A 2NS