CORRESPONDENCE

Genetic counselling and genetic registers A E H Emery, FRCPED..... 637	Orf in Britain G H Tee, FRCPATH..... 641	National Health Service Consultants' Association S P Baxter, MRCPsych..... 643
Metabolic bone disease in Asians J C Waterlow, FRCP..... 637	Carpal tunnel syndrome in the blind N H Harris, FRCS..... 641	Consultants' contract J M Cundy, FFARCS..... 643
Hazards of smallpox vaccination F Kellerman, MD..... 638	Volvulus of the small bowel in a diabetic patient H Freund, MD..... 641	Industrial action P L Hays, MB; D C Hogg, MB; Margaret C Watkinson, MB; P D Griffiths, MD; D M Bowers, LRCPED..... 644
Replacement of the knee joint J T Scales, FRCS..... 638	Porcine dermis dressing versus Bisgaard therapy for leg ulcers Christine I Harrington, MRCP..... 641	Consultants' increments A P J Ross, FRCS; M J Brown, MRCPATH..... 645
Cancer of the oesophagus K Moghissi, FRCSED..... 638	Acute gastric dilatation after trauma S U Rahman, FRCS..... 641	Madness of the profession H Cairns, MRCP..... 645
Rheumatoid atlantoaxial subluxation J A Mathews, MRCP..... 638	Effects of pethidine on the newborn R S Brown, FCP(SA)..... 642	Juniors' contract dispute R F Massey, MB; K J J Tayton, FRCS, and others..... 646
Oral irritation with gentian violet J L Verbov, MD..... 639	Extrapyramidal side effects of lithium treatment B Johnels, MD, and others..... 642	United profession E R C Walker, FRCPED..... 646
Hypoglycaemia in children undergoing operations E N S Fry, FFARCS; S P Deacon, MB..... 639	Pyloric stenosis and sustained-release iron tablets I P Carne-Ross, MB..... 642	Private practice and the NHS K T H Moore, FRCS..... 646
Mechanism of action of antiallergic drugs J C Foreman, PHD, and L G Garland, PHD..... 640	A hazard of apple-eating T Havland, MD, and G Frithz, MD..... 643	Medical manpower and hospital staffing I M Librach, MB..... 646
Thyrotoxic vomiting J M Temperley, MD..... 640	Doctors, contraception, and sterilisation A R Hill, FRCSED..... 643	Professional unity R A Keable-Elliott, FRCGP..... 646
Lung cancer and smoking: Is there proof? P R J Burch, PHD..... 640		
Out-of-hours calls in general practice H M Halle, MRCP; P I Vardy, MRCP..... 640		

Correspondents are urged to write briefly so that readers may be offered as wide a selection of letters as possible. So many are being received that the omission of some is inevitable. Letters should be signed personally by all their authors.

important if a genetic register system is to be really effective in helping prevent serious genetic disease within the community.

ALAN E H EMERY

University Department of
Human Genetics,
Western General Hospital,
Edinburgh

¹ WHO Technical Report Series No 497. Geneva, World Health Organisation, 1972.

² Emery, A E H, et al, *Journal of Medical Genetics*, 1974, 11, 145.

³ *Registers for the Detection and Prevention of Genetic Disease*, ed A E H Emery and J R Miller. Symposia Specialists Medical Books. New York, Stratton International. In press.

Genetic counselling and genetic registers

SIR,—Dr A J Caro and his colleagues (14 August, p 420) have recently emphasised the need for detecting individuals at risk of having children who might develop Huntington's chorea so that they may be given genetic counselling. While I entirely agree with these sentiments, the problem is in fact much wider than this, for it exists with many other dominant disorders with late onset (such as polyposis coli, polycystic kidney disease, and myotonic dystrophy) and several X-linked disorders (such as haemophilia and Duchenne muscular dystrophy) where there are often many family members at risk of having affected children who may be unaware of this. In a study of the families of individuals referred to this department for genetic counselling over a four-year period we estimated that there were 950 relatives at high risk (greater than 1 in 10) of having a child with a serious genetic disorder, yet of these only 138 (15%) had had any genetic counselling. Further, 81 affected children were born to parents who a priori were at high risk of having affected children and at least a proportion of these might have been prevented by genetic counselling.

From these findings it would seem reasonable to assume that the extent of the problem is much greater than is reflected by the number of individuals who attend a genetic counselling clinic. There is at present no defined procedure for ascertaining and follow-

ing up such individuals. Herein lies the value of a genetic register system which, for this reason, has been advocated by a WHO scientific group.¹ Such a system was first established in Edinburgh in 1970 under the acronym RAPID (Register for the Ascertainment and Prevention of Inherited Disease)² and has been fully operational for at least a year. For ease of storage, updating, and retrieval of family data the system has been computerised, and strict safeguards for confidentiality have been incorporated into the system. Individuals are included in the register system only with their full approval and written consent (or that of their parents). No relatives are contacted without the permission of the index case and also the relatives' family doctor. The latter is considered especially important as there may be factors unknown to the medical geneticist or index case which would make it imprudent or unnecessary to contact certain family members. Particularly important in such a system is the maintenance of data on individuals who are at high risk of having affected children or of developing a genetic disorder but as yet are too young for counselling.

In the past few years a number of similar genetic register systems have been established in various countries, including Canada, Norway, the United States, and Belgium.³ The latter is particularly interesting because it has been organised on a national basis. This is

Metabolic bone disease in Asians

SIR,—Your leading article on this subject (21 August, p 442) was welcome in drawing attention to a problem which should be a primary concern to all those working in public health. As the article points out, Asians appear to be peculiarly susceptible to rickets, a deficiency disease for which we have long known the cure and the means of prevention. The situation referred to in Bradford was an estimate based on a study of hospital admissions done over four years ago, but there are many other fully documented examples (such as that of Goel *et al*)¹ of a state of affairs which should not be allowed to continue.

Fortification of chapati flour is one way of tackling the problem. However, this solution has its limitations: very young children, who are the members of the family probably most in need of extra vitamin D, would receive least of the fortified flour, while there are Asians who eat rice in preference to chapatis. There

may well be mistrust of a modification to a familiar dietary staple. Furthermore, other sections of the population, such as the house-bound elderly and the chronic sick of all ages, are also at risk of vitamin D deficiency through lack of exposure to sunlight, and it is possible that some other item of the diet, such as milk, would be a more appropriate vehicle for fortification and would confer a benefit over a wider range of consumers.

Fortification of flour is legally and administratively complicated. One difficulty is that we do not know the extent of the problem and hence the benefit likely to be conferred by fortification compared with other measures. Your leading article refers, reasonably enough, to the clinical cases as the tip of an iceberg. We do not know how big the iceberg is on a national scale. For example, I know of no recent information about the prevalence of rickets in the very large Asian communities of southern England comparable with the studies which have been made in the midlands and the north. There are no clear criteria for the biochemical diagnosis of hypovitaminosis D. Certainly, as Goel *et al*¹ point out, raised serum alkaline phosphatase by itself is not adequate evidence of deficiency.^{2,3} Measurement of 25-hydroxycholecalciferol concentration in the serum offers promise, but there is at present no standardisation among the different groups who make this measurement so that we do not have "normal" or reference levels which could be used in community surveys.

This emphasis on what we do not know is not a plea for doing nothing, pending more research. I hope that your leading article will stimulate more inquiries by paediatricians, community physicians, and others who come into contact with the groups at risk. I fear, however, that concentration on one particular remedy—fortification of chapati flour—which cannot be put into effect quickly may divert attention from other measures already available which could be immediately effective. As you suggest, the school health services and community health teams, in collaboration with the Indian and Pakistani community leaders, could play a large part here. Goel *et al*¹ have already advocated that a supplement should be widely available for children up to the age of 16 years and this should be extensively advertised. It is then for the local authorities in areas where immigrants are concentrated to ensure that advantage is taken of it.

J C WATERLOW

Department of Human Nutrition,
London School of Hygiene and
Tropical Medicine,
London WC1

¹ Goel, K M, *et al*, *Lancet*, 1976, 1, 1141.

² Round, J M, *British Medical Journal*, 1973, 3, 137.

³ Stephen, J M L, and Stephenson, P, *Archives of Disease in Childhood*, 1971, 46, 185.

Hazards of smallpox vaccination

SIR,—It is regrettable that Mr G T Watts (28 August, p 530) should give bad advice, contrary to the evidence regarding the epidemiology of smallpox. Sir James Howie's view (24 July, p 217) that "smallpox is definitely on the way out" is not optimistic; it is realistic and based on facts.¹ On the other hand, Mr Watts's misconception of "areas where the disease is still endemic" is sheer fantasy. Today such areas do not exist.² The last remaining endemic focus in the world is in

Ethiopia and as the mortality rate is about 1% the type of disease is more likely to be variola minor. Far from having a "large social interchange between this country and areas where the disease is still endemic"—as Mr Watts would like us to believe—the exact opposite is true: it would be practically impossible for the ordinary traveller to get there. WHO observers who carried out a surveillance of smallpox in Ethiopia since 1971 pointed out that the eradication of smallpox in Ethiopia has been less rapid than in other countries because the endemic foci are in "remote and inaccessible" parts of the country.³

It is unfortunately true that "many countries will insist on vaccination," but they should not and it is our duty as a profession to resist this demand, which is based on out-of-date, ill-informed, inexpert advice, and support with all our might Sir John Howie's wisdom in urging us not to indulge in potentially dangerous vaccination aimed at the prevention of a non-existing disease.

F KELLERMAN

Colchester

¹ *British Medical Journal*, 1976, 1, 1219.

² *WHO Chronicle*, 1976, 30, 152.

Replacement of the knee joint

SIR,—I read with interest your leading article on this subject (21 August, p 443).

It is correct to say that replacement of the knee joint is not being carried out as frequently as total hip replacement. A prosthesis must relieve pain, provide a useful range of joint movement, provide stability, and correct deformity. However, it is open to debate whether the movement provided by the artificial joint should simulate anatomical joint movements. The knee joint is essentially an unstable joint configuration relying more on muscle and ligamentous control for movements than a ball and socket joint. To achieve maximum stability there must be some means of locking the joint in full extension, and this probably is best achieved in the case of a joint stabilised by soft tissues by the use of a torsional system such as the external rotation which occurs at the knee joint. Probably one of the functions of the semilunar cartilages is to allow this torsion to occur under load without undue shear stresses developing in the tibial and femoral cartilages.

A number of hinged knee joints which allow movement around a fixed axis have been functioning very satisfactorily for a number of years. The joints which allow gliding movements on flexion and extension, with some rotation and lateral mobility, may be useful for patients in whom there is no gross joint destruction. However, in patients with gross destruction of the joint there is also generally torsion of the tibia. This anatomical change of the tibia, which develops over the years, results in muscle forces of such a magnitude acting in such directions that they may necessitate a fixed hinge system to prevent the recurrence of a deformity or undue wear on the components of a total knee replacement rather than a prosthesis of the semi- or non-constrained type. The total knee replacement consisting of two parts which are linked in such a manner as to provide freedom of movement in three planes may be suitable for certain cases, provided material failure does not occur in the years ahead.

In no way am I attempting to decry the pioneering work of those such as Sheehan, Attenborough, Deane, or Gschwend *et al*, who are all concerned with the development of "physiological" knee joints. However, it would be unwise for those who are concerned with advising patients on replacement of the knee joint to do other than explain to them that there is a variety of designs available and that no one type is at the present time demonstrating an overwhelming clinical advantage.

JOHN T SCALES

Department of Biomedical Engineering,
Institute of Orthopaedics,
Royal National Orthopaedic Hospital,
Stanmore, Middx

Cancer of the oesophagus

SIR,—Your leading article on this subject (17 July, p 135) is misleading in that it conveys the message that radiotherapy is preferred for mid-oesophageal cancer. As no reference is provided one wonders whose opinion is quoted.

Your readers may be interested to know that in a recent symposium in Valencia on this subject a questionnaire was sent by the organiser, Dr F Paris, to more than 80 surgeons in Europe, including some of the surgeons carrying out oesophageal surgery in Britain. Two surgeons used radiotherapy for mid-oesophageal carcinoma, but only when the tumour histology was one of anaplastic carcinoma; three others used radiotherapy with surgery. No one appeared to prefer radiotherapy as the method of choice.

You also omit to state the morbidity and mortality of radiotherapy.

K MOGHISSE

Cardiothoracic Surgery Unit,
Castle Hill Hospital,
Cottingham,
N Humberside

*.*Presumably it is acceptable that if a mid-oesophageal tumour cannot be resected by a surgeon the patient should be offered radiotherapy to enable him to swallow properly. If x-rays help in these extreme cases surely they are of use in earlier cases. In one series of "inoperable" mid-oesophageal carcinomas treated by radiotherapy a 20% 5-year survival was obtained.¹ It would appear that Dr Paris's questionnaire was sent only to surgeons. What would have happened if, instead, 80 radiotherapists had been asked what the ideal treatment was? The answer must lie somewhere in the middle between two extremes. Our article emphasised that good results can be obtained by experts in early cases. In Britain these specialists should be surgeons working in close co-operation with a radiotherapy department. The best results have not yet been achieved because patients are treated as a rarity in different hospitals without using all our facilities.

¹ Pearson, J G, *Clinical Radiology*, 1966, 17, 242.

Rheumatoid atlantoaxial subluxation

SIR,—I am pleased to see that continuing interest in rheumatoid arthritis of the cervical spine has stimulated you to publish a second leading article (24 July, p 200) on this topic in under 3½ years. The recent article discussed the problem in a way that suggests an experience