Today's Treatment

Diseases of the alimentary system

Significance of persistent HBs antigenaemia

I L WOOLF, ROGER WILLIAMS

British Medical Journal, 1976, 2, 807-808

It is well established that the presence of hepatitis B surface antigen (HBsAg) in blood indicates infection with the hepatitis B virus. HBsAg constitutes the surface of the 42-nm Dane particle, which may well be the complete virus, and is also the antigen of the 20-nm spherical and tubular forms that exist, probably as a result of overproduction of this surface component. HBsAg may be detected during the incubation period of acute hepatitis type B, and in cases that follow the usual course to complete recovery it has invariably disappeared three months after the onset of the illness. Subsequently the corresponding antibody (HBsAb) may be detected, which persists for months or even years. The core of the Dane particle contains a second antigen, hepatitis B core antigen (HBcAg), which has also been found in the nuclei of infected hepatocytes. Free HBcAg has not been detected in blood but its antibody (HBcAb) appears during or just before the onset of acute hepatitis type B and probably persists indefinitely.

Who are HBsAg-positive?

HBsAg, however, does persist in the serum in certain conditions. A few patients, for example, may remain HBsAg-positive after acute hepatitis type B. This occurred in 11 out of 253 such patients (4.3%) in Copenhagen, and all showed evidence of chronic liver disease.¹ In haemodialysis and renal transplantation units infection with hepatitis B virus is often followed by persistent HBs antigenaemia. This may result from an impaired immune response to hepatitis B viral antigens through either the toxic effect of renal failure in the dialysis patients or treatment with immunosuppressives in those patients who have had transplants. This persistent HBs antigenaemia is often although not invariably associated with the development of a chronic hepatitis.

Some patients with evidence of chronic liver disease, with or without a history of hepatitis, will be HBsAg-positive. The proportion will depend on the population studied. Thus with the use of a sensitive radioimmunoassay technique only 13% of a series of British-born patients with chronic active hepatitis were shown to be HBsAg-positive, whereas among a similar group of patients originating from either the Mediterranean area or the Middle East the proportion was 60%.²

Finally, HBsAg may be present persistently in asymptomatic people without a history of hepatitis or signs of chronic liver disease. Such people are usually referred to as "carriers." Carriers are common in institutions for the mentally handicapped, particularly among adults committed since childhood, but they also exist in the normal population. For instance, the prevalence of HBsAg among blood donors in Britain is about 1/1000. Higher rates (up to 1%) have been recorded in commercial blood donors in the USA, reflecting the high proportion of drug addicts and the low socioeconomic status of such donor populations, while rates of up to 6% have been recorded in parts of Africa.

HBsAg-positive chronic liver disease

Detailed investigations of HBsAg-positive blood donors have shown evidence of liver disease in some, particularly when many drug addicts have been included among the donors investigated. In a study in north-west England,³ 40 out of 110 HBsAg-positive blood donors (36%), none of whom were parenteral drug users, showed abnormalities of liver function, particularly raised aminotransferase levels. In only six instances, however, did liver biopsy show a serious condition such as chronic aggressive hepatitis or cirrhosis, although further follow-up showed progression from persistent hepatitis to chronic aggressive hepatitis in two more. With a longer period of follow-up further evidence of progression may be found, but it is worth noting that the "healthy" carrier differs from the patient with HBsAg-positive chronic liver disease in several respects.

Firstly, there are notable differences in the distribution of both HBsAg and HBcAg in the liver. Intranuclear HBcAg may be found in many hepatocytes from patients with chronic liver disease but in few if any hepatocytes from healthy carriers. HBsAg, which is found in the cytoplasm, is present in less than 5% of hepatocytes in chronic liver disease and is scattered in random fashion throughout each lobule. In healthy carriers up to half the cells may contain HBsAg, which tends to occur in clumps or large sheets of affected hepatocytes. This accords with our understanding of the basic immunological defect in those carriers in whom no cellular immunity to HBsAg is detectable. This failure of immunity presumably allows the virus to exist in harmless symbiosis with its host and direct cell-to-cell spread of virus may consequently occur. In HBsAg-positive chronic active hepatitis, on the other hand, cellular immunity to HBsAg is present. Infected hepatocytes with viral determinants on their surface are consequently destroyed, which accounts for the finding of comparatively few HBsAg-containing cells. Indeed, in one study on eight patients with this condition, HBsAg was never found on the liver cell surface but only in the cytoplasm, where it is presumably immune from attack. Similarly, the many infected cells that contain core antigen in their nuclei, but which have not yet synthesised the coat protein, cannot be destroyed.

Liver Unit, King's College Hospital and Medical School, London SE5 8RX

I L WOOLF, BM, MRCP, lecturer
ROGER WILLIAMS, MD, FRCP, director and consultant physician

The absence of core particles from the nuclei of hepatocytes from healthy carriers is mirrored by an absence of Dane particles from the serum when examined with the electron microscope. Conversely, Dane particles are usually present in the serum of patients with HBsAg-positive chronic liver disease.

e Antigen—another antigen-antibody system in HBsAg-positive sera, but serologically distinct from HBsAg, was described in 1972⁴ and designated "e." This e antigen-antibody system has provided another difference between the healthy carrier and the patient with HBsAg-positive chronic liver disease. e Antigen is not found in healthy carriers but is often present in patients with chronic liver disease, particularly those with chronic aggressive hepatitis. Some correlation between the detection of e antigen and the presence of Dane particles also exists. e Antigen may be associated with DNA polymerase, which is detectable in sera containing Dane particles.

Dangers from HBsAg carriers

From a public health point of view, the potential infectivity of the healthy carrier is of great importance. The HBsAg carrier has been described as "the leper of the 20th century," which, since his detection usually results from an act of altruism—namely, donating blood—is a cruel irony. Is this description justified? Clearly HBsAg-positive blood cannot be accepted for transfusion, but should further restrictions be placed on the activities of a carrier? It is certainly advisable to exclude such people from working in haemodialysis and renal transplantation units. Both patients and staff in these units should be tested regularly for HBsAg and those found to be positive treated in isolation. The institution of such a policy in Britain has led to the virtual elimination of type B hepatitis from these units. In other parts of the world the infection in renal units continues to be rife.

Nevertheless, despite the accumulating evidence for the non-parenteral transmission of hepatitis B virus as a major mode of spread, the possibility that healthy carriers may transmit infection remains debatable. Serological studies of the families and household contacts of patients with HBsAg positive chronic liver disease have indicated that transmission of infection often occurs through intimate domestic and sexual contact. By contrast, in the families and household contacts of healthy carriers evidence of exposure to hepatitis B virus is rarely found. In a retrospective investigation in Sweden only two out of 16 long-term HBsAg carriers were implicated in cases of transfusion hepatitis, although all of them had previously donated much blood before their HBsAg state was discovered.⁵ Furthermore, at the time of the investigation transfusion hepatitis was usually associated with the ay subtype of HBsAg, whereas most of the long-term carriers were of subtype ad, which was the predominant subtype of HBsAg associated with type B hepatitis in Sweden 20 years previously. Similarly, although transmission of hepatitis B from an HBsAg-positive mother to her newborn infant is a recognised hazard, it is much more likely to occur to the infant of a woman contracting type B hepatitis in the last trimester than to the child of a healthy carrier. In the Far East, where HBsAg is much more prevalent, transmission from asymptomatic carrier mothers to their infants is not uncommon, but when it does occur e antigen has been detected in the maternal serum, suggesting that these women are not truly healthy carriers.

There is little evidence that persistent HBsAg carriers concerned in medical care transmit hepatitis B. Alter *et al*⁶ followed up 171 patients who were contacts of three healthy HBsAg carriers—two doctors and one nurse—for six to nine months. None of these patients developed type B hepatitis. A recent survey of dental surgeons at King's College Hospital, prompted by reports that the prevalence of HBsAg carriage in dentists was higher than in the general population, showed no evidence of transmission of hepatitis from an HBsAg-positive dentist to his patients. Although one such case has been reported, the dentist

concerned was suffering from chronic liver disease. Thus apart from the special circumstances of renal dialysis units and blood transfusion laboratories, where the risk of hepatitis is great, there appears to be no justification for restricting the activities of a healthy HBsAg-positive health care worker unless he has been definitely implicated in an outbreak of hepatitis.

The detailed investigation of the apparently healthy carrier is important in two respects. Primarily it will permit the identification of occult chronic liver disease, which may require future treatment, at an early stage. Secondly, in most instances where no such abnormality is found, these people may be reassured that they are neither harbouring a serious illness nor a great danger to the health of others.

Hepatitis B vaccine

Finally, mention should be made of efforts to produce a hepatitis B vaccine. Since HBsAb is a protective antibody, vaccines containing HBsAg in a non-infectious form may prove valuable. Maupas *et al* have reported the results of a vaccination programme using a vaccine prepared from the sera of patients with persistent HBs antigenaemia that were free of Dane particles and negative for e antigen.⁷ As an added precaution the vaccine was inactivated with 0.1% formalin. In 35 out of 46 people vaccinated an antibody response was induced. No adverse reactions occurred. Although these results are encouraging, their preliminary nature must be emphasised. In particular, the difficulty of ensuring complete inactivation of the virus and the possibility that host components may be present in the vaccine inducing harmful immunological reactions have been pointed out. Nevertheless, the way is clear for further work in this important area and, far from being a pariah, the healthy HBsAg carrier, through being the source of a hepatitis B vaccine, may help to eradicate one of the most serious infectious diseases remaining in the world.

References

- ¹ Nielson, J O, *et al*, *New England Journal of Medicine*, 1971, **285**, 1157.
- ² Reed, W D, *et al*, *Lancet*, 1973, **2**, 690.
- ³ Woolf, I L, Symposium on the Carrier State, Aviemore, 1974, British Society of Gastroenterology.
- ⁴ Magnus, L O, and Espmark, J A, *Journal of Immunology*, 1972, **109**, 1017.
- ⁵ Iwarson, S, *et al*, *British Medical Journal*, 1973, **1**, 84.
- ⁶ Alter, H J, *et al*, *New England Journal of Medicine*, 1975, **292**, 455.
- ⁷ Maupas, P, *et al*, *Lancet*, 1976, **1**, 1367.

What treatment is advised for a 34-year-old woman with marked premenstrual tension in whom the following has been serially tried without success: (a) fluphenazine and bendroflumethiazide, (b) frusemide, and (c) norethisterone daily for the last 10 days of the cycle?

The psychological, physiological, and endocrine basis of premenstrual tension is far from clear. If there is no pelvic abnormality the psychosocial history needs exploration and appropriate advice should be given. Fluid retention varies from patient to patient, and in some seems of little importance. The place of sex steroids in aetiology also varies. Progestogens have been the most popular treatment, not always with success, as in this case. Some authorities have suggested that either ethisterone or dydrogesterone may be more helpful than norethisterone. A combined oral contraceptive pill is worth a try. Possibly the fluctuation in the output of hormones is responsible for triggering the symptoms. The pill prevents these changes through the cycle and may allay worries about becoming pregnant. Less likely to be helpful than counselling and hormones are non-specific treatments such as alterations in diet, vitamins, and various tranquillisers, all of which have at some time been recommended, but probably because nothing else could be thought of.