CORRESPONDENCE

The end of excellence? J W Todd, FRCP.....1130	Professional standards M C T Morrison, FRCS.....1133	Not cricket J P Sparks, MD.....1135
DNCB, chemical laboratory workers, and chloramphenicol R J Pye, MRCP, and J L Burton, MD.....1130	Traumatic rupture of ear drum D McCracken, FRCP.....1133	The hospitals we need P R J Vickers, FRCS.....1135
Reactions to current influenza vaccine T W Hoskins, MB.....1131	Direct translation H-D Pache, MD.....1134	Closure of cottage hospitals D M Baker, MD.....1136
Side effects of prazosin M J Bendall, MRCP; C Rosendorff, MRCP.....1131	Oculo-mucocutaneous reactions to β-adrenoceptor antagonists J C Petrie, MRCP.....1134	Unpopular specialties F M Lanigan-O'Keeffe, MD.....1136
Chemotherapy in breast cancer J Wilkinson.....1132	Misuse of tubular elasticated bandages J B Bourke, FRCS.....1134	JHDA and HJSC D F H Guèret Wardle, MRCP, and others.....1136
Compulsory seat belts M J W Barker, MA.....1132	In vitro lymphocyte reactivity during pregnancy Y W Loke, MD, and others.....1134	Loss of earnings through committee attendances G S Duncan, MRCP.....1137
Neonatal electrocardiogram H Watson, FRCP; D P Southall, MRCP.....1132	Intravenous and aerosol salbutamol S P Deacon, MB.....1134	Health Service charges V Cooper, MB.....1137
"Press that bruise" P J Kirk, MB.....1132	Can hepatitis B be transmitted sexually? N R Grist, FRCPED.....1134	Hospital practitioner grade W D Waller, MB, DO.....1137
Hyposensitisation with house dust mite vaccine in bronchial asthma J Gaddie, MD, and others.....1132	Propranolol absorption in Crohn's disease and coeliac disease B T Cooper, MRCP, and others.....1135	National Insurance deductions J W Jackson, FRCS.....1137
Risk of venepuncture D Kohn, MD, and others.....1133	Sedatives and driving J M Plumb, MRCP.....1135	Pay differentials A A McInnes, MRCP.....1137
Thyrotoxic vomiting P Skrabanek, LAH; B I Chazan, FRCP.....1133	Detection of paraquat in urine B Widdop, PHD.....1135	BMA and HCSA Margaret C Watkinson, MB.....1137
Coronary artery disease and coalworkers' pneumoconiosis G S Andrews, FRCPATH.....1133	White coat D J Vicary, MRCP.....1135	Suspended consultant A J G Davis, MB.....1137
		Royal Medical Benevolent Fund Christmas appeal Sir Geoffrey Bateman, FRCS.....1137

Correspondents are urged to write briefly so that readers may be offered as wide a selection of letters as possible. So many are being received that the omission of some is inevitable. Letters should be signed personally by all their authors.

The end of excellence?

SIR,—Your leading article (2 October, p 779) begins: "Throughout the world there are a few hospital complexes where the quality of care given to patients, the teaching, and the academic research reach the highest possible standards." You also refer to "excellence," which requires a "concentration of medical talent and technological hardware", and comment that the centres of excellence improve standards and advance medicine far beyond their boundaries.

What are the criteria which determine that an institution is excellent? And how can we judge quality of care and standards of management? A few years ago in the most famous hospitals patients with small cancers of the breast were having radical mastectomies, patients with ankylosing spondylitis were immobilised for months on plaster beds, and patients with low backache were having their round ligaments shortened so as to anteverte their retroverted uteruses. Patients with uncomplicated duodenal ulcer were kept in bed for weeks being fed initially on citrated milk every two hours and later with the gradual addition of gruel, minced steamed fish, lightly poached eggs, rusks (to be well chewed), etc, until they were finally advised for ever after to avoid roast and fried dishes, salads, spices and seasoning, and most of the other features which make food attractive. Patients with cardiac infarcts were kept at absolute rest for a week or two and strictly in bed for six weeks and thereafter advised to live semi-invalid lives.

Patients with diverticulitis were given residue-free diets, and patients who were poisoned were given analeptic drugs and the poison was flushed out of them by forced diuresis. Were the hospitals where all this took place excellent? And were these patients being given care of the highest possible standard?

A commonly seen patient in the outpatient department is the middle-aged woman with multiple symptoms such as unpleasant head sensations, persistent low backache, being always tired and run down, widespread "rheumatism," and "indigestion." If she is seen by a physician who believes it is not his business to delve into his patients' emotions, concentrates his attention on her bodily symptoms, and admits her for a vast series of investigations is she being treated by the highest standards? Indeed, a feature of modern medicine as practised in the most famous hospitals is the ever-increasing number of investigations, many of which are done as routine. Is this an example of excellence?

The great majority of patients are not helped by being treated in an institution filled with the latest technological hardware and staffed with a galaxy of talent. They merely need a doctor with nous and sympathetic understanding. And the most excellent place for most of them is their own homes. Even if there are grounds for their being in hospital the most important of these grounds, if they have a medical illness, is often social; they may be living alone or old, with the main need

of nursing care. Those with most surgical conditions do just as well in a basic hospital; their prime need is a competent surgeon who performs the right operation at the right time.

May I finally suggest that we should all be wary of using such words as "excellence" and "standards"? We constantly hear assertions about standards—how they are rising or, more often, falling—according to the particular axe that the writer is grinding. Such assertions are mostly quite meaningless.

JOHN W TODD

Farnham, Surrey

DNCB, chemical laboratory workers, and chloramphenicol

SIR,—2, 4-Dinitrochlorobenzene (DNCB) is a potent sensitizer which is widely used in medicine to test the integrity of the immune response in experimental animals or human patients. Most organic chemists appear to be unaware of its sensitising properties, and DNCB is a standard laboratory preparation which is used for the routine preparation of derivatives.

We recently treated two laboratory technicians who had both taken a practical organic chemistry examination which involved the preparation of DNCB. The first patient was a 23-year-old man who had not allowed the DNCB to come into direct contact with his skin, but he had spilt some of the prepared DNCB solution on to his examination paper, which he had then put into the back pocket of his trousers to discuss with his friends after the examination. Some 24 h later he developed a widespread urticated eczematous eruption which was maximal over the right buttock and which also involved the chest, abdomen, arms, and hands. The rash subsided rapidly with prednisone 50 mg daily

for one week but flared up again on stopping treatment. His skin eventually returned to normal with the topical application of a fluorinated steroid for several weeks. The second patient was a 21-year-old woman who developed a severe eczema of the hands and neck which started on the day after her chemistry examination. She was sure that no caustics had spilt on her skin but thought that the DNCB solution might have come into contact with her hands. She was treated with potassium permanganate soaks, a steroid cream, and prednisone 20 mg daily for 4 days, but despite this the condition became more severe. It eventually responded to prednisone 50 mg daily for one week, which was then rapidly withdrawn, and her skin has since been normal. In view of the severity of the reaction produced we felt it inadvisable to perform patch tests on either patient.

Both patients worked in chemistry laboratories and believed they might have handled DNCB in the past and neither had been warned of its sensitising potential. An inquiry at the institution which organised the examination revealed that none of the organic chemists on the staff knew that DNCB is a sensitiser and the fact is not mentioned in the standard practical textbooks of organic chemistry.

We feel it unlikely that a dermatologist would overlook the possibility of DNCB sensitivity in a chemist or laboratory technician presenting with a contact dermatitis, but it may not be so well known that DNCB cross-reacts with chloramphenicol.¹ The preparation of DNCB on one or more occasions could presumably therefore sensitise a person to chloramphenicol, with the subsequent provocation of a severe eczematous reaction following systemic or topical exposure to the drug. In view of the increasing use of topical chloramphenicol we feel that the potential hazards of handling DNCB should be widely publicised to doctors and chemists.

R J PYE
J L BURTON

Department of Dermatology,
Bristol Royal Infirmary,
Bristol

¹ Schwank, R, and Jirasek, L, *Hautarzt*, 1963, 14, 24.

Reactions to current influenza vaccine

SIR,—For some years the Medical Officers of Schools Association has supported the belief that boarding-school children should be among those offered annual influenza vaccination. This has been supported by evidence of its efficacy.^{1 2} One of the assumptions behind this policy is that the rate of systemic reactions to influenza vaccine is low. For some years the rate has been below 1%.

This year there have been reports of high rates of reaction to the vaccine, which contains influenza A/New Jersey/76 for the first time. From my contacts with members of this association I have obtained data from 14 schools which have given one of the two commercially available (and antigenically identical) vaccines to over 4600 pupils. The overall rate of systemic reactions is 15%. This is clearly a crude figure as rates of reporting and criteria of diagnosing reactions must vary. Nevertheless, the picture differs so markedly from previous years that I feel it is worth bringing to your attention.

The reactions described include pyrexia, reaching 39.4°C in some cases, headache, nausea, vomiting, faintness, and myalgia. In most schools they have been more frequent in the younger pupils, even when the under-13s

have been given half the adult dose. None of the reactions lasted longer than 36 h and no sequelae have been reported to me.

This unexpectedly high reaction rate must lead to a reassessment of vaccination policy, including reconsideration of the intradermal route recommended by Payler and Skirrow.³

T W HOSKINS
Honorary Secretary,
Medical Officers of Schools Association

The Infirmary,
Christ's Hospital,
Horsham, Sussex

¹ Hoskins, T W, et al, *Lancet*, 1973, 2, 116.

² Hoskins, T W, et al, *Lancet*, 1976, 1, 105.

³ Payler, D K, and Skirrow, M B, *British Medical Journal*, 1974, 2, 727.

Side effects of prazocin

SIR,—The recent short report by Professor Clive Rosendorff (28 August, p 508) indicates that, at least symptomatically, commencement of prazosin therapy for hypertension with a 0.5-mg dose instead of the previously recommended 2 mg may be accompanied by a lower incidence of side effects a short time after taking the first dose. However, there are several points in the paper which I feel are worthy of comment and clarification.

Though the author states that supine and standing blood pressures were recorded before and at intervals after administration of prazosin, the only results given are for standing pressures. Hypotension as a result of antihypertensive therapy is often a relative phenomenon. It may manifest itself with change in posture from the lying to the standing position. Also in previously hypertensive patients, although lowering of the blood pressure may not result in typically hypotensive levels, a degree of lowering may be achieved that for that patient may produce symptoms of hypotension—this depending on the state of the cerebral vasculature and the rate at which the blood pressure is lowered. Though the dose of drug used was small, the depression of systolic blood pressure in one patient (case 12) was dramatic at $\frac{1}{2}$ h (62 mm Hg). This implies that for certain patients the drug is still very potent even in small dosage. In all, four of the seven patients who were rechallenged had standing systolic pressures lowered by more than 30 mm Hg within $1\frac{1}{2}$ h of administration and in three of these four the fall was in excess of 45 mm Hg, again attesting to the potency of the drug in some patients. Consequently in the context of this paper it is difficult to assess the meaning of the statement, "The standing blood pressure fell in all cases, but in no patient was this fall to anything that could be described as a hypotensive level."

In spite of the low incidence of side effects (only two patients had transient mild dizziness) only three of the seven patients were continued on prazosin for 1-12 weeks (whatever that means). If these seven patients were hypertensives in need of treatment by the criteria used in the clinic, why was prazosin not continued in more of them after a small starting dose had apparently been well tolerated? And which of the seven patients were continued and for how long?

Why, of the twelve patients who had first-dose reactions, did only seven volunteer for rechallenge with a smaller dose of the drug? One must assume that either the side effects were so severe on the first exposure that the medical staff felt it unwise to request further

exposure or that the symptoms were of such severity that five patients felt unable to give further consent for taking part. By either criterion these patients must be regarded as failures and this consequently makes nonsense of the predicted incidence of first-dose reactions in the original population of 74 patients as being 4.6%. In fact the incidence taking into account these five patients would be at least 6.9%, and taking into account the two patients who had reactions on the smaller dose and those who were not continued after the challenge with a smaller dose the incidence of the first reaction may be as high as 10%. This is clearly unacceptable.

I fear that this paper does little to allay the concern regarding prazosin. Although only two patients experienced mild dizziness, it is unwise to extrapolate the findings from this small, randomly selected group of patients being studied in a controlled clinical situation for a relatively short period of time to that pertaining to the usual circumstances of commencement of antihypertensive therapy—that is, outpatients leading normal active lives. Its routine use in the treatment of mild and moderate hypertension is far from established when there are other preparations with less, or potentially less, catastrophic side effects which have been used for a number of years and are trusted, in so far as one can trust any drug, by a large number of physicians. Its general acceptance must await adequate clinical trials with a 0.5-mg starting dose in which it is compared with other, established, antihypertensive agents.

I also feel that the decision to rechallenge these patients with a drug known to have severe initial side effects and in whom the drug has already produced such an initial reaction, is ethically unacceptable.

M J BENDALL

Department of Geriatric Medicine,
St Mary's Hospital,
Colchester

* * * We sent a copy of this letter to Professor Rosendorff, whose reply is printed below.—
Ed, *BMJ*.

SIR,—I am grateful to Dr Bendall for raising some points which were impossible to cover in the "short report" format.

An excessive fall in blood pressure on changing from the supine to the standing position obviously may cause dizziness. For the seven patients in the study the mean pretreatment postural fall in blood pressure was $6/2 \pm 7/4$ (SEM) mm Hg. Corresponding values for the period after 0.5 mg prazosin were: $\frac{1}{2}$ h, $12/4 \pm 9/2$ mm Hg; 1 h, $2/2 \pm 10/4$ mm Hg; and $1\frac{1}{2}$ h, $12/2 \pm 4/4$ mm Hg. The pre- and post-treatment values were not significantly different, nor were the post-treatment values significantly different from zero, on paired *t*-testing. In other words we were unable to demonstrate a consistent or significant postural fall in blood pressure in the study. This is not to say that in individual cases there was not a large postural drop; the two patients who did show some dizziness had quite large postural falls of blood pressure. In one case the systolic blood pressure fell from 194 mm Hg supine to 160 mm Hg erect and in another from 168 mm Hg supine to 128 mm Hg erect. This underlines the need for caution in the use of this drug; there are patients who are sensitive to its effects and some patients are more sensitive than others. Prazosin is a potent antihypertensive (isn't that what we are after?),