Local adaptations in populations of rock-dwelling haplochromines (Pisces: Cichlidae) from southern Lake Victoria

N. Bouton, F. Witte, J. J. M. van Alphen, A. Schenk and O. Seehausen

Institute of Evolutionary and Ecological Sciences, Leiden University, PO Box 9516, 2300 RA Leiden, The Netherlands
(bouton@rulsfb.leidenuniv.nl)

The radiation of cichlid fishes in the African great lakes is often described as adaptive, because, at a superficial level, cichlid fishes seem adapted to the ecological niches they occupy. However, adaptiveness has rarely been studied. We investigated to what extent island populations of three species of the rock-dwelling genus *Neochromis*, endemic to Lake Victoria, are adapted anatomically to exploit locally abundant resources. Specifically, we asked whether different resource environments were reflected in differences in the feeding apparatus, both within species and between species. In populations of two specialized biters, the algae scrapers *N. rufocaudalis* and *N. omnicaeruleus*, the biting force of the lower jaw increased with increasing amount of items that require biting in the diet. *N. greenwoodi* is a less specialized biter; we found differences between two populations in the hyoid position and in the premaxilla that enhance suction feeding. These adaptations were related to the amount of items requiring suction. Comparing across three sympatric pairs of species, in each case different diets were reflected in differences in anatomy.

Keywords: adaptation; cichlid; diet; Lake Victoria; morphology; phenotypic plasticity

1. INTRODUCTION

The question of why there are so many species of cichlid fishes in the African great lakes has been debated for several decades. Discussion revolves around two ideas: (i) cichlid speciation is promoted strongly by sexual selection (Dominey 1984; Turner & Burrows 1995; Seehausen *et al.* 1997); and (ii) cichlid 'design' provides a unique substrate for rapid anatomical change (Liem 1973; Greenwood 1991; Galis & Metz 1998). These ideas are not conflicting, but are complementary (Galis & Metz 1998). Sexual selection is a mechanism that may produce reproductively isolated forms (or species), but cannot account for the adaptive radiation of cichlids which permits them to use virtually all resources in the great lakes. Highly adapted anatomical features and body shapes can be achieved through relatively gradual modifications, involving slight heterochronic shifts in the ontogeny (Strauss 1984; Meyer 1987; Greenwood 1991).

Philopatry, small brood sizes, and parental care, leading to low dispersal, may create the possibility of intralacustrine, allopatric speciation. Rock-dwelling cichlids, which are often reluctant to cross stretches of open water or other substrates are good candidates for such processes (McKaye & Gray 1984). Fine-scale genetic differentiation has been shown for many of these species (Dorit 1990; Verheyen *et al.* 1996; Bowers *et al.* 1994; van Oppen *et al.* 1997), although in some species, which might be better dispersers, no such a differentiation was found (Meyer *et al.* 1996). However, fine-scale genetic differentiation in populations of rock-dwelling species in itself

does not necessarily lead to allopatric speciation, because selection pressures may be similar across the range of a species (McPeck (1996) and references therein). Conversely, if populations in different parts of a species' range experience different selection regimes, populations in these different areas may evolve to the point of differentiation that can lead to speciation (McPeck (1996) and references therein). The aim of this paper is to trace adaptations in natural populations of cichlid fishes. An adaptation is a trait that enhances fitness and that arose historically as a result of natural selection for its current biological role (Lauder 1996). We analyse adaptations in the prey-capture apparatus using models from functional morphology to predict how shapes of structures should change to improve exploitation of certain resources. Adaptiveness of traits has, to our knowledge, never been tested in natural populations of cichlids, but the functional anatomy of the feeding apparatus has been well described (Liem 1980, 1991; Otten 1983; van Leeuwen & Spoor 1987; de Visser & Barel 1996, 1998). Feeding of cichlids can roughly be divided into biting and suction (Barel 1983; Liem 1991). A third category (ram feeding) distinguished by Liem (1991), is not found among the species described here. Biting is used when any forceful manipulation of the food by the oral jaws is needed. However, suction feeding is the most efficient way to collect elusive prey. Both biting and suction place conflicting demands on the anatomy (Barel 1983). Two models (Otten 1983; van Leeuwen & Spoor 1987) make predictions about efficient biting, the third model (de Visser & Barel 1996, 1998) concentrates on suction feeding.

The genus studied, *Neochromis*, is a lineage of rock-dwelling biters, feeding largely on epilithic algae, but the amount of algae in the diet is variable among species and among populations of the same species. The proportions of different food types vary among rocky islands. We hypothesized that if food intake varies with resource availability, adaptive differences in trophic traits will evolve.

2. MATERIALS AND METHODS

(a) Research area, sampling, species and populations

From August 1990 until August 1991 we sampled *Neochromis* at four rocky islands on a south–north transect in southern Lake Victoria: Python and Anchor Islands in the Mwanza Gulf, and Makobe and Ruti Islands in the Speke Gulf. The distances between these islands are: Python–Anchor, 8.2 km; Anchor–Makobe, 21.6 km; and Makobe–Ruti, 11.5 km. We sampled by angling close to, and in-between, the rock boulders around the islands, and by gill-netting in slightly deeper water (2–7 m), 530 m off the shoreline. Fishes were put on ice, fixed in 10% formalin neutralized with sodium tetraborate (Borax), and after several months of fixation transferred to 70% ethanol.

We studied seven populations of three species (table 1; Seehausen *et al.* 1998), viz. *Neochromis rufocaudalis* Seehausen and Bouton 1998 from Python, Anchor and Makobe (absent at Ruti), *Neochromis greenwoodi* Seehausen and Bouton 1998 from Python and Anchor (absent at Makobe and Ruti), and *Neochromis omnicaruleus* Seehausen and Bouton 1998 from Makobe and Ruti (absent at Python and Anchor). *N. greenwoodi* and *N. omnicaruleus* make up an allopatric sibling species pair. Both species have a colour morph (orange blotched) not found in other *Neochromis* species, and their distribution is complementary (Seehausen 1996).

(b) Diet and resource base

To determine diets we analysed stomach contents of 36–43 specimens per population (table 1), 8–12 specimens from four different seasons, namely March–May, June–August, September–November and December–February. In the case of *N. omnicaruleus* from Ruti Island, we analysed 24 specimens caught in one season (June–August), because the remote island could only be reached in that period. We disregarded specimens with empty stomachs or stomachs with very few items (filled at less than 40%). We calculated the volumetric proportion of attached organisms, i.e. filamentous algae, bryozoans and sponges, which are collected by biting techniques. The remainder of the food, the non-attached organisms, such as insect larvae, zooplankton, and diatoms, are collected by feeding techniques that involve suction.

Direct measurement of relative contributions of food items to the resource base is hampered by the infrastructure of the rocky islands, which are rich in rock holes and crevices. Therefore, we used an indirect approach. Fish densities are very high at rocky islands (11.6 m⁻² at Makobe, Seehausen & Bouton (1997)). Thus, food is probably limiting densities. Since the up to 30 species of haplochromine cichlids at each island exploit virtually all resources, relative biomass of species exploiting certain resources will reflect the relative contribution of these resources to the resource base. We used the proportion of fishes that fed on attached organisms as an estimation of the proportion of attached organisms in the resource base (table 1). The method described above is only an approximation, but qualitative obser-

Table 1. *Test of the hypothesis that the proportion of attached organisms in the diet increases with the proportion available in the resource base*

(Islands are placed in order of increasing proportion of attached organisms in the resource base; we therefore expect a similar order in the proportion of attached organisms in the diet. The body of the table gives mean proportions of attached organisms in the diet and numbers of specimens used. Bracketed values indicate specimen number from which they were selected. The *p*-values are based on the Jonckheere (J) or the Wilcoxon–Mann–Whitney test (W).)

	Python	Anchor	Ruti	Makobe	
attached organisms in the resource base	0.173	0.418	0.523	0.648	
species	attached organisms in the diet				<i>p</i> -value
<i>N. rufocaudalis</i>	0.529	0.656	—	0.8	<0.001 (J)
number	24 (36)	25 (40)	—	19 (43)	
<i>N. greenwoodi</i>	0.215	0.413	—	—	0.017 (W)
number	33 (40)	23 (40)	—	—	
<i>N. omnicaruleus</i>	—	—	0.543	0.640	0.104 (W)
number	—	—	14 (24)	30 (43)	

vations during catches support the results (see table 1 in Bouton *et al.* (1997)). We used non-parametric tests: consequently, only rank data have been used. To test the hypothesis that the proportion of attached organisms in the diet increases with the proportion available in the resource base, we used Jonckheere's test for the case of three populations of one species, and the one-tailed Wilcoxon–Mann–Whitney test for two populations (see Siegel & Castellan 1988).

(c) Anatomical models

Five male individuals of about 90 mm standard length were used from each population. Linear measurements were divided by a measure of neurocranium length for standardization. We did an anatomical analysis of the prey-capture apparatus, which consists of the oral jaws and their musculature and the suspensory apparatus. Anatomical biting force F_T of the lower jaw is calculated as (van Leeuwen & Spoor 1987)

$$F_T = F_M * \frac{(R \text{ to } Z)}{(R \text{ to } T)} = \frac{V_m}{(C \text{ to } D)} * \frac{(R \text{ to } Z)}{(R \text{ to } T)} = V_m * \frac{(R \text{ to } Z)}{(C \text{ to } D)} * \frac{1}{(R \text{ to } T)}$$

where F_M is the power that can be exerted by pars 2 of the *musculus adductor mandibulae* (figure 1a), calculated as its dry weight (V_m) divided by fibre length (C to D, figure 1a), R to Z is the length of the arm from the centre of rotation (R, figure 1a) perpendicular to a muscle fibre to the point where it crosses this fibre (Z, figure 1a), R to T is the length of the lower jaw from centre of rotation (R) to the tip of the most exterior tooth (T, figure 1a). The ratio of arm (R to Z) and fibre length (C to D) is approximately constant throughout the muscle. We segmented the muscle surface in six parts and averaged the ratio over these segments.

For the upper jaw or premaxilla, trade-offs between suction feeding and biting are straightforward. Protrusion of the premaxilla, of importance for suction feeding, will be enhanced if the ascending arm of the premaxilla is long and the angle β between the ascending arm and the dentigerous arm is small

Figure 1. (a) The biting force model (after Van Leeuwen & Spoor 1987). The shaded area is pars 2 of the *musculus adductor mandibulae*; F_T , biting force of the lower jaw; R to T, lower jaw length; C to D, muscle fibre length; R to Z, length of arm perpendicular to muscle fibre. (b) The premaxilla, β is the angle between ascending arm and dentigerous arm, A to B is the length of the ascending arm.

(figure 1b). In contrast, transmission of biting force is optimized by shortening the ascending arm and increasing β (Otten 1983).

Head expansion during suction feeding in fishes involves three main movements: (i) mouth opening; (ii) outward rotation of the lateral walls (i.e. suspensorium and gill cover); and (iii) depression of the bottom of the mouth. Evidently, mouth opening should start prior to the other two movements. From their kinematic model of the expansion apparatus in fishes, De Visser & Barel (1996) derived a quantitative measure for the degree of efficiency by which the apparatus of a specimen was able to fulfil this demand. A simplified representation of this measure and its kinematic background is given here (figure 2). The three-dimensional model concerns two doors, representing left and right suspensorium, each suspended by a hinge joint from a ceiling (the ventral face of the neurocranium). The hyoid bones articulate with the inner side of each door. The other ends of the hyoids articulate with each other in the medial plane. On this symphysis operates the expansion force (F_E), which via complex anatomical chains with the hyoid as a central element causes all expansion movements. For instance, the expansion force will cause an outward rotation of the suspensorium (figure 2). The model allows two types of movements: (i) rotation of the two hyoids together (the hyoid chevron) around a hypothetical axis through its left and right articulation with the suspensorium and (ii) widening of the angle between the two hyoids. Hyoid chevron rotation and angle widening are

Figure 2. (a) Schematic representation of the kinematic model of the expansion apparatus (after De Visser & Barel 1996, 1998). Note that the expansion force F_E and σ rotation are in the medial plane, while α rotation is in the plane of the two hyoids. (b) α , σ curve, $d\alpha$ is the distance to the curve, which gives optimality of the hyoid position for all combinations of α and σ .

measured, respectively, by the angular parameters σ and α , the anatomical definitions of which are not needed here. Note that changing α necessarily involves suspensorial rotation. Otherwise stated: for every α there is a corresponding position of the suspensorium. From the foregoing it may be understood, that among all α , σ positions feasible there will be several for which the hyoids are positioned in the plane of the (model's) suspensorium. In these positions the expansion force will cause no outward rotation of the suspensorium and, ideally, at the onset of suction feeding the hyoid should be in such a position (de Visser & Barel 1998). However, anatomical constraints may preclude the hyoid from obtaining this optimal starting position. As an estimate of the degree of deviation from this optimality, the distance of the real anatomical hyoid position to the curve of optimal positions ($d\alpha$) could be measured (see De Visser & Barel (1996, 1998) for details).

Constructional constraints restrict the possibility of optimizing both suction feeding and biting at the same time, e.g. F_T and $d\alpha$. Biting force (F_T) may be improved by an increase of the cross-section of the *m. adductor mandibulae*. To accommodate this thicker muscle the head has to widen, which implies a lateral displacement of the touching point between the hyoid and the suspensorium (figure 2). This displacement increases the angle between the hyoids, which in turn affects the optimality of the starting position of the hyoid ($d\alpha$) negatively (de Visser & Barel 1996).

(d) *Inter- and intraspecific morphological comparisons*

We did three pairwise interspecific comparisons of sympatric populations: (i) *N. rufocaudalis* and *N. greenwoodi* from Python Island; (ii) *N. rufocaudalis* and *N. greenwoodi* from Anchor Island;

Figure 3. Anatomical comparisons. Populations are placed in order of increasing proportion of attached organisms in the diet. Indicated are ranges (mean $\pm$ s.d.) for the populations. (a) Biting force of the lower jaw. (b) Premaxilla: β and ascending arm length (scaled downward). (c) hyoid position $d\alpha$. Interspecific comparisons of sympatric populations are made by comparing columns with the same shading pattern. Mean values of populations of the same species from different locations are connected with lines for ease and significant differences ($p < 0.05$) are denoted (*).

and (iii) *N. rufocaudalis* and *N. omnicaeruleus* from Makobe Island. For intraspecific comparisons we used three populations of *N. rufocaudalis*, and two of both *N. greenwoodi* and *N. omnicaeruleus* (table 1). To test whether there were differences in the predicted direction we used one-tailed analyses of variance (ANOVAs), performed in Statgraphics. We did a Bonferroni procedure, because all groups are used in two comparisons.

3. RESULTS

(a) Resource availability and diet

In *N. rufocaudalis* and *N. greenwoodi* the proportion of attached organisms in the diet increases significantly with the estimated proportion in the resource base (table 1). It is likely that the same holds for *N. omnicaeruleus*, but the sample of Ruti Island is unfortunately rather small (table 1). For all populations of *N. rufocaudalis*, proportions of attached organisms in the diet were higher than in the local resource base, while for populations of *N. greenwoodi* and *N. omnicaeruleus* proportions in the diet and the

resource base were roughly the same (table 1). Among sympatric populations, *N. rufocaudalis* always had the highest proportion of attached organisms in its diet.

(b) Anatomical comparisons of sympatric species

Of the three species pairs we predict that *N. rufocaudalis* will be better adapted to biting and *N. greenwoodi* or *N. omnicaeruleus* to suction. The species pair that differed most in diet, i.e. *N. greenwoodi* and *N. rufocaudalis* from Python Island, differed in three anatomical aspects in the predicted direction (figure 3): (i) the hyoid position ($d\alpha$) of *N. greenwoodi* is better suited for suction ($p < 0.1$); (ii) the ascending arm of the premaxilla is longer ($p < 0.01$); (iii) the angle β is smaller in *N. greenwoodi* than in *N. rufocaudalis* ($p < 0.01$). The latter two aspects of the premaxilla are better suited for suction in *N. greenwoodi* and for biting in *N. rufocaudalis*. According to predictions anatomical biting force of the lower jaw is larger in *N. rufocaudalis* than in *N. greenwoodi* from Anchor Island ($p < 0.1$). *N. omnicaeruleus* from Makobe island is better adapted to suction feeding than *N. rufocaudalis* from the same island, since $d\alpha$ is smaller ($p < 0.1$) and the ascending arm of the premaxilla is longer ($p < 0.05$, figure 3), which is also in accordance with predictions.

(c) Intraspecific anatomical comparisons

For all three species we predict that populations adapt to either biting or suction with respectively increasing or decreasing amounts of attached organisms in the diet. The two populations of *N. greenwoodi* differed in the predicted direction in three out of four aspects of the anatomy: hyoid position $d\alpha$ ($p < 0.05$) and the two aspects of the premaxilla, the angle β ($p < 0.05$) and the length of the ascending arm ($p < 0.05$, figure 3). In the two populations of *N. omnicaeruleus* we found an increase in anatomical biting force of the lower jaw in concordance with increasing proportion of attached organisms in the diet ($p < 0.05$, figure 3). In the three populations of *N. rufocaudalis* there is an increasing trend of anatomical biting force of the lower jaw in concordance with diet (figure 3).

4. DISCUSSION

(a) Trophic diversification

Neochromis species change the amount of attached organisms in their diet with the estimated availability in the environment. This may indicate that resource availability in the environment provides a stronger selection force than other forces within the community, such as interspecific interactions. The major part of the diet of *N. rufocaudalis* and *N. omnicaeruleus* always consisted of attached organisms. We found trends indicating that biting force of the lower jaw increased with the proportion of attached organisms in the diet, while no such trends were found in the other aspects of the anatomy studied in these species. The major part of the diet of *N. greenwoodi* consisted of non-attached organisms; this was more extreme in the population from Python than in the population from Anchor Island. We found adaptations towards suction feeding in the population from Python Island, while no differences between populations were found in anatomical biting force of the lower jaw. Thus, in

all three species the dominant way of feeding, biting in *N. rufocaudalis* and *N. omnicaeruleus* and suction feeding in *N. greenwoodi*, seems to induce changes in the feeding apparatus. We did not find trade-offs between specialization for suction (hyoid position $d\alpha$) and biting force of the lower jaw. Probably changes are too small to find such an effect. For instance, the range of hyoid positions ($d\alpha$) is about 10–16 among the three *Neochromis* species, while for all Lake Victoria species this range is about 0–18 (de Visser & Barel 1996).

(b) Genetic divergence versus phenotypic plasticity

Two processes may cause anatomical differences between populations of rock-dwelling haplochromines, namely (i) genetic divergence due to local adaptation or genetic drift and (ii) developmental plasticity. Measuring heritabilities of anatomical properties is necessary to assess the relative roles of the two processes. However, there is some evidence that genetic divergence plays a role. Populations of *N. greenwoodi*, including the ones studied here, did show different mitochondrial haplotype frequencies (Dorit 1990), which indicates a restricted gene flow between populations and allows genetic divergence. *N. greenwoodi* from Python Island and Anchor Island, which were bred in tanks under similar conditions in our laboratory, did not show visible character convergence for two generations. Developmental plasticity of the feeding apparatus could be demonstrated experimentally in haplochromine cichlids including *N. rufocaudalis* and *N. greenwoodi*, when siblings were reared on different food types (Witte *et al.* 1990; N. Bouton, personal observation). Thus, adaptations of populations in the field probably result from genetic divergence, while developmental plasticity may have an augmentative effect. In conclusion, adaptive divergence, as described here, can lead to intra-lacustrine, allopatric speciation and may also play a role in adaptive radiation after sympatric and parapatric speciation by sexual selection.

Sarah Collins, Jaap Dubbeldam, Patsy Haccou, Les Kaufman, Peter Mayhew, Dolph Schluter and an anonymous referee gave useful comments on earlier drafts of the manuscript. Johan van Leeuwen and Kees Barel advised on biomechanical models. We thank Martin Brittijn for drawing the figures. The research was supported by WOTRO grant W84-282. We thank the Tanzanian Fisheries Research Institute (TAFIRI) for its hospitality, and Moja Kayeba and Reuben Enoka for invaluable field assistance.

REFERENCES

- Barel, C. D. N. 1983 Towards a constructional morphology of cichlid fishes (Teleostei, Perciformes). *Neth. J. Zool.* **33**, 357–424.
- Bouton, N., Seehausen, O. & van Alphen, J. J. M. 1997 Resource partitioning among rock-dwelling haplochromines (Pisces: Cichlidae) from Lake Victoria. *Ecol. Freshw. Fish.* **6**, 225–240.
- Bowers, N., Stauffer, J. R. & Kocher, T. D. 1994 Intra- and inter-specific mitochondrial DNA sequence variation within two species of rock-dwelling cichlids (Teleostei: Cichlidae) from Lake Malawi, Africa. *Mol. Phyl. Evol.* **3**, 75–82.
- de Visser, J. & Barel, C. D. N. 1996 Architectonic constraints on the hyoid's optimal starting position of suction feeding of fish. *J. Morph.* **228**, 1–18.
- de Visser, J. & Barel, C. D. N. 1998 The expansion apparatus in fish heads, a 3-D kinetic deduction. *Neth. J. Zool.* **48**, 361–395.
- Dominey, W. J. 1984 Effects of sexual selection and life history on speciation: species flocks in African cichlids and Hawaiian *Drosophila*. In *Evolution of fish species flocks* (ed. A. A. Echelle & I. Kornfield), pp. 231–249. University of Maine: Orono Press.
- Dorit, R. L. 1990 The correlates of high diversity in Lake Victoria haplochromine cichlids: a neontological perspective. In *Causes of evolution: a paleontological perspective* (ed. R. M. Ross & W. D. Allmon), pp. 322–353. University of Chicago Press.
- Galis, F. & Metz, J. A. J. 1998 Why are there so many cichlid species? *Trends Ecol. Evol.* **13**, 1–2.
- Greenwood, P. H. 1991 Speciation. In *Cichlid fishes: behaviour, ecology and evolution* (ed. M. H. A. Keenleyside), pp. 86–102. London: Chapman & Hall.
- Lauder, G. V. 1996 The argument from design. In *Adaptation* (ed. M. R. Rose & G. V. Lauder), pp. 55–91. San Diego: Academic Press.
- Liem, K. F. 1973 Evolutionary strategies and morphological innovations: cichlid pharyngeal jaws. *Syst. Zool.* **22**, 425–441.
- Liem, K. F. 1980 Adaptive significance of intra- and interspecific differences in the feeding repertoires of cichlid fishes. *Am. Zool.* **20**, 295–314.
- Liem, K. F. 1991 Functional morphology. In *Cichlid fishes: behaviour, ecology and evolution* (ed. M. H. A. Keenleyside), pp. 129–150. London: Chapman & Hall.
- McKaye, K. R. & Gray, W. N. 1984 Extrinsic barriers to gene flow in rock-dwelling cichlid species flocks in African Lakes. In *Evolution of fish species flocks* (ed. A. A. Echelle & I. Kornfield), pp. 169–183. University of Maine: Orono Press.
- McPeck, M. A. 1996 Linking local species interactions to rates of speciation in communities. *Ecology* **77**, 1355–1366.
- Meyer, A. 1987 Phenotypic plasticity and heterochrony in *Cichlasoma managuense* (Pisces, Cichlidae) and their implications for speciation in cichlid fishes. *Evolution* **41**, 1357–1369.
- Meyer, A., Knowles, L. L. & Verheyen, E. 1996 Widespread geographical distribution of mitochondrial haplotypes in rock-dwelling cichlid fishes from Lake Tanganyika. *Mol. Ecol.* **5**, 341–350.
- Otten, E. 1983 The jaw mechanism during growth of a generalized *Haplochromis* species: *H. elegans* Trewavas 1933 (Pisces: Cichlidae). *Neth. J. Zool.* **33**, 55–98.
- Seehausen, O. 1996 *Lake Victoria rock cichlids: taxonomy, ecology, and distribution*. Zevenhuizen, Netherlands: Verduyn Cichlids.
- Seehausen, O. & Bouton, N. 1997 Microdistribution and fluctuations in niche overlap in a rocky shore cichlid community in Lake Victoria. *Ecol. Freshw. Fish.* **6**, 161–173.
- Seehausen, O., van Alphen, J. J. M. & Witte, F. 1997 Cichlid fish diversity threatened by eutrophication that curbs sexual selection. *Science* **277**, 1808–1811.
- Seehausen, O., Lippitsch, E., Bouton, N. & Zwennes, H. 1998 Mbipi, the rock-dwelling cichlids of Lake Victoria: description of three new genera and 15 new species (Teleostei). *Ichth. Expl. Freshw.* **9**, 129–228.
- Siegel, S. & Castellan, N. J. Jr. 1988 *Nonparametric statistics for the behavioral sciences*, 2nd edn. New York: McGraw-Hill.
- Strauss, R. E. 1984 Allometry and functional feeding morphology in haplochromine cichlids. In *Evolution of fish species flocks* (ed. A. A. Echelle & I. Kornfield), pp. 217–229. University of Maine: Orono Press.
- Turner, G. F. & Burrows, M. T. 1995 A model of sympatric speciation by sexual selection. *Proc. R. Soc. Lond. B* **260**, 287–292.
- van Leeuwen, J. L. & Spoor, C. W. 1987 Biting force in cichlids: a discussion of the planar model of Barel (1983). *Neth. J. Zool.* **37**, 307–314.

- van Oppen, M. J. H., Turner, G. F., Rico, C., Deutsch, J. C., Ibrahim, K. M., Robinson, R. L. & Hewitt, G. M. 1997 Unusually fine-scale genetic structuring found in rapidly speciating Malawi cichlid fishes. *Proc. R. Soc. Lond. B* **264**, 1803–1812.
- Verheyen, E., Rüber, L. & Snoeks, J. 1996 Mitochondrial phylogeography of rock-dwelling cichlid fishes reveals evolutionary influence of historical lake level fluctuations of Lake Tanganyika, Africa. *Phil. Trans. R. Soc. Lond. B* **351**, 797–805.
- Witte, F., Barel, C. D. N., Hoogerhoud, R. J. C. 1990 Phenotypic plasticity of anatomical structures and its ecomorphological significance. *Neth. J. Zool.* **40**, 278–298.