imagining himself to be volumes and so years ahead of the rest of us is that you have erroneously imputed such notions to him; though as between the possibility of the editor of *World Medicine* having delusions of grandeur and that of the editor of the *British Medical Journal* ignoring convention I should prefer not to speculate. Perhaps next time, however, you will put your editorial heads together and so avoid supplying an occasion for their being banded together later.

JOHN S BRADSHAW

How Caple,
Hereford

¹ Fowler, H W, *A Dictionary of Modern English Usage*, 2nd edn, revised E Gowers. London, Oxford University Press, 1965.
² *British Medical Journal*, 1976, 1, 6.

Possible immunological effect of oxprenolol

SIR,—I am writing about a possible adverse reaction to oxprenolol (Trasicor).

A 40-year-old male patient of mine who has severe hypertension is taking oxprenolol and polythiazide. Two weeks after his son's attack of chickenpox he and his daughter developed chickenpox. While the children's illness was of average severity his, however, was of a quite extreme degree.

I reported this on a yellow card to the Committee on Safety of Medicines and in an admirably prompt and courteous reply they mentioned that they had had one previous report of an immunological abnormality in a person on oxprenolol—a woman who developed lupus erythematosus after being on oxprenolol for two weeks.

In view of the fact that people on practolol may have developed immunological abnormalities it is perhaps worth wondering if beta-blockers in general may affect the immunological responses. It would be interesting to hear if any other readers have experienced similar problems.

L F W McMAHON

Whitchurch, S Glam

Cimetidine and ulcers

SIR,—I read your leading article on this subject (27 November, p 1275) with interest. I thoroughly concur that cimetidine is no panacea for all dyspepsia, and share your concern about the possibility of indiscriminate use. The "undeserved ill repute" you mention as a possible consequence of this may also, however, follow incorrect use of the drug in perfectly conventional indications. Confusion may arise on points where your recommendations appear to be at variance with those of the data sheet for the product. As the data sheet recommendations are based strictly on the terms of the product licence under which cimetidine is released for routine clinical use, some comments may be useful.

Firstly, the impression is left that the timing of doses of cimetidine is not important. I should emphasise that the recommended dosage for most indications is 200 mg three times a day *with meals* and 400 mg *at bedtime*. This regimen was used in most of the clinical trials and should therefore lead to predictable results. Secondly, I am puzzled as to the source of your recommendation for two months' treatment, when a month to six weeks

has proved sufficient to allow healing of most duodenal ulcers and evidence on two months' treatment is scanty. I was also surprised by your abrupt dismissal of the evidence concerning the role of cimetidine in the treatment of reflux oesophagitis, one of the main permitted indications listed in the data sheet. Although this evidence may concern small numbers as yet, it is not lacking in quality.

Your recommendation concerning monthly serum creatinine estimations raises the question of what a doctor should do if he finds an increased concentration. The rises in serum creatinine reported in trials were not generally considered clinically significant; patients continued treatment with no demonstrable ill effect. Cimetidine was withdrawn in only three cases for this reason; two of these patients had pre-existing renal disease and one had an abnormal creatinine level before starting treatment. This test would seem, therefore, to be of little practical value for routine use.

Finally, you may be interpreted at one point as implying that a considerable hazard attends the use of cimetidine. No such hazard has been detected, but this should not detract from the desirability of all doctors reporting any possible adverse reactions they see, as this is the only way in which the hazards, or lack of them, of long-term and widespread use can be defined.

E M JACKSON-MOORE

Medical Director,
Smith, Kline, and French Laboratories Ltd

Welwyn Garden City,
Herts

Screening for breast cancer

SIR,—It was with much interest that I read the report by Mr W D George and others from Manchester (9 October, p 858) of the feasibility of conducting a screening programme for breast cancer using non-medical staff to carry out clinical and mammographic examinations.

If introduced, any such mass screening programme aiming to improve substantially the long-term prognosis of this disease must be capable of detecting the majority of cancers while the primary is still under 2 cm in greatest diameter, when 10-year survival rates of 80-90% may be obtained.¹ Thus the ability of non-medical staff to detect cancers at this stage must be assessed, since the methods being used are capable of doing this in medical hands, as has been demonstrated at West London and Guildford.^{2,3} The staging of the cancers detected at Manchester would therefore be most helpful in assessing the performance of the non-medical examiners, since it is possible that there could be differences between the groups in their detection rates of T₁N₀ cancers even if both detected more advanced cancers.

Presumably the "referred women" at Manchester are equivalent to the usual outpatient referrals and contain a very high proportion of later cancers. Experience at Guildford suggests that of the cancers detected in women attending the screening clinic, those occurring in symptomatic women are mainly over 2 cm in diameter with or without nodal involvement, while the majority of those detected in asymptomatic women are under 2 cm in diameter and node-free, and represent the true screening cases in which most benefit is to be expected.

Considering the Manchester results, in practice "referred women" are seen by a medical team. Of the "invited" and "self-referred," the 782 symptomatic women having 14 cancers represent a group which should ideally consult their own doctors and these cancers are, at least potentially, detectable by a health education programme without sophisticated screening as an initial stage. It is therefore only the nine women having cancers out of the 1944 asymptomatic "invited" and "self-referred" who really benefit from the screening procedure and who are most likely to have the early lesions.

Consideration of the use of non-medical staff for screening clinics must surely depend largely upon their ability to detect such cases. Here the Manchester data seem incomplete, since we are told only that the non-medical staff detected among the "invited" women, both symptomatic and asymptomatic, 10 out of 16 cancers clinically and 13 out of 16 radiologically, eight of the 16 being symptomatic. The most pessimistic interpretation of these data means that the non-medical staff could have missed three or possibly four (if they missed the only cancer in the asymptomatic "self-referred") of the nine cancers previously defined. This would presumably not be acceptable. Clearly the full data are needed, though even then any judgment based on only nine cases seems premature.

On the other hand there is clearly a high accuracy when non-medical staff assess referred women. Follow-up will be of course of interest to establish any false-negative rate of medical or non-medical staff.

BARBARA A THOMAS

Buryfields Breast Screening Clinic,
Guildford, Surrey

¹ Cutler, S J, in *Prognostic Factors in Human Breast Cancer*, ed A P M Forest and P B Kunkler, p 20. Edinburgh, Livingstone, 1968.
² Chamberlain, J, *et al*, *Lancet*, 1975, 2, 1026.
³ Thomas, B A, *Lancet*, 1975, 2, 914.

Thoracic duct cannulation for bleeding oesophageal varices

SIR,—In your leading article on "Bleeding oesophageal varices" (11 September, p 603) it was suggested that the least traumatic operation should be performed in order to control the bleeding.

Such a minor surgical procedure, not listed in your article, is cannulation of the thoracic duct and lymph drainage. This operation, carried out under local anaesthesia, was first suggested by Dumont and Mulholland.¹ They demonstrated that in cirrhotic patients cannulation of the thoracic duct and establishment of free lymph flow may reduce portal pressure, decrease ascites and liver size, and control bleeding from oesophageal varices. Since 10 out of 15 patients who bled massively from oesophageal varices stopped bleeding after cannulation of the thoracic duct through a supraclavicular approach it was concluded that this procedure was as effective as a portacaval shunt in decreasing the portal vein pressure and controlling the oesophageal bleeding. A similar enthusiasm was expressed by Bowers *et al*,² who proposed a portacaval shunt operation within 72 h after bleeding has been successfully controlled by cannulation of the thoracic duct.

This simple surgical procedure has been