The surface glycoconjugates of trypanosomatid parasites

MICHAEL A. J. FERGUSON

Department of Biochemistry, University of Dundee, Dundee DD1 4HN, UK

SUMMARY

Insect-transmitted protozoan parasites of the order Kinetoplastida, suborder Trypanosomatina, include *Trypanosoma brucei* (aetiological agent of African sleeping sickness), *Trypanosoma cruzi* (aetiological agent of Chagas' disease in South and Central America) and *Leishmania* spp. (aetiological agents of a variety of diseases throughout the tropics and sub-tropics). The structures of the most abundant cell-surface molecules of these organisms is reviewed and correlated with the different modes of parasitism of the three groups of parasites. The major surface molecules are all glycosylphosphatidylinositol (GPI)-anchored glycoproteins, such as the variant surface glycoproteins of *T. brucei* and the surface mucins of *T. cruzi*, or complex glycopospholipids, such as the lipophosphoglycans and glycoinositolphospholipids of the leishmanias. Significantly, all of the aforementioned structures share a motif of Man α 1-4GlcNAc-6-*myo*-inositol-1-HPO₄-lipid and can therefore be considered to be members of a GPI superfamily.

1. INTRODUCTION

In this article, the most abundant surface molecules (more than 100 000 copies per cell) of three trypanosomatid parasites, *Trypanosoma brucei*, *Trypanosoma cruzi* and *Leishmania* spp., will be described and correlated with their modes of parasitism. These surface molecules are glycosylphosphatidylinositol (GPI)-anchored glycoproteins and GPI-related complex glycopospholipids. Many other interesting and important surface molecules are found in these parasites but these will not be discussed here because their lower abundance means that they contribute less to the general surface architecture of the parasites.

2. THE SURFACE OF *TRYPANOSOMA BRUCEI*

The bloodstream forms of the African trypanosomes express an unusual surface coat (Vickerman & Luckins 1969) composed of about 10⁷ copies of variant surface glycoprotein (VSG) (Cross 1975) arranged as a dense monolayer of homodimers on the parasite surface. The VSG coat acts as a macromolecular diffusion barrier that prevents the approach of macromolecules, such as the components of the alternative complement pathway, to the plasma membrane while allowing the free diffusion of small nutrient molecules to underlying transmembrane transporter systems. The trypanosome genome contains several hundred VSG genes that encode immunologically distinct VSGs that are expressed one at a time by an individual cell. The sequential expression of different VSG genes (antigenic variation) enables the parasite population to evade specific humoral immune attack; the mechanism of antigenic variation has been reviewed recently (Cross 1996).

Each VSG monomer has a molecular mass of approximately 55 kDa and contains typically one or two N-linked oligosaccharides (Zamze *et al.* 1991). The amino-terminal domains of the VSGs represent about 75% of the mature polypeptide sequences and show little primary sequence homology, thereby allowing the VSGs their unique immunological identities. However, X-ray crystallographic studies indicate that they fold into similar elongated shapes, with two extended α -helical bundles per monomer (Blum *et al.* 1993), that pack into dense monolayer coat arrays. This arrangement allows the coat to be quite thick (15 nm) and means that only a limited subset of VSG B-cell-stimulating epitopes are exposed on the surface of the living trypanosome. The carboxy-terminal domains of the VSGs display some sequence homology and the VSGs can be grouped into two main subtypes based on these sequences; however, the three-dimensional structures of these domains have not been solved. The VSG molecules are finally attached to the plasma membrane through a covalent linkage from the carboxy-terminal amino acid α -carboxy group to a GPI membrane anchor (figure 1a) (Ferguson *et al.* 1988). The mature VSG GPI anchors contain the core structure of ethanolamine-HPO₄-6Man α 1-2Man α 1-6Man α 1-4GlcNAc-6PI, common to all GPI membrane anchors, but this is modified by an α -galactose side chain that is unique to VSGs. The predicted three-dimensional structure of the complete anchor suggests that it forms a large, dense plate-like structure of carbohydrate on which the VSG polypeptide sits (Homans *et al.* 1989). Thus the VSG coat can be considered as a dense GPI glycocalyx immediately adjacent to the membrane covered with the proteinaceous part of the VSG layer (see figure 2).

The surface of the procyclic form of *T. brucei*, which lives in the tsetse fly midgut, is covered with a more

(a) *Trypanosoma brucei* VSG

(b) *Trypanosoma brucei* PARP

(c) *Leishmania major* PSP

(d) *Trypanosoma cruzi* mucins

Figure 1. The structures of some trypanosomatid GPI-protein anchors. The structures of the GPI anchors of (a) *T. brucei* bloodstream-form VSG, (b) *T. brucei* procyclic PARP, (c) *L. major* PSP, and (d) *T. cruzi* metacyclic and epimastigote mucins are shown.

Figure 2. A representation of the cell surface molecular architecture of three trypanosomatid parasites. Schematic representations of the major surface molecules and a 20 nm $\times$ 20 nm section of plasma membrane of (left) a *T. brucei* bloodstream-form trypomastigote, (middle) a *Leishmania* metacyclic promastigote and (right) a *T. cruzi* metacyclic trypomastigote.

diffuse coat made up of glycoproteins called procyclins or procyclic acidic repetitive proteins (PARPs). Most trypanosome clones contain four pairs of PARP genes that encode three PARP types: two with repeat domains of -Glu-Pro- (EP-PARPs), one of which contains an N-glycosylation site adjacent to the polyanionic EP-repeat domain, and one with a repeat domain of -Gly-Pro-Glu-Glu-Thr- (GPEET-PARP) (Roditi *et al.* 1987; Mowatt & Clayton 1988). Both the N-glycosylated EP-PARP species and the GPEET-PARP species have been isolated and structurally characterized. Both forms are believed to adopt highly extended rod-like structures (Roditi *et al.* 1989; Treumann *et al.* 1997) and both contain GPI anchors with an unusual lipid structure and large polydisperse carbohydrate side chains made of branched poly-lactosamine repeats, i.e. Gal β 1-4GlcNAc repeat units (figure 1b) (Field *et al.* 1991; Ferguson *et al.* 1993; Treumann *et al.* 1997). When the parasites are grown in culture, the β -galactose termini of these side chains are substituted with sialic acid through the action of a cell surface trans-sialidase enzyme that transfers α 2-3-linked sialic acid from fetal calf serum glycoconjugates (Engstler *et al.* 1993). The significance of trans-sialylation in *T. brucei* procyclics, and the extent to which it occurs in the tsetse fly, are not clear. Overall, the PARP coat can be considered as a dense GPI glycocalyx immediately adjacent

to the membrane above which are projected the highly extended polyanionic rod-like structures of the PARP polypeptides. The significance of the three different PARP forms, described above, is unclear. However, when bloodstream-form trypanosomes transform into procyclics *in vitro* the cells generally express predominantly N-glycosylated EP-PARP, which is gradually replaced by GPEET-PARP over a period of months (Treumann *et al.* 1997). It will be interesting to analyse which forms of PARP are being expressed by trypanosomes during their complex life-cycle within the tsetse fly. Recently, tsetse fly lectins have been suggested to be involved in controlling the parasite population in the tsetse fly (Maudlin & Welburn 1994; Welburn *et al.* 1996) and it is likely that the PARPs serve as the principal ligands for these molecules.

3. THE SURFACE OF THE *LEISHMANIA* SPP.

The most abundant cell-surface molecules of the promastigote stage of the *Leishmania* are a GPI-anchored glycoprotein called promastigote surface protease (PSP), a family of glycoinositol-phospholipids (GIPLs) and lipophosphoglycan (LPG) (McConville & Ferguson 1993). The PSP GPI anchor is a very simple one (figure 1c) and the GIPLs fall into three types according to their glycan structure (McConville

et al. 1993; McConville & Ferguson 1993). The Type-1 GIPLs are more similar to the GPI protein anchors and contain a core of $\text{Man}\alpha 1\text{-6}\text{Man}\alpha 1\text{-4GlcNAc}\alpha 1\text{-6PI}$, the Type-2 GIPLs contain a core of $\text{Man}\alpha 1\text{-3}\text{Man}\alpha 1\text{-4GlcNAc}\alpha 1\text{-6PI}$, and the hybrid-GIPLs contain a branched core of $\text{Man}\alpha 1\text{-3}(\text{Man}\alpha 1\text{-6})\text{Man}\alpha 1\text{-4GlcNAc}\alpha 1\text{-6PI}$. Representative examples of the *Leishmania* GIPLs are shown in figure 3. The relative proportions of the different GIPL species varies between *Leishmania* species and between the promastigote and amastigote forms of the parasite (McConville *et al.* 1990; McConville & Blackwell 1991; McConville & Ferguson 1993; Schneider *et al.* 1993, 1994; Winter *et al.* 1994). The LPGs consist of a Type-2 GIPL membrane anchor attached to a long phosphosaccharide-repeat domain that is capped by a neutral oligosaccharide (McConville & Ferguson 1993; McConville *et al.* 1995) (figure 3). The *Leishmania* promastigote surface therefore consists of a dense GIPL glycocalyx immediately adjacent to the membrane through which the PSP and polyanionic LPG molecules are projected (figure 2).

The physicochemical similarity between the *Leishmania* cell surface and the African trypanosome procyclic cell surface is quite striking in that both surfaces have dense layers of carbohydrate immediately adjacent to the membrane with polyanionic rods displayed above. In the trypanosomes this is achieved with a single molecule (PARP) that supplies the glycocalyx in the form of its highly modified GPI anchor and the polyanionic rods in the form of extended acidic polypeptide repeats, whereas the *Leishmania* use GIPLs to form the glycocalyx and separate LPG molecules to supply the polyanionic rods in the form of sugar-phosphate repeats.

With the exception of *L. major* (Moody *et al.* 1993), the intracellular amastigote stages expresses little or no LPG and no major cell-surface glycoproteins (McConville & Blackwell 1991; Bahr *et al.* 1993). The phosphosaccharide-repeat domains of the LPGs carry side chain modifications that are species-specific (figure 3) (McConville *et al.* 1995). These differences in LPG structure seem to have a key role in defining the sandfly vector range of these organisms, suggesting that LPGs mediate parasite adhesion to putative sandfly midgut epithelium lectins (Pimenta *et al.* 1992, 1994). In *L. major* and *L. donovani*, differentiation of the non-infectious dividing procyclic promastigotes to infectious non-dividing metacyclic promastigotes is correlated with an approximate doubling in the length of the phosphosaccharide repeat domains and (for *L. major*) a change in the phosphosaccharide side chains (McConville *et al.* 1992; Sacks *et al.* 1995). These changes in LPG structure are responsible for the detachment of the parasites from the vector midgut epithelium and for their acquisition of resistance to complement. Interestingly, the metacyclic promastigote LPG is a potent activator of the classical complement pathway, and the opsonization of the metacyclic promastigote LPG layer with C3b and C3bi is essential for the successful invasion of host macrophages via the macrophage CR1/CR3 complement receptor system, reviewed in Brittingham *et al.* (1996). The resistance to complement-mediated lysis is thought to be due (a) to the length of the LPG molecule, which holds the

complement cascade far enough away from the parasite membrane to prevent insertion of the membrane attack complex, and (b) to the action of the PSP protease that can degrade C3b to C3bi-like products. The activation of complement by the classical pathway is thought to be due to the nature of the LPG cap structure (which contains α -mannose residues) that efficiently binds serum mannose-binding protein (MBP), which has a complement-activating Clq-like domain (Green *et al.* 1994). The LPG has also been implicated in inhibiting the macrophage oxidative burst, providing further protection for the parasite once inside the macrophage (Turco & Descoteaux 1992). Recently, the GIPLs and LPG (and synthetic LPG fragments) have been implicated in modulating the macrophage inducible nitric oxide synthase (iNOS) (Proudfoot *et al.* 1995, 1996) and these features are reviewed elsewhere in this volume by Liew *et al.*

The novel phosphosaccharide repeats found in the LPGs are also found attached to protein as part of the so-called proteophosphoglycans that are secreted by *Leishmania* promastigotes and amastigotes as large macromolecular filaments and networks (Ilg *et al.* 1994a,b, 1996). These large molecules might have an important role in protecting the amastigote parasite inside the macrophage (Ilg *et al.* 1996).

4. THE SURFACE OF *TRYPANOSOMA CRUZI*

The most abundant cell-surface molecules on the epimastigote and metacyclic forms of this parasite are a small family of Type-1 GIPLs (figure 3) (Previato *et al.* 1990; Lederkremer *et al.* 1991, 1993; Carreira *et al.* 1996) and a family of GPI-anchored heavily O-glycosylated mucin-like glycoproteins (Schenkman *et al.* 1993; Previato *et al.* 1994, 1995; Serrano *et al.* 1995). The *T. cruzi* GIPLs are predicted to form a dense glycocalyx immediately adjacent to the membrane, with the mucins projecting out above this layer (figure 2). The GPI anchors of the mucins are relatively simple (figure 1d) but there is an interesting change in their lipid structure when the epimastigotes differentiate into metacyclic trypomastigotes. The epimastigote mucin GPI anchors contain exclusively *sn*-1-alkyl-2-acylglycerol, whereas over 70% of the metacyclic mucin GPI anchors contain ceramide lipids (Serrano *et al.* 1995). The significance of this change in lipid structure is unclear but it might correlate with the ability of the mucins to insert into the host cell membrane during invasion, whereas another metacyclic GPI-anchored glycoprotein (the IG7 antigen), which contains predominantly *sn*-1-alkyl-2-acylglycerol (Heise *et al.* 1995), remains attached to the parasite (Schenkman *et al.* 1993).

The bloodstream forms of *T. cruzi* also express abundant GPI-anchored mucins, although these seem to be larger than the epimastigote/metacyclic mucins (Almeida *et al.* 1994), and the GPI anchor lipid component in this case is *sn*-1-alkyl-2-acylglycerol. The acyl (fatty acid) chains of the bloodstream trypomastigote GPI lipid are mostly unsaturated fatty acids ($\text{C}_{18:1}$ and $\text{C}_{18:2}$), whereas the *sn*-1-alkyl-2-acylglycerol lipids from the other life-cycle stages contain exclusively saturated

Figure 3. 1 Representative examples of the Type-1 and Type-2 GIPLs and LPGs. The arrows on the *Leishmania* GIPLs indicate the names of the smaller GIPL species (McConville & Ferguson 1993). The structure of the side chains (R) and the average number of phosphosaccharide repeats (n) of the *Leishmania* LPGs varies depending on the species and developmental stage of the parasite (McConville & Ferguson 1993; McConville *et al.* 1995). (a) *T. cruzi* Type-1 GIPL, (b) *Leishmania* Type-1 GIPLs, (c) *Leishmania* hybrid GIPLs, (d) *Leishmania* Type-2 GIPLs, and (e) *Leishmania* LPGs.

fatty acids. This subtle difference in lipid structure might be immunologically significant because the bloodstream trypomastigote mucins, but not the epimastigote and metacyclic mucins, are potent inducers of host macrophage tumour necrosis factor α and IL-12 (Carmago *et al.* 1997). The mucin O-linked oligosaccharides seem to play an important role in *T. cruzi* metacyclic and bloodstream trypomastigote infectivity. The mucins are the major acceptor substrates for the cell surface trans-sialidase enzymes of the parasite, which transfer α 2-3-linked sialic acid from host glycoconjugates to the parasite surface (Schenkman *et al.* 1994). Trans-sialylation seems to be necessary for host cell invasion (Schenkman *et al.* 1991), and the O-linked oligosaccharides that terminate in β -galactopyranose residues (e.g. Gal β 1-3(Gal β 1-6)Gal β 1-6(Gal β 1-4)GlcNAc-Thr) are the specific sialic acid acceptor sites (Previano *et al.* 1994, 1995; Serrano *et al.* 1995). The available structural data show that the O-linked oligosaccharides of the epimastigote mucins show some variation between *T. cruzi* strains (Previano *et al.* 1994, 1995) but that within a given strain the epimastigote and metacyclic structures are identical (Serrano *et al.* 1995). In contrast, the bloodstream trypomastigote mucin oligosaccharides uniquely contain α -galactose residues that are responsible for eliciting the high-levels of anti- α Gal antibodies seen in patients with Chagas' disease (Almedia *et al.* 1994). Several *T. cruzi* mucin genes belonging to a multi-gene family have been cloned (Noia *et al.* 1995; Salazar *et al.* 1996) but it is not yet clear which of them are expressed in the various life-cycle stages.

Although a small number of *Leishmania* amastigote surface glycoproteins have been identified (Andrews *et al.* 1988; Teixeira *et al.* 1994), relatively little is known about the surface architecture of this stage of the parasite.

5. OTHER TRYPANOSOMATID PARASITES AND GIPL PHYLOGENY

A number of other trypanosomatid parasites have been analysed for glycolipid/glycoconjugate content and structure. In all cases, the major molecular species belong to the GPI family; for example, Type-1 GIPLs have been described in *Phytomonas* (Redman *et al.* 1995) and Type-2 and/or hybrid GIPLs have been described in *Leptomonas samueli* (Previano *et al.* 1992), *Endotrypanum schaudinni* and *Herpetomonas samuelpeessoai* (Routier *et al.* 1995), and *Crithidia fasciculata* (Schneider *et al.* 1996). A wide variety of glycosyl groups are found attached to the aforementioned GIPLs, including α Man, α Gal, β Gal, α Glc, β Xyl, α Rha, α GlcA and D-Arap residues, and a number of phosphorus-containing substituents are also found, including ethanolamine phosphate, 2-aminoethylphosphonate and Glc α 1-HPO $_4$ -. A recent analysis of trypanosomatid small-subunit rRNA sequences (Fernandes *et al.* 1993) suggests that: (i) *T. cruzi* is more closely related to *Phytomonas* than to *Leptomonas*, *Endotrypanum* and *Leishmania*, (ii) *Leptomonas*, *Endotrypanum* and *Leishmania* are more closely related to each other than to *Phytomonas* and *T. cruzi*, and (iii) *T. brucei* is the most ancient and distantly related of all the trypanosomatids. In this

Figure 4. The phylogeny of GIPL types in the trypanosomatids. The phylogenetic tree, based on the average percentage of small-subunit ribosomal RNA divergence (X axis), is adapted from Fernandes *et al.* (1993). The monogenetic (M) and digenetic (D) life-cycles are indicated after the species name, followed by the GIPL types expressed by the species. Type-1 indicates exclusively Type-1 GIPLs and Type-2 indicates Type-2 and/or hybrid-GIPLs; see the text for details.

context, it is worth noting that: (i) although *T. brucei* expresses very high levels of protein-linked GPI anchors it does not make free GIPLs, (ii) *T. cruzi* and *Phytomonas* seem to make only Type-1 GIPLs (the type most closely related to protein-linked GPI anchors), and (iii) *Leptomonas*, *Endotrypanum*, *Leishmania* and *Crithidia* make predominantly Type-2 and/or hybrid GIPLs. Thus, one could suggest that the GIPLs have arisen through adaptations of GPI anchor biosynthesis (ubiquitous in the eukaryotes) and that the Type-1 GIPLs preceded the Type-2 GIPLs and related structures such as the *Leishmania* LPGs (figure 4).

There does not seem to be any correlation between the presence or types of GIPL expressed by trypanosomatids and their monogenetic (single host) or digenetic (host and vector) life-cycles. For example, the digenetic parasite *T. brucei* does not express GIPLs, and the monogenetic parasite *Leptomonas* expresses high levels of complex Type-2 GIPLs. The precise functions of cell-surface GIPLs in trypanosomatid parasites are largely unknown. Their abundance suggests that they might have a general protective function by providing a dense, negatively charged, glycocalyx close to the surface of the plasma membrane through which other macromolecules project. However, this function alone does not readily explain the stage- and species-specific nature of GIPL structures.

6. GPI BIOSYNTHESIS

The GPI family of molecules (GPI-anchored glycoproteins and/or the LPGs and/or the GIPLs) are the most abundant molecules on the surface of all the trypanosomatid parasites. Although these molecules show great structural diversity they also share the common core structure, Man α 1-4GlcNAc1-6myo-inositol-1-PO $_4$ -lipid. Thus, the biosynthetic enzymes responsible for the synthesis of this common core are attractive targets for the development of anti-trypanosomatid agents. In contrast, some of the unique features of GPI-anchor biosynthesis in the African trypanosomes, such as fatty acid remodelling and inositol acylation/deacylation, offer attractive targets for the

development of trypanocidal agents specific to the VSG-coated African trypanosomes. These issues are discussed in detail in Güther & Ferguson (1995) and Smith *et al.* (1996).

This work was supported by The Wellcome Trust. I thank my many colleagues and collaborators involved in the structural analyses described in this article, especially: Steve Homans, Malcolm McConville, Pascal Schneider, Angela Mehler, Achim Treumann, Nicole Zitzmann, Andreas Hülsmeier, Chris Redman, Trevor Rutherford, Julia Richardson, Jane Thomas-Oates, Jerry Thomas, Thomas Ilg, Peter Overath, Sam Turco, David Sacks, Sergio Schenkman, Nobuko Yoshida, Luiz Travassos, Alvaro Acosta Serrano, Igor de Almeida, Lucia de Almeida and Norton Heise. The author is a Howard Hughes International Research Scholar.

REFERENCES

- Almeida, I. C., Ferguson, M. A. J., Schenkman, S. & Travassos, L. R. 1994 Lytic anti- α -galactosyl antibodies from patients with chronic Chagas' disease recognize novel O-linked oligosaccharides on mucin-like glycosyl-phosphatidylinositol-anchored glycoproteins of *Trypanosoma cruzi*. *Biochem. J.* **304**, 793–802.
- Andrews, N. W., Robbins, E. S., Ley, V., Hong, K. S. & Nussenzweig, V. 1988 Developmentally regulated, phospholipase C-mediated release of the major surface glycoprotein of amastigotes of *Trypanosoma cruzi*. *J. Exp. Med.* **167**, 300–314.
- Bahr, V., Stierhof, Y.-D., Ilg, T., Demar, M., Quinten, M. & Overath, P. 1993 Expression of lipophosphoglycan, high-molecular weight phosphoglycan and glycoprotein gp63 in promastigotes and amastigotes of *Leishmania mexicana*. *Molec. Biochem. Parasitol.* **58**, 107–122.
- Blum, M. L., Down, J. A., Gurnett, A. M., Carrington, M., Turner, M. J. & Wiley, D. C. 1993 A structural motif in the variant surface glycoproteins of *Trypanosoma brucei*. *Nature* **362**, 603–609.
- Brittingham, A. & Mosser, D. M. 1996 Exploitation of the complement system by *Leishmania* promastigotes. *Parasitol. Today* **12**, 444–447.
- Carmargo, M. M., Almeida, I. C., Pereira, M. E. S., Ferguson, M. A. J., Travassos, L. R. & Gazzinelli, R. T. 1997 GPI-anchored mucin-like glycoproteins isolated from *Trypanosoma cruzi* trypomastigotes initiate the synthesis of pro-inflammatory cytokines by macrophages. *J. Immun.* (In the press.)
- Carreira, J. C., Jones, C., Wait, R., Previato, J. O. & Mendonca-Previato, L. 1996 Structural variations in the glycoinositolphospholipids of different strains of *Trypanosoma cruzi*. *Glycoconj. J.* **13**, 955–966.
- Cross, G. A. M. 1975 Identification, purification and properties of clone-specific glycoprotein antigens constituting the surface coat of *Trypanosoma brucei*. *Parasitology* **71**, 393–417.
- Cross, G. A. M. 1996 Antigenic variation in trypanosomes: secrets surface slowly. *BioEssays* **18**, 283–291.
- Engstler, M., Reuter, G. & Schauer, R. 1993 The developmentally regulated trans-sialidase from *Trypanosoma brucei* sialylates the procyclic acidic repetitive protein. *Molec. Biochem. Parasitol.* **61**, 114.
- Ferguson, M. A. J., Homans, S. W., Dwek, R. A. & Rademacher, T. W. 1988 Glycosyl-phosphatidylinositol moiety that anchors *Trypanosoma brucei* variant surface glycoprotein to the membrane. *Science* **239**, 753–759.
- Ferguson, M. A. J., Murray, P., Rutherford, H. & McConville, M. J. 1993 A simple purification of procyclic acidic repetitive protein and demonstration of a sialylated glycosylphosphatidylinositol membrane anchor. *Biochem. J.* **291**, 51–55.
- Fernandes, A. P., Nelson, K. & Beverley, S. M. 1993 Evolution of nuclear ribosomal RNAs in kinetoplastid protozoa: perspectives on the age and origins of parasitism. *Proc. Natn. Acad. Sci. USA* **90**, 11608–11612.
- Field, M. C., Menon, A. K. & Cross, G. A. M. 1991 A glycosylphosphatidylinositol protein anchor from procyclic stage *Trypanosoma brucei*: lipid structure and biosynthesis. *EMBO J.* **10**, 2731–2739.
- Green, P. J., Feizi, T., Stoll, M. S., Theil, S., Prescott, A. & McConville, M. J. 1994 Recognition of the major cell surface glycoconjugates of *Leishmania* parasites by the human serum mannan-binding protein. *Molec. Biochem. Parasitol.* **66**, 319–328.
- Güther, M. L. S. & Ferguson, M. A. J. 1995 The role of inositol-acylation and inositol-deacylation in GPI biosynthesis in *Trypanosoma brucei*. *EMBO J.* **14**, 3080–3093.
- Heise, N., Cardoso de Almeida, M. L. & Ferguson, M. A. J. 1995 Characterization of the lipid moiety of the glycosyl-phosphatidylinositol anchor of *Trypanosoma cruzi* IG7-antigen. *Molec. Biochem. Parasitol.* **70**, 71–84.
- Homans, S. W., Edge, C. J., Ferguson, M. A. J., Dwek, R. A. & Rademacher, T. W. 1989 Solution structure of the glycosylphosphatidylinositol membrane anchor glycan of *Trypanosoma brucei* variant surface glycoprotein. *Biochemistry* **28**, 2881–2887.
- Ilg, T., Stierhof, Y.-D., Weise, M., McConville, M. J. & Overath, P. 1994a Characterization of phosphoglycan-containing secretory products of *Leishmania*. *Parasitology* **108**, S63–S71.
- Ilg, T., Overath, P. O., Ferguson, M. A. J., Rutherford, T., Campbell, D. G. & McConville, M. J. 1994b O- and N-glycosylation of the *Leishmania mexicana* secreted acid phosphatase: characterization of a new class of phosphoserine linked glycans. *J. Biol. Chem.* **269**, 24073–24081.
- Ilg, T., Stierhof, Y.-D., Craik, D., Simpson, R., Bacic, A. & Handman, E. 1996 Purification and structural characterization of a filamentous, mucin-like proteophosphoglycan secreted by *Leishmania* parasites. *J. Biol. Chem.* **271**, 21583–21596.
- Lederkremer, R. M., Lima, C. E., Ramirez, M. I., Goncalves, M. F. & Colli, W. 1993 Hexadecylpalmitoylglycerol or ceramide is linked to similar glycoposphatidylinositol anchor-like structures in *Trypanosoma cruzi*. *Eur. J. Biochem.* **218**, 929–936.
- Lederkremer, R. M., Lima, C., Ramirez, M. I., Ferguson, M. A. J., Homans, S. W. & Thomas-Oates, J. E. 1991 Complete structure of the glycan of lipopeptidophosphoglycan from *Trypanosoma cruzi* epimastigotes. *J. Biol. Chem.* **266**, 23670–23675.
- Maudlin, I. & Welburn, S. C. 1994 Maturation of trypanosome infections in tsetse. *Expl Parasitol.* **79**, 202–205.
- McConville, M. J. & Blackwell, J. M. 1991 Developmental changes in the glycosylated phosphatidylinositols of *Leishmania donovani*. Characterization of the promastigote and amastigote glycolipids. *J. Biol. Chem.* **266**, 15170–15179.
- McConville, M. J., Collidge, T. A. C., Ferguson, M. A. J. & Schneider, P. 1993 The glycoinositolphospholipids of *Leishmania mexicana* promastigotes: evidence for the presence of three distinct pathways of biosynthesis. *J. Biol. Chem.* **268**, 15595–15604.
- McConville, M. J. & Ferguson, M. A. J. 1993 The structure, biosynthesis and function of glycosyl-phosphatidylinositols in the parasitic protozoa and higher-eukaryotes. *Biochem. J.* **294**, 305–324.
- McConville, M. J., Homans, S. W., Thomas-Oates, J. E., Dell, A. & Bacic, A. 1990 Structures of the glycoinositolphospholipids from *Leishmania major*: a family of novel galactofuranose-containing glycolipids. *J. Biol. Chem.* **265**, 7385–7394.
- McConville, M. J., Schnur, L. F., Jaffe, C. & Schneider, P. 1995 Structure of *Leishmania* lipophosphoglycan: inter- and intra-specific polymorphism in Old World species. *Biochem. J.* **310**, 807–818.
- McConville, M. J., Turco, S. J., Ferguson, M. A. J. & Sacks, D. L. 1992 Developmental modification of lipophosphoglycan

- during the differentiation of *Leishmania major* promastigotes to an infectious stage. *EMBO J.* **11**, 3593–3600.
- Moody, S. F., Handman, E., McConville, M. J. & Bacic, A. 1993 The structure of *Leishmania major* amastigote lipophosphoglycan. *J. Biol. Chem.* **268**, 18457–18466.
- Mowatt, M. R. & Clayton, C. E. 1988 Polymorphism in the procyclic acidic repetitive protein gene family of *Trypanosoma brucei*. *Molec. Cell. Biol.* **8**, 4055–4062.
- Noia, J. M., Sanchez, D. O. & Frasch, A. C. 1995 The protozoan *Trypanosoma cruzi* has a family of genes resembling the mucin genes of mammalian cells. *J. Biol. Chem.* **270**, 24146–24149.
- Pimenta, P. F., Turco, S. J., McConville, M. J., Lawyer, P. G., Perkins, P. V. & Sacks, D. L. 1992 Stage-specific adhesion of *Leishmania* promastigotes to the sandfly midgut. *Science* **256**, 1812–1815.
- Pimenta, P. F., Saraiva, E. M. B., Rowton, E., Modi, G. B., Garraway, L. A., Beverley, S. M., Turco, S. J. & Sacks, D. L. 1994 Evidence that the vectorial competence of phlebotomine sand flies for different species of *Leishmania* is controlled by structural polymorphisms in the surface lipophosphoglycan. *Proc. Natn. Acad. Sci. USA* **91**, 9155–9159.
- Previato, J. O., Mendonca-Previato, L., Jones, C. & Fournet, B. 1992 Structural characterisation of a novel class of glycosphosphingolipids from the protozoan *Leptomonas samueli*. *J. Biol. Chem.* **267**, 24279–24286.
- Previato, J. O., Jones, C., Goncalves, L. P. B., Wait, R., Travassos, L. R. & Medonca-Previato, L. 1994 O-glycosidically linked N-acetylglucosamine-bound oligosaccharides from glycoproteins of *Trypanosoma cruzi*. *Biochem. J.* **301**, 151–159.
- Previato, J. O., Gorin, P. A. J., Mazurek, M., Xavier, M. T., Fournet, B., Wieruszkes, J. M. & Mendonca-Previato, L. 1990 Primary structure of the oligosaccharide chain of lipopeptidophosphoglycan of epimastigote forms of *Trypanosoma cruzi*. *J. Biol. Chem.* **265**, 2518–2526.
- Previato, J. O., Jones, C., Xavier, M. T., Wait, R., Travassos, L. R., Parodi, A. J. & Mendonca-Previato, L. 1995 Structural characterisation of the major glycosylphosphatidylinositol membrane anchored glycoprotein from epimastigote forms of *Trypanosoma cruzi* strains. *J. Biol. Chem.* **270**, 7241–7250.
- Proudfoot, L., Nikolaev, A. V., Ferguson, M. A. J., Brimacombe, J. S. & Liew, F. Y. 1996 Regulation of the expression of nitric oxide synthase and leishmanicidal activity by glycoconjugates of *Leishmania* lipophosphoglycan in murine macrophages. *Proc. Natn. Acad. Sci. USA* **93**, 10984–10989.
- Proudfoot, L., O'Donnell, C. A. & Liew, F. Y. 1995 Glycoinositolphospholipids of *Leishmania major* inhibit nitric oxide synthesis and reduce leishmanicidal activity in murine macrophages. *Eur. J. Immunol.* **25**, 745–750.
- Redman, C. A., Schneider, P., Mehlert, A. & Ferguson, M. A. J. 1995 The glycoinositol-phospholipids of *Phytomonas*. *Biochem. J.* **311**, 495–503.
- Roditi, I., Carrington, M. & Turner, M. 1987 Expression of a polypeptide containing a dipeptide repeat is confined to the insect stage of *Trypanosoma brucei*. *Nature* **325**, 272–274.
- Roditi, I., Schwarz, H., Pearson, T. W., Beecroft, R. P., Liu, M. K., Richardson, J. P., Bühring, H.-J., Pleiss, J., Bülow, R., Williams, R. O. & Overath, P. 1989 Procyclin gene expression and loss of the variant surface glycoprotein during differentiation of *Trypanosoma brucei*. *J. Cell Biol.* **108**, 737–746.
- Routier, F. H., da Silveira, E. X., Wait, R., Jones, C., Previato, J. O. & Mendonca-Previato, L. 1995 Chemical characterisation of glycosylinositolphospholipids of *Herpetomonas samuelipessoai*. *Molec. Biochem. Parasitol.* **69**, 81–92.
- Sacks, D. L., Pimenta, P. F., McConville, M. J., Schneider, P. & Turco, S. J. 1995 Stage-specific binding of *Leishmania donovani* to the sand fly vector midgut is regulated by conformational changes in the abundant surface lipophosphoglycan. *J. Exp. Med.* **181**, 685–697.
- Salazar, N. A., Mondragon, A. & Kelley, J. M. 1996 Mucin-like glycoprotein genes are closely linked to members of the trans-sialidase super-family at multiple sites in the *Trypanosoma cruzi* genome. *Molec. Biochem. Parasitol.* **76**, 91–103.
- Schenkman, S., Eichinger, D., Pereira, M. A. & Nussenzweig, V. 1994 Structural and functional properties of *Trypanosoma* trans-sialidase. *A. Rev. Microbiol.* **48**, 499–523.
- Schenkman, S., Ferguson, M. A. J., Heise, N., Cardoso de Almeida, M. L., Mortara, R. A. & Yoshida, N. 1993 Mucin-like glycoproteins linked to the membrane by glycosyl-phosphatidylinositol anchor are the major acceptors of sialic acid in a reaction catalysed by trans-sialidase in metacyclic forms of *Trypanosoma cruzi*. *Molec. Biochem. Parasitol.* **59**, 293–304.
- Schenkman, S., Jiang, M.-S., Hart, G. W. & Nussenzweig, V. 1991 A novel cell surface trans-sialidase of *Trypanosoma cruzi* generates a stage-specific epitope required for invasion of mammalian cells. *Cell* **65**, 1117–1125.
- Schneider, P., Rosal, J.-P., Ransijn, A., Ferguson, M. A. J. & McConville, M. J. 1993 Characterization of glycoinositol phospholipids in the amastigote stage of the protozoan parasite *Leishmania major*. *Biochem. J.* **295**, 555–564.
- Schneider, P., Schnur, L. F., Jaffe, C. L., Ferguson, M. A. J. & McConville, M. J. 1994 Glycoinositol-phospholipid profiles of four serotypically distinct Old World *Leishmania* strains. *Biochem. J.* **304**, 603–609.
- Schneider, P., Treumann, A., Milne, K. G., McConville, M. J., Zitzmann, N. & Ferguson, M. A. J. 1996 Structural studies on a lipoarabinogalactan of *Crithidia fasciculata*. *Biochem. J.* **313**, 963–971.
- Serrano, A. A., Schenkman, S., Yoshida, N., Mehlert, A., Richardson, J. M. & Ferguson, M. A. J. 1995 The lipid structure of the GPI-anchored mucin-like sialic acid acceptors of *Trypanosoma cruzi* changes during parasite differentiation from epimastigotes to infective metacyclic trypomastigote forms. *J. Biol. Chem.* **270**, 27244–27253.
- Smith, T. K., Cottaz, S., Brimacombe, J. S. & Ferguson, M. A. J. 1996 Substrate specificity of the dolichol phosphate mannose: glucosaminyl phosphatidylinositol α 1-4 mannosyltransferase of the glycosylphosphatidylinositol biosynthetic pathway of African trypanosomes. *J. Biol. Chem.* **271**, 6476–6482.
- Teixeira, S. M., Russell, D. G., Kirchhoff, L. V. & Donelson, J. E. 1994 A differentially expressed gene family encoding 'amastin', a surface protein of *Trypanosoma cruzi* amastigotes. *J. Biol. Chem.* **269**, 20509–20516.
- Treumann, A., Zitzmann, N., Hülsmeier, A., Prescott, A. R., Almond, A., Sheehan, J. & Ferguson, M. A. J. 1997 Structural characterisation of two forms of procyclic acidic repetitive protein expressed by procyclic forms of *Trypanosoma brucei*. *J. Molec. Biol.* **269**, 529–547.
- Turco, S. J. & Descoteaux, A. 1992 The lipophosphoglycan of *Leishmania* parasites. *A. Rev. Microbiol.* **46**, 65–94.
- Vickerman, K. & Luckins, A. G. 1969 Localization of variable antigens in the surface coat of *Trypanosoma brucei* using ferritin-conjugated antibody. *Nature* **224**, 1125–1126.
- Welburn, S. C., Dale, C., Ellis, D., Beecroft, R. & Pearson, T. W. 1996 Apoptosis in procyclic *Trypanosoma brucei rhodesiense* *in vitro*. *Cell Death Differentn* **3**, 229–236.
- Winter, G., Fuchs, M., McConville, M. J., Stierhof, Y.-D. & Overath, P. 1994 Surface antigens of *Leishmania mexicana* amastigotes: characterization of glycoinositol phospholipids and a macrophage-derived glycosphingolipid. *J. Cell Sci.* **107**, 2471–2482.
- Zamze, S. E., Ashford, D. A., Wooten, E. W., Rademacher, T. W. & Dwek, R. A. 1991 Structural characterization of the asparagine-linked oligosaccharides from *Trypanosoma brucei* type II and type III variant surface glycoproteins. *J. Biol. Chem.* **266**, 20244–20261.