Wound healing in the liver with particular reference to stem cells

Malcolm Alison, Matthew Golding, El-Nasir Lalani and Catherine Sarraf

Histopathology Department, Division of Investigative Science, Imperial College School of Medicine at Hammersmith Campus, Du Cane Road, London W12 0NN, UK

The efficiency of liver regeneration in response to the loss of hepatocytes is widely acknowledged, and this is usually accomplished by the triggering of normally proliferatively quiescent hepatocytes into the cell cycle. However, when regeneration is defective, tortuous ductular structures, initially continuous with the biliary tree, proliferate and migrate into the surrounding hepatocyte parenchyma. In humans, these biliary cells have variously been referred to as ductular structures, neoductules and neocholangioles, and have been observed in many forms of chronic liver disease, including cancer. In experimental animals, similar ductal cells are usually called oval cells, and their association with impaired regeneration has led to the conclusion that they are the progeny of facultative stem cells. Oval cells are of considerable biological interest as they may represent a target population for hepatic carcinogens, and they may also be useful vehicles for *ex vivo* gene therapy for the correction of inborn errors of metabolism.

This review proposes that the liver harbours stem cells that are located in the biliary epithelium, that oval cells are the progeny of these stem cells, and that these cells can undergo massive expansion in their numbers before differentiating into hepatocytes. This is a conditional process that only occurs when the regenerative capacity of hepatocytes is overwhelmed, and thus, unlike the intestinal epithelium, the liver is not behaving as a classical, continually renewing, stem cell-fed lineage. We focus on the biliary network, not merely as a conduit for bile, but also as a cell compartment with the ability to proliferate under appropriate conditions and give rise to fully differentiated hepatocytes and other cell types.

Keywords: oval cells; liver; bile ducts; hepatocellular carcinoma; cholangiocarcinoma; stem cells

1. LIVER DEVELOPMENT

The liver primordium consists of endodermal and mesodermal components. Ventral foregut endoderm gives rise to hepatoblasts that form the parenchyma, whereas both sinusoidal-lining cells and connective tissue components originate from the mesenchymal tissue invaded by the hepatic cords (Shiojiri *et al.* 1991). The primitive hepatoblasts that surround the portal mesenchyme form a double-layered cylinder of hepatocytes, the 'ductal plate', which remodels and migrates into the mesenchyme to form the intrahepatic bile ducts (Van-Eyken *et al.* 1988a; Shiojiri *et al.* 1991). Rat hepatoblasts synthesize the hepatocyte-specific proteins alpha-fetoprotein (AFP) and albumin as they migrate into the portal stroma and, in addition, begin to express the bile-duct-specific enzyme, gamma glutamyl transferase (GGT) activity. Initially, in the intrahepatic bile ducts, intermediate filament expression is restricted to cytokeratins (CK) 8 and 18, but during the later stages of ductular morphogenesis, the ductal cells begin to express the characteristic biliary CKs 7, 8, 18 and 19 (Moll *et al.* 1982), although some continue to express hepatocyte traits such as AFP and albumin synthesis for the first 7–14 days after birth (Shiojiri *et al.* 1991). Thus, the hepatoblasts adjacent to the portal mesenchyme, which give rise to the intrahepatic bile ducts, transiently express both hepatocyte and ductal markers and can be considered to be oval cell equivalents. Hepatoblasts not in

contact with the portal mesenchyme differentiate into hepatocytes that form the liver cell plates, although they continue to have GGT activity until birth. The hepatocyte cytokeratin complement is, of course, restricted to 8 and 18.

In humans at 8–14 weeks gestation, bipotential progenitor cells may be both CK14- and CK19-positive, and commitment to the biliary lineage might be signalled by increased CK19 expression, loss of CK14 and transient expression of vimentin (Haruna *et al.* 1996). Short-lived expression of vimentin is also a feature of oval cells before they differentiate. In the mouse too, hepatoblasts from the liver diverticulum on day 9.5 of gestation are bipotential, and *in vitro* such cells can be made to differentiate into either hepatocytes or biliary cells (Rogler 1997).

2. ORGANIZATION OF THE BILIARY TREE

Periportal hepatoblasts give rise to bile ducts during liver development (the hepatocytogenetic theory of bile duct development), and in the adult the bile ducts remain connected to the hepatocyte parenchyma via the complex anastomoses of the bile canaliculi (intralobular passages bound by parenchymal cells). The canicular network drains the bile produced by the hepatocytes to the portal tract interface. The bile initially passes through ductules composed of a limiting plate hepatocyte and specialized duct cells (Steiner & Carruthers 1961), and more distally

cholangioles (also called canals of Hering) lined exclusively by squamous ductular cells (see figure 2). From here bile passes into the peripheral or marginal interlobular ducts, and then into larger septal bile ducts (Millward-Sadler & Jezequel 1992), ultimately entering into the duodenum via the extrahepatic bile ducts.

3. THE PROLIFERATIVE ORGANIZATION OF THE ADULT LIVER

There has been a considerable debate concerning the proliferative organization of the liver. In rapidly renewing epithelia (gut and skin), the stem cells represent the ancestors of the migrating pathways (Potten & Loeffler 1990), and in the liver too, it has been suggested that the hepatic plates represent the trajectory along which cells migrate centrifugally from the periportal areas to the centrilobular regions, where they ultimately die by apoptosis (Zajicek *et al.* 1985). Following the fate of tritiated thymidine-labelled cells in adult rat liver, it was claimed that over 5 weeks hepatocytes moved a distance of 46 μm (*ca.* 1 cell) from the portal rim towards the hepatic vein. The so-called 'streaming liver' hypothesis has engendered considerable debate and not a little heat (Grisham 1994; Correspondence 1995), and models of hepatic organization have been based on the hypothesis (Sigal *et al.* 1992). Unfortunately, depicting the plates of the hepatic muralium as straight cords is misleading and, moreover, models that divide plates into proliferative, maturing and terminally differentiated compartments are too simplistic. All remaining hepatocytes can enter the cell cycle after a two-thirds partial hepatectomy (2/3PH). Furthermore, all hepatocytes are equally proliferation-competent by 48 h after surgery, and even the well-known periportal concentration of proliferative cells in the early post-operative period can be reversed by altering the direction of the afferent blood supply (Wright & Alison 1984). Studies using genetic labelling with an *Escherichia coli* β -galactosidase gene certainly do not support the 'streaming liver' hypothesis. For example, when rat hepatocytes are labelled *in vivo* at 24 h after a 2/3PH by an amphotropic retrovirus carrying the β -galactosidase gene coupled to a nuclear localization signal, then initially most labelled cells are located in the periportal and mid-zonal regions; as the distribution of labelled cells did not change over the proceeding 15 months, this strongly argued against the streaming liver hypothesis (Bralet *et al.* 1994). A similar conclusion was reached from studies of transgenic mice where the β -galactosidase gene was driven by a human α 1-antitrypsin promoter (Kennedy *et al.* 1995). Here, neither postnatal growth nor a subsequent partial hepatectomy changed the distribution of labelled cells; the labelled cells simply presented as larger cell clusters with the passage of time.

4. GROWTH REACTIONS IN THE LIVER

Despite the very limited proliferative activity in the adult liver, probably resulting from 'wear and tear' loss, the liver is able to mount a prompt proliferative response to hepatotoxic insult and other hyperplastic stimuli. These reactions may be categorized as follows: (i) regenerative or compensatory hyperplasia; (ii) additive liver growth; and (iii) stem cell-driven (oval cell) reactions.

A wide variety of xenobiotics, e.g. paracetamol and carbon tetrachloride, cause parenchymal cell death. Such xenobiotics invariably, although not exclusively, cause centrilobular necrosis as the affected cells possess the enzymic capacity to metabolize the compounds to their hepatotoxic metabolites. A useful model to study regeneration has been the rat liver after a 2/3PH. Resection triggers the hepatocytes in the remnant lobes to exit the G_0 phase and, after a lag of at least 15 h, to enter DNA synthesis (Gerlach *et al.* 1997). Because this response actually occurs in quite separate lobes, this is strictly speaking viewed as a compensatory hyperplasia, although in practice the terms 'regeneration' and 'compensatory hyperplasia' are used synonymously. Figure 1a illustrates the spatial heterogeneity in hepatocyte proliferation at 24 h after 2/3PH, but at later times cell proliferative activity is randomly located across the liver (Wright & Alison 1984; Alison 1986). Of course, the reconstitution of the liver architecture also involves the proliferation of all other cell types normally present in the liver, although their amplification tends to be delayed in comparison to hepatocytes. A second type of growth reaction occurs in response to a wide variety of xenobiotics that are not cytotoxic, but instead can be substrates (inducers) for the cytochrome P450 family of enzymes (phenobarbitone) or peroxisome proliferators such as the hypolipidaemic agent called nafenopin. These agents stimulate hypertrophy and/or hyperplasia (figure 1b–d), but withdrawal of the stimulus triggers a rapid reversion to normal size by invoking hepatocyte apoptosis.

A third category of liver growth does not involve hepatocytes, emanating instead from interlobular bile ducts. The extensive formation of biliary-derived ductules emerging from the portal tracts of rodents fed chemical carcinogens was first noted in 1944 (Opie 1944). Since then a number of experimental regimes have been used to induce bile ductular hyperplasia in the rodent liver (figure 1e), and these have been reviewed elsewhere (Aterman 1992; Burt & MacSween 1993; Sell 1994). Farber (1955) used the ambiguous term 'oval cells' to describe the cells that formed the expanding biliary network during such feeding regimes. The importance of such biliary hyperplasia was not appreciated until 1958 (Wilson & Leduc 1958) when it was shown by electron microscopy that oval cells, or as they called them, 'cholangioles', ultimately differentiated into cells that had all the morphological features of mature hepatocytes. As we shall see, the evidence that these oval cells can differentiate into hepatocytes is now overwhelming (Onoe *et al.* 1975; Evarts *et al.* 1987, 1989; Vandersteenhoven *et al.* 1990; Lemire *et al.* 1991; Novikoff *et al.* 1991; Nomoto *et al.* 1992; Dabeva & Shafritz 1993; Tarsetti *et al.* 1993; Factor *et al.* 1994; He *et al.* 1994; Sarraf *et al.* 1994; Sirica *et al.* 1994; Golding *et al.* 1995; Yavorkovsky *et al.* 1995; Alison *et al.* 1996, 1997; Yasui *et al.* 1997).

5. OVAL CELLS ARE THE PROGENY OF BILIARY-LOCATED, FACULTATIVE STEM CELLS

During embryological development, AFP transcription is the earliest example of liver-specific gene expression in determined endodermal cells (Shiojiri *et al.* 1991); it is followed one day later by albumin synthesis, during

Figure 1. The variety of growth reactions that can be induced in the rat liver. (a) At 24 h after a 2/3 PH many hepatocytes, primarily located around the portal tracts (PT), are in the phase of DNA synthesis as demonstrated here by the presence of histone mRNA (non-isotopic *in situ* hybridization) in the cytoplasm. (b) The appearance of a typical hepatocyte with perinuclear glycogen (g). Bar, 3.4 μ m. (c) In contrast, a rat liver exposed to nafenopin is characterized by hypertrophic hepatocytes, which undergo mitosis (m); and contain numerous electron-dense peroxisomes (p). Bar, 5.9 μ m. (d) Rats treated with a cytochrome P450 enzyme inducer such as phenobarbitone have hypertrophic hepatocytes with augmented smooth endoplasmic reticulum (s). Bar, 1.4 μ m. (e) The appearance of oval cells in an appropriately stimulated animal, see table 1. The cells typically have euchromatic nuclei, a high nuclear/cytoplasmic ratio and sparse organelles; mitoses (m) are common. Bar, 3.6 μ m.

which time formation of the hepatic cords begins (Nagy *et al.* 1994). The primitive intrahepatic bile ducts also express biliary epithelial markers in addition to these hepatocyte proteins, and so have been called 'transitional cells' (Fausto *et al.* 1993; Sell & Pierce 1994), which are believed to remain in the adult liver as bipotential progenitors for

both hepatocytes and biliary cells (Sell 1990; Sell *et al.* 1991; Shiojiri *et al.* 1991; Grisham *et al.* 1993; Blouin *et al.* 1995). Indeed, the lineage commitment of foetal hepatoblasts can be altered *in vitro* by manipulating specific culture conditions (Germain *et al.* 1988b; Marceau 1994; Blouin *et al.* 1995).

Figure 2. Possible locations of facultative liver stem cells within the biliary tree. Bile passes through canaliculi, which are channels between adjacent hepatocytes, and enters (1) cholangioles, which progressively become composed of specialized ductular cells, before feeding into (2) the small marginal interlobular ducts, which eventually coalesce into the larger (3) septal ducts. Cell kinetic and functional studies suggest that the stem cells are primarily in the cholangioles and small marginal ducts.

Foetal liver-specific gene expression is recapitulated during oval cell proliferation by the re-emergence of transitional cell 'equivalents', represented by oval cells (Sell 1980; Sell *et al.* 1981; Germain *et al.* 1988a; Lemire & Fausto 1991; Golding *et al.* 1995). Oval cells represent the progeny of activated stem cells (Sell 1993b), a belief supported by the fact that they express markers considered characteristic of stem cells, including stem cell factor (SCF) (Fujio *et al.* 1994) and its receptor, c-kit (Fujio *et al.* 1994), bcl-2 (Burt & MacSween 1993) and cytokeratin 14 (Thorgeirsson *et al.* 1993; Bisgaard *et al.* 1994; Haque *et al.* 1996). It is not clear if liver stem cells have a specific location, but there are several possible sites (figure 2). The simple fact remains that all biliary cells in the smaller interlobular ducts are heavily implicated, at least in the modified Solt–Farber procedure (see figure 4), while the larger ducts appear to be mere bystanders. Nevertheless, several studies have alluded to stem cells being a small sub-population of ductular cells. Novikoff *et al.* (1996) made an ultrastructural analysis of oval cell proliferation during the Solt–Farber protocol, and identified occasional 'blast-like cells' beneath the lining epithelial cells of biliary ductules. These cells were only 3–5 µm in diameter, smaller than the smallest cholangiocytes (6 µm) detected in rat liver (Alpini *et al.* 1996). These blast-like cells were not in contact with either the basal lamina or the lumen of the bile ductule, but were enveloped by neighbouring ductular cells. These cells are certainly possible candidates for the liver stem cells; they are unpolarized and proliferate, they have a very dense heterochromatic nucleus, do not possess any junctional complexes, and lack expression of any differentiation markers, e.g. CK19 or even the ubiquitously expressed cytoskeletal protein, actin. We have also demonstrated the presence of occasional CK19-negative and also intensely CK18-positive cholangiocytes in small bile ducts of rats treated by the modified Solt–Farber procedure, a model in which animals are fed

Figure 3. Photomicrograph illustrating two intensely CK18-immunoreactive biliary cells (arrows) within a portal tract; such cells could represent a distinct subpopulation of potential liver stem cells. Magnification × 200.

the hepatotoxin 2-acetylaminofluorene (AAF) before and after a 2/3PH (figure 3). A possible reason for the selective upregulation of CK18 in these rare cholangiocytes can be made from analogy with the cytokeratin profile of the intestinal epithelium. Flint *et al.* (1994) showed that CK8 and CK19 are uniformly distributed along the crypt–villus axis, but that CK18 is restricted to the proliferative crypt, whereas Calnek & Quaroni (1993) have shown that CK18 and CK19 mRNA are restricted to crypt cells as opposed to CK8 transcripts, which showed a uniform distribution along the crypt–villus axis. Interestingly, similarly intensely CK18-immunoreactive 'brush cells' with a speculated chemoreceptive function have been found randomly scattered among much weaker-staining epithelial cells of the pancreatic ducts (Hofer & Drenckhahn 1996). Thus, an abundance of CK18 in particular may signify a different proliferative potential among a hitherto morphologically similar population of ductular cells.

Various opinions exist as to the site of the stem cells. Cells situated within or in contact with the portal stroma (Yavorkovsky *et al.* 1995), small non-descript cells around cholangioles, the 'periductular cells' (Sell & Salman 1984; Sell 1993b), the cholangioles (sometimes called canals of Hering, terminal biliary ductules or transition ducts; see figure 2) (Grisham & Porta 1964; Sell 1990, 1993b; Lemire *et al.* 1991; Factor *et al.* 1994), and small interlobular ducts (Nomoto *et al.* 1992; Anilkumar *et al.* 1995; Golding *et al.* 1995) have all been mooted. Having said this, it is still likely that under particular circumstances any component of the intrahepatic biliary tree can give rise to oval cells (Lenzi *et al.* 1992; Golding *et al.* 1995). This is consistent with the observation that hepatocellular carcinomas can develop from extrahepatic bile ducts (Park *et al.* 1991), even though their embryological development is independent of intrahepatic bile ducts (Shiojiri *et al.* 1991).

6. MODELS OF OVAL CELL ACTIVATION

Irrespective of the experimental regime used, oval cells form cords and ductules continuous with interlobular bile ducts (Dunsford *et al.* 1985; Makino *et al.* 1988; Lenzi *et al.* 1992; Sarraf *et al.* 1994; see figure 4), and seemingly

Figure 4. Characteristics of an oval cell reaction induced in the rat liver using the AAF/PH protocol (see text for details). (a) The reaction begins in the small bile ducts (arrows) in the portal space, which signal their commitment towards the hepatocyte lineage by massive upregulation of AFP expression. Magnification $\times 200$. (b) Oval cells form branching duct-like structures that migrate outwards from each portal tract (PT). Magnification $\times 200$. (c) These ductular oval cells (o) are found in the space of Disse, bordered on one side by an endothelial cell (e) with attenuated cytoplasm (arrows), and on the other side by stellate cells (s), which produce collagen. Bar, $2.2\ \mu\text{m}$. (d) The ductular oval cells, radiating from each portal tract (PT), are intensely CK19-immunoreactive, signifying their biliary heritage. Magnification $\times 100$. (e) In most oval cell models, the ductular oval cells are the only epithelial cells to proliferate, as demonstrated here by bromodeoxyuridine immunoreactivity confined to the profusion of ductular cells. Note that the ductules have not quite reached the draining hepatic veins (HV). Magnification $\times 100$.

similar bile ductular proliferation is frequently encountered in a variety of human liver diseases where substantial parenchymal damage occurs (Vandersteenhoven *et al.* 1990; De-Vos & Desmet 1992; Hsia *et al.* 1992; Koukoulis *et al.* 1992; Ray *et al.* 1993). Hepatocyte differentiation can even occur within the interlobular bile ducts in human liver (Nomoto *et al.* 1992), observations that have led to the belief that the small biliary cells that repopulate severely damaged liver parenchyma can function as a progenitor cell population for new hepatocytes. Certainly in rodents, early reactive bile ductules do not resemble hepatocytes (figure 1e), but later acquire features of hepatocytes (see figure 5).

Table 1 summarizes the commonly adopted rodent models for oval cell proliferation; all produce similar early morphological changes, although the potential of the ductular cells to differentiate into hepatocytes varies greatly. For example, feeding rats with furan (Elmore & Sirica 1991; Sirica *et al.* 1994) induces massive biliary cell proliferation with a few cells expressing hepatocyte features. In contrast, feeding rodents dipin (Engelhardt *et al.* 1990; Factor *et al.* 1994), acetylaminofluorene (AAF) (Sarraf *et al.* 1994; Anilkumar *et al.* 1995; Golding *et al.* 1995), or a choline-deficient ethionine-containing (CDE) diet (Hayner *et al.* 1984; Yaswen *et al.* 1985; Lenzi *et al.* 1992; Hiruma *et al.* 1993), results in similar ductular

Figure 5. The differentiation of oval cell ductules into small hepatocytes. (a) Electron micrograph of a 'juvenile' hepatocyte in the space of Disse adjacent to the sinusoid lumen (L). In contrast to oval cell progenitors (see figure 1e), here there are organelles such as rough endoplasmic reticulum (r) and microvillus border (mv). Bar, 2 μ m. (b) Small hepatocytes now surround each portal tract (PT) and expression of the biliary 'marker', CK19, is restricted to the periphery of these islands of new cells, where the biliary ductules have yet to differentiate into hepatocytes. Contrast with figure 4d, where the CK19-positive, intralobular biliary ducts are in obvious continuity with the 'parent' ducts in the portal tracts. Magnification $\times 200$.

reactions with more substantial hepatocyte differentiation. Alternatively, allyl alcohol-induced periportal necrosis causes proliferation of biliary cells that do not form typical ductular structures, yet they still rapidly differentiate into small hepatocytes (Yavorkovsky *et al.* 1995). In contrast, ligation of the common bile duct (Sirica *et al.* 1985), feeding 4,4'-diaminodiphenylmethane or α -naphthyl isothiocyanate (ANIT) (Dunsford *et al.* 1985) induces bile duct proliferation with apparently no hepatocyte differentiation. It is important to distinguish these authentic biliary proliferations from those derived from a ductular metaplasia of hepatocytes. Metaplasia is a common consequence of chronic parenchymal damage, and, for example, can be induced in guinea pigs by

feeding ANIT (Bhathal & Christie 1969), where bile canalicular ectasia was considered responsible for the 'tubularization' of hepatic plates. In diseased human liver, a similar tubularization of hepatic plates is believed to result from a ductular metaplasia of hepatocytes, and not through hepatocytic differentiation of ductular cells (Uchida & Peters 1983; Van-Eyken *et al.* 1988b, 1989; Meybehm *et al.* 1993; Delladetsima *et al.* 1995), even though some metaplasias express dual biliary/hepatocyte phenotypes (Van-Eyken *et al.* 1988b, 1989; Vandersteenhoven *et al.* 1990).

The characteristics of each ductular reaction are clearly dependent on the experimental regime used, primarily the administered hepatotoxin that impedes hepatocyte regeneration in the face of liver injury (Germain *et al.* 1988a). Each hepatotoxin could be acting in a highly specific manner resulting in the selective expansion of one or more ductal cell populations. For example, CDE diets induces cells of a 'transitional' biliary/hepatocyte phenotype, whereas rats fed on dimethylaminobenzidine (Germain *et al.* 1988b), or ANIT (Dunsford *et al.* 1985), results predominantly in a proliferation of cells committed to the biliary lineage. Feeding rats with furan induces a massive proliferation of ductular structures containing single 'hepatocyte-like cells' (Sirica *et al.* 1994). This suggests that furan induces a preferential proliferation of cells committed to the biliary phenotype, and/or inhibits proliferation/differentiation of hepatocyte precursor cells. The species, sex and strain of animal also have profound influences on the fate of ductular reactions (He *et al.* 1994), probably owing to different metabolic sensitivities to chemical carcinogens (Huitfeldt *et al.* 1990, 1991; Alpini *et al.* 1992; Yang *et al.* 1993). This is related to the differential expression of the various activating and detoxification systems, the phase I and phase II metabolic enzymes (Aterman 1992). For example, AAF used in the modified Solt-Farber regime (AAF/PH) is metabolized to the cytotoxic/mitoinhibitory *N*-hydroxy derivative by phase I metabolic enzymes (Kroese *et al.* 1990). Biliary cells and oval cells express very low amounts of phase I, and high levels of phase II enzymes compared with hepatocytes, and this favours detoxification over activation of hepatocarcinogens (Mathis *et al.* 1989; Sirica *et al.* 1990). Therefore, cultured oval cells, unlike hepatocytes, are resistant to the toxic effects of carcinogens (Ledda *et al.* 1983). Thus, in the presence of AAF, oval cells rather than hepatocytes proliferate after 2/3PH. In this model, [3 H]-thymidine-labelled cholangioles are seen as early as 4 h after PH (Thorgeirsson *et al.* 1993), and proliferation is well underway by 24 h (Evarts *et al.* 1993; Thorgeirsson *et al.* 1993; Anilkumar *et al.* 1995). The larger septal bile ducts, however, follow the time-course of biliary proliferation seen after an uncomplicated PH (Wright & Alison 1984), with significant numbers entering S-phase only 72 h after PH (Evarts *et al.* 1993; Thorgeirsson *et al.* 1993).

Thus, purely on kinetic grounds we can distinguish oval cell proliferation from the essentially intraportal bile duct proliferation accompanying a simple 2/3PH, which is probably activated by an expansion of lobular size (Grisham 1962; Fabrikant 1968). Furthermore, there are clear morphological and phenotypic differences between oval cell proliferation and some other bile duct hyperplasias. While oval cells radiate from the portal areas

Table 1. *Experimental manoeuvres that cause biliary epithelial cell hyperplasia in the rodent liver, and the subsequent fate of these newborn cells*

regime	observations	reference
bile duct ligation		
ligation of the common bile duct of male rats for 1 day to 4 weeks	massive bile duct hyperplasia with moderate peri-biliary fibrosis and no hepatocyte differentiation or necrosis. Bile ducts have well-defined ductular profiles, do not penetrate parenchyma but remain confined to expanded portal stroma	Lenzi <i>et al.</i> (1992)
	as above, but occasional proliferating biliary cells express albumin and AFP mRNA	Alpini <i>et al.</i> (1992)
	as above, but did not express hepatocyte-enriched transcription factors or AFP mRNA	Bisgaard <i>et al.</i> (1996)
4,4'-diaminodiphenylmethane (DDPM)		
male rats fed basal diet containing 0.15% 4,4'-diaminodiphenylmethane for 1–12 weeks	as for bile duct ligation, with no expression of AFP or albumin	Sell (1983)
α-naphthyl isothiocyanate (ANIT)		
male rats fed casein diet containing 0.6% ANIT for 14–28 days	similar reaction to bile ligated livers, no AFP expression	Dunsford <i>et al.</i> (1985)
male rats fed standard diet containing 0.1% ANIT for up to 30 days	as above, but occasional cells in hyperplastic ducts express albumin and AFP mRNA	Alpini <i>et al.</i> (1992)
male rats fed ANIT by oral gavage (50 mg kg ⁻¹ body weight (BW) per day) for 3–20 days	necrosis of interlobular bile ducts within 3 days, followed by biliary cell proliferation and fibrosis by day 5. 'Indistinguishable' from normal bile duct hyperplasia	Carthew <i>et al.</i> (1989)
2-acetylaminofluorene alone		
male and female rats dosed by daily oral gavage of 10 mg kg ⁻¹ BW of either 2-AF, 2-AAF or N-OH-AAF for 1–9 days	no parenchymal necrosis. Proliferation of Hering duct cells and periductular cells, which express AFP, albumin and hepatocyte transcription factors within 24 h of carcinogen exposure. Potency of stimulus: N-OH-AAF > 2-AAF > 2-AF. No extensive proliferation and migration of these oval cells	Bisgaard <i>et al.</i> (1996)
2-acetylaminofluorene and CCl₄		
male rats fed a standard diet containing 0.02% 2-AAF for 2 weeks. CCl ₄ (2 ml kg ⁻¹ BW) fed by oral gavage on day 7 and killed 1–14 days after CCl ₄	classic zone-3 parenchymal necrosis with massive oval cell proliferation from portal tracts. Oval cells disappear after withdrawal of 2-AAF supplement	Ghoshal <i>et al.</i> (1983)
2-AAF in a choline devoid diet (CD/2-AAF)		
male rats fed CD diet supplemented with 0.05% 2-AAF for 12 days, returned to basal diet and killed up to 28 days after commencing the CD/2-AAF diet	essentially similar pattern of oval cell proliferation as for 2-AAF/PH regime. Albumin and AFP expression, but no evidence of complete hepatocyte differentiation	Sell (1983), Dunsford <i>et al.</i> (1985), Sell <i>et al.</i> (1981), Sell & Salman (1984)
2-acetylaminofluorene and partial hepatectomy(2-AAF/PH)		
oral gavage of male rats with 2-AAF for 5 days in first week, PH and feeding continued for further 4 days (average daily 2-AAF dose: 6.6 mg kg ⁻¹ BW). Rats killed up to 13 days after PH	no parenchymal necrosis. Gradual increase in number of oval cells after PH, with strong expression of albumin. [³ H]-thymidine initially incorporated exclusively in oval cells became 'transferred' to newly formed basophilic hepatocytes; i.e. the latter differentiated from the oval cells (the 'precursor-product relationship')	Evarts <i>et al.</i> (1987)
	oval cells express AFP and undergo intestinal metaplasia; metaplastic tissue did not express AFP	Bisgaard <i>et al.</i> (1994)
male rats fed a basal diet containing 0.02% 2-AAF for 2 weeks and PH administered at the midway point. Rats killed up to 9 weeks after PH	oval cell proliferation into parenchyma, with intestinal metaplasia common after 3 weeks. These metaplasias develop into cholangiofibrotic lesions. No hepatocyte differentiation	Tatematsu <i>et al.</i> (1985)
male rats fed 2-AAF (10 mg kg ⁻¹ BW) daily by oral gavage for 14 days, with PH performed at midway point. Rats killed up to 7 days after PH	massive oval cell proliferation with intestinal metaplasia by 1 week after PH. Oval cells express AFP and albumin, and differentiating CYP450-expressing hepatocytes develop from some of these ducts	Sarraf <i>et al.</i> (1994), Golding <i>et al.</i> (1995)

(continued)

Table 1. (*continued*)

regime	observations	reference
male rats fed 2-AAF at either 5 mg kg ⁻¹ or 2.5 mg kg ⁻¹ BW from 6 days prior to and up to 7 days after PH. Rats killed up to 14 days after PH	the higher of the two doses of 2-AAF produces a similar response as that observed using a 10 mg kg ⁻¹ dose (see above). The lower dose of 2-AAF gives a similar result up to day 7 after PH but without intestinal metaplasia, but by day 14 the majority of oval cells differentiate into small hepatocytes	Alison <i>et al.</i> (1996, 1997)
choline-devoid ethionine (CDE) diet male rats fed a standard choline-devoid diet supplemented with 0.05–0.1% ethionine for 1 day to 12 weeks	0.1% ethionine: proliferation of α -FP and of albumin-expressing, ductular oval cells which, unlike bile duct ligation, ANIT or DDPM treatment, do not commonly form well-defined ductular profiles. Hepatocyte differentiation very rare and no necrosis 0.07% ethionine: oval cells proliferated as cords of cells initially, ducts by 2 weeks and some formed intestinal-like cells by week 5. Oval cells expressed both foetal and adult hepatocyte enzymes 0.05% or 0.1% ethionine: parenchymal necrosis, massive oval cell proliferation and cholangio-fibrosis. Extent and onset of oval cell proliferation is greater and earlier at higher dose. No hepatocyte differentiation within ducts ultrastructurally, oval cells seen to migrate along sinusoids and also form intimate associations (desmosomal and tight junctions, canalicular formation, and interdigitation of microvilli) with neighbouring hepatocytes	Lenzi <i>et al.</i> (1992), Alpini <i>et al.</i> (1992) Tee <i>et al.</i> (1994) Tarsetti <i>et al.</i> (1993) Novikoff <i>et al.</i> (1991)
3'-methyl diaminobenzidine (3'-Me-DAB) male rats fed a standard diet containing 0.06% 3'-Me-DAB for up to 12 weeks	no parenchymal necrosis. Massive oval cell proliferation that penetrates parenchyma as ill-defined ductules. Cholangiofibrosis and intestinal-like differentiation after 12 weeks 1–2% of proliferating oval cells expressing AFP proliferation of oval cells accompanied by hyperplasia of small hepatocytes (derived from oval cells?) expressing the bile duct marker gamma glutamyl transpeptidase (GGT). By week 7, oval cell numbers decrease to be replaced by hyperplastic hepatocyte foci that express canalicular GGT	Carthew <i>et al.</i> (1989) Germain <i>et al.</i> (1985) Suzuki <i>et al.</i> (1987)
galactosamine male rats given a single intraperitoneal (i.p.) dose of galactosamine (700 mg kg ⁻¹ BW) and killed 1–8 days after	parenchymal necrosis and proliferation of small clusters and ductular oval cells, expressing AFP and albumin. Transitional hepatocytes differentiate from these cells and lose biliary specific markers	Dabeva & Shafritz (1993), Dabeva <i>et al.</i> (1993)
two i.p. doses of galactosamine (750 mg kg ⁻¹ BW), 6 h apart and killed 1–10 days after	as above with oval cells expressing foetal hepatocyte enzyme markers and normal bile ducts. Small hepatocytes and 'hepatocytes lining atypical duct structures' present	Lemire <i>et al.</i> (1991)
dipin a single i.p. injection of dipin (60 mg kg ⁻¹ BW) into mice followed 2 h later with a PH. Livers examined up to 18 months thereafter	initial massive parenchymal necrosis, oval cell proliferation from which small foci of hepatocytes are derived. Differentiating hepatocytes occasionally seen in oval cell ducts. Damaged parenchyma replaced by 'newly formed hepatocytes'	Factor <i>et al.</i> (1994)

(*continued*)

Table 1. (*continued*)

regime	observations	reference
furan male rats fed furan (15–60 mg kg ⁻¹ BW) by oral gavage, once a day, 5 days a week for up to 13 weeks	liver necrosis and replacement by 'hyperplastic bile ductular structures' with abundant fibrosis. 'Ductular hepatocytic cells' expressing dual bile duct/hepatocyte phenotypes differentiating within some ducts. Development of intestinal-like glands and preneoplastic hepatocyte foci and hepatocellular carcinomas at later time points	Elmore & Sirica (1991, 1992, 1993), Sirica <i>et al.</i> (1994)
chloroform inhalation male and female rats exposed to chloroform vapour (300 p.p.m.) for 6 h per day, 5–7 days per week, for up to 13 weeks	parenchymal necrosis and regeneration by hepatocytes with biliary duct hyperplasia confined to portal stroma. Intestinal metaplasia with fibrosis developed from 6 weeks onwards	Jamison <i>et al.</i> (1996)
carbon tetrachloride inhalation inhalation of CCl ₄ by mice, (no details of dose given)	classic centrilobular necrosis with AFP expressing oval cells that appear as small ducts or as small clusters of cells in the perinecrotic regions	Engelhardt <i>et al.</i> (1984)
diethylnitrosamine (DEN) three strains of male mice injected i.p. with 10–150 mg kg ⁻¹ BW DEN and examined 1–7 days thereafter	hepatic necrosis and oval cell proliferation. 'Transitional cells' (differentiating hepatocytes) observed within ducts. Different magnitudes of necrosis, oval cell proliferation and hepatocyte differentiation between strains	He <i>et al.</i> (1994)
long-term exposure to ethanol rats fed for 1–24 months on a totally liquid diet containing 5% ethanol	sparse numbers of very slowly proliferating oval cells expressing markers of foetal and mature hepatocytes. Similar changes as seen using the CDE protocol, yet over a much longer time frame; e.g. magnitude of response after 6–9 months on ethanol diet is similar to that after only 3 weeks on CDE regime	Smith <i>et al.</i> (1996)
allyl alcohol female rats injected with a single i.p. dose of allyl alcohol (0.62 mmol kg ⁻¹ BW) and killed after 6 h to 6 days	periportal necrosis induced within 6–12 h. Damaged parenchyma is reconstituted within 6 days by small hepatocyte-like cells (non-ductular oval cell progeny?) derived from 'intraportal stem cells'	Yavorkovsky <i>et al.</i> (1995)
diethylnitrosamine and <i>Clonorchis sinensis</i> Syrian hamsters infected with CS metacercariae and DMN (15 p.p.m.) in drinking water for 28 days	intense oval cell proliferation, followed by ductular dysplasia and development of cholangio-carcinoma	Lee <i>et al.</i> (1997)
p21^{CIP1/WAF1} transgenic mouse p21 overexpression targeted to the liver	retarded liver development with oval cell proliferation	Wu <i>et al.</i> (1996)
Long-Evans Cinnamon rat defect in Wilson disease gene (<i>ATP7B</i>) resulting in toxic accumulation of copper	acute hepatitis and death; survivors get chronic hepatitis accompanied by oval cell proliferation leading to development of hepatocellular carcinoma and cholangiocarcinoma	Betto <i>et al.</i> (1996), Yasui <i>et al.</i> (1997)

(figure 4*b,d*), invading deep into the hepatic parenchyma as tortuous arborizing ductules (Alpini *et al.* 1992; Sarraf *et al.* 1994), and expressing hepatocyte proteins (Golding *et al.* 1995), bile duct ligation induces bile stasis and also biliary cell hyperplasia, but with the important distinction that such hyperplastic ducts remain within the confines of the portal space (Milani *et al.* 1989; Alpini *et al.* 1992; Lenzi *et al.* 1992). In addition, these hyperplastic ducts do not differentiate into hepatocytes, neither do they express liver-enriched transcription factors that regulate AFP and albumin expression (Bisgaard *et al.* 1996), unlike

AAF-induced oval cells. Moreover, while bile stasis in adult liver also fails to induce biliary expression of traditional oval cell 'markers' such as AFP, SCF and c-kit, these molecules are upregulated following bile duct ligation of very young rats (Omori *et al.* 1997). Perhaps this indicates the greater bipotentiality of early postnatal biliary epithelia, but the fact that these molecules were particularly upregulated in the smaller ductules, reinforces the opinion that the small ducts are the prime instigators of oval cell reactions.

Figure 4 illustrates the characteristics of a typical oval cell reaction with its demonstrable biliary heritage (figure

4d); the oval cells also express vimentin. Co-expression of CKs and vimentin is seen in many carcinomas, a combination of intermediate filaments considered to endow epithelial cancer cells with a more invasive and possibly metastatic phenotype (Chu *et al.* 1996; Gilles *et al.* 1996; Hendrix *et al.* 1997). An increased migratory phenotype is, of course, a useful attribute for oval cells. Likewise, cultured neonatal hepatocytes express both CKs and vimentin coincident with the acquisition of a more fibroblast-like morphology (Pagan *et al.* 1995); EGF promoted this co-expression, whereas the differentiating agent, dimethyl sulphoxide, inhibited the increase in vimentin (Pagan *et al.* 1997).

Many studies have attempted to identify the first cells to proliferate at the onset of oval cell reactions, the rationale being that the initial 'proliferators' represent the stem cells. Cell proliferation may begin both in the cholangioles (Grisham & Porta 1964; Lenzi *et al.* 1992; Alison *et al.* 1996) and small marginal bile ducts (Evarts *et al.* 1993; Anilkumar *et al.* 1995), with concomitant expression of AFP and albumin (Golding *et al.* 1995). Some studies report that it is not the biliary epithelium that responds first, but the biliary-associated periductular cells (Sell 1990; Lenzi *et al.* 1992). These cells may give rise to oval cells themselves (Evarts *et al.* 1990), as well as components of the biliary stroma (Grisham & Porta 1964; Lenzi *et al.* 1992). Some periductular cells express AFP mRNA in normal liver (Alpini *et al.* 1992) and oval cell markers in carcinogen-treated livers (Evarts *et al.* 1990). They maybe Ito cells or myofibroblasts that co-proliferate with oval cells, and these have been distinguished from the latter by expression of desmin; on this basis their role as oval cell progenitors has been more or less discounted (Thorgeirsson *et al.* 1993). During foetal bile-duct development in humans, α -smooth muscle actin-positive Ito cells envelope the new ducts as they form from the ductal plate (Cocjin *et al.* 1996), emphasizing the close association between the two cell types.

Oval cell ductules eventually differentiate into small hepatocytes (figure 5a), losing biliary traits (CK7 and CK19 expression, figure 5b), and ceasing expression of vimentin and the oncofoetal glycoprotein, AFP.

7. STROMAL INFLUENCES ON REACTIVE BILE DUCTULES

Reactive ductules in human liver are surrounded by activated mesenchymal cells consisting primarily of myofibroblasts and perisinusoidal Ito cells (Burt & MacSween 1993). After hepatic injury in humans, Ito cells acquire a myofibroblast-like phenotype expressing desmin and α -smooth muscle actin (Schmitt-Graf *et al.* 1991; Arthur & Iredale 1994). Similarly activated Ito cells appear in rat livers damaged by carbon tetrachloride (Johnson *et al.* 1992), and in oval cell activated rat livers (Alison *et al.* 1993b; Sarraf *et al.* 1994; Anilkumar *et al.* 1995).

Ito cells are the first to proliferate in the AAF/PH model (Thorgeirsson *et al.* 1993), and are intimately associated with oval cells, enveloping them in a dense meshwork (Evarts *et al.* 1990). Ito cells are regarded as the principal source of extracellular matrix (ECM) proteins during hepatic regeneration both in rats (Johnson *et al.* 1992) and humans (Griffiths *et al.* 1992); additionally they

secrete metalloproteinases specific for the ECM, in particular for basement membrane proteins (Arthur *et al.* 1989). Secretion of matrix-busting enzymes may be critical for initiating oval cell invasion by 'clearing a path' through the damaged parenchyma to facilitate ductular proliferation, migration and morphogenesis, reactions promoted by Ito cell-produced HGF (see below). The oval cells themselves might also degrade matrix proteins, as primitive biliary cells migrating from the ductal plate can express matrix metalloproteinases during liver development in humans (Terada *et al.* 1995). However, ECM degradation during oval cell migration may not always be so pressing as oval cells frequently migrate along the space of Disse (Betto *et al.* 1996; Alison *et al.* 1997; and see figure 4b). Ito cells also synthesize laminin as they penetrate clusters of normal regenerating hepatocytes, which might provide a stimulus for the ingrowth of endothelial cells to re-establish the sinusoidal vasculature (Martinez-Hernandez & Amenta 1995). Therefore, Ito cells among reactive biliary epithelia might function to stimulate both oval cell migration during the early stages of regeneration, as well as restoring the normal lobular vasculature after hepatocyte differentiation. Ito cells may also insulate oval cells from the hepatocyte milieu during the early stages of cell migration, thereby preventing premature hepatocyte differentiation induced by the parenchymal microenvironment; cultured oval-like cells, when transplanted into the liver parenchyma of syngeneic rats swiftly integrate into the hepatic plates and acquire features of fully mature hepatocytes (Coleman *et al.* 1993; Grisham *et al.* 1993). The deterministic role of the hepatic microenvironment is further highlighted by the fact that it can even decrease the tumorigenicity of malignantly transformed oval-like cells (Coleman *et al.* 1993; Grisham *et al.* 1993). When these cells were transplanted subcutaneously they commonly formed highly aggressive, poorly differentiated tumours. On the other hand, those transplanted into the liver either lost the malignant phenotype or became more differentiated.

The close association of oval cells with stromal elements places them in a situation mimicking the biliary epithelium embedded in the portal mesenchyme. This may have a deterministic role in oval cell differentiation, retaining a biliary-like phenotype during the early stages of the regenerative response, which latterly breaks down, removing the 'biliary cell lineage restraint' and permitting differentiation into hepatocytes.

Cell-cell recognition and cell-ECM interactions are critical for cellular aggregation, segregation and migration. Contact with ECM components is fundamental in retaining cell polarity and regulating tissue-specific histogenesis and gene transcription, as well as maintaining tissue architecture (Rescan *et al.* 1993; Stamatoglou & Hughes 1994). Changes in matrix and intercellular adhesion very likely influence the pronounced tissue remodelling and alterations in gene expression that occur during oval cell responses. The influence of the ECM on gene transcription is highlighted by the well-known rapid hepatocyte retrodifferentiation to a more primitive phenotype when cultured under simple conditions, a process that can be counteracted by addition of ECM proteins (Stamatoglou & Hughes 1994). For example, albumin synthesis is maintained when hepatocytes contact type IV collagen,

but substitution of this protein with laminin results in the expression of a more primitive AFP-expressing phenotype, akin to limiting plate hepatocytes *in vivo* (Shah & Gerber 1990). The hepatic parenchyma is unique compared with other epithelia in that it does not rest on a true basement membrane, although the hepatocytes rest on a 'basement membrane-like matrix' consisting of collagen type IV, laminin, fibronectin and various proteoglycans (Stamatoglou & Hughes 1994). In addition to cell-matrix interactions, cell-cell associations are also likely to have an influence on the hepatocyte phenotype.

8. GROWTH FACTORS INVOLVED IN OVAL CELL ACTIVATION

The growth factors involved in oval cell reactions are essentially the same as those seemingly involved during normal hepatocyte regeneration. Nevertheless, the complexity of the remodelling process during ductular oval cell hyperplasia and differentiation clearly requires a sophisticated and highly coordinated temporal and spatial expression of both stimulatory and inhibitory growth factors, which in turn will influence cell-cell and cell-ECM interactions.

Parathyroid hormone-related peptide (PTHrP) is believed to be important in the growth and differentiation of several human tissues including liver. In humans, PTHrP is expressed in reactive bile ductules of cholestatic and regenerating liver (Roskams *et al.* 1993a), as well as by cholangiocarcinomas (Roskams *et al.* 1993b). The fact that growth factors such as EGF can rapidly induce PTHrP synthesis in cultured ductular cells has led to the belief that PTHrP is a member of the early response gene family, and that the peptide may have an autocrine role in bile ductular reactions (Roskams *et al.* 1995).

Oval cell proliferation and differentiation is mediated by complex growth factor loops that coordinate epithelial and stromal cell responses, highlighted by the expression of TGF β by reactive bile ductules, a growth factor known to stimulate perisinusoidal cells to produce ECM proteins (Milani *et al.* 1991). TGF β is also expressed by periductular Ito cells (Nagy *et al.* 1989; Evarts *et al.* 1990), and the abundance of this growth factor acting in an autocrine/paracrine fashion has been implicated in differentiation along the hepatocyte lineage, at least *in vitro* (Nagy *et al.* 1989). Ito cells also express TGF α , aFGF, HGF (Thorgeirsson *et al.* 1993) and stem cell factor (SCF) (Fujio *et al.* 1994). Oval cells express all of these growth factors apart from HGF (Evarts *et al.* 1992; Alison *et al.* 1993a) and the receptors for these ligands (Hu *et al.* 1993; Thorgeirsson *et al.* 1993; Fujio *et al.* 1994), and thus may be autonomous in their requirement for mitogens and morphogens. Indeed, cultured hepatoblasts—putative, embryological oval cell equivalents—are able to proliferate and migrate in serum-free conditions (Pagan *et al.* 1995), suggesting auto-crine regulation.

HGF was originally called 'scatter factor' (Stoker & Perryman 1985) because of its ability to promote the scattering and spreading of a variety of epithelial cells *in vitro*. Significantly, HGF is a powerful stimulator of human biliary epithelial cells *in vitro* (Joplin *et al.* 1992; Strain *et al.* 1995) and, furthermore, HGF is the most potent known mitogen for hepatocytes *in vitro* (Strain *et al.* 1991).

HGF is also a powerful morphogen, particularly during embryological tissue development (Barros *et al.* 1995), and therefore probably has a fundamental influence on oval cell behaviour (Alison *et al.* 1993b). The branching tubulogenesis of oval cells is undoubtedly augmented by HGF and TGF α , both of which have been shown to be critical in promoting and regulating ductular morphogenesis during epithelial tissue development (Barros *et al.* 1995), and may regulate this process via paracrine (Schirmacher *et al.* 1992; Hu *et al.* 1993) or autocrine (Burr *et al.* 1993) mechanisms. Indeed, HGF has been shown to induce glandular and ductular morphogenesis in a wide variety of epithelial cells growing in culture, including cell lines from colon, pancreas, mammary gland, prostate and lung (Brinkmann *et al.* 1995). In the AAF/PH model, aFGF mRNA expression by ductular oval cells coincides with their differentiation into hepatocytes (Marsden *et al.* 1992). In contrast, TGF β and PTHrP, which are also expressed during liver regeneration (Thorgeirsson *et al.* 1993) and by human reactive bile ductules (Roskams *et al.* 1993a), can inhibit ductal branching morphogenesis (Miettinen *et al.* 1994). HGF has, in addition, been shown to dissociate epithelial sheets to form mesenchymal-like cells, inducing a spindle cell-like morphology and increasing motility, migration and invasion, the latter phenomenon probably being accomplished by increased urokinase expression which mediates degradation of the ECM (Rosen *et al.* 1994).

HGF also plays a fundamental role in the control of proliferation and differentiation of erythroid progenitor cells in foetal liver, and acts in synergy with SCF (Galimi *et al.* 1994), an important effector of stem cell activation (Keshet *et al.* 1991). Both of these growth factors are expressed during oval cell proliferation (Thorgeirsson *et al.* 1993; Fujio *et al.* 1994), perhaps explaining why extramedullary haemopoiesis, probably from ordinarily resident pluripotential haemopoietic stem cells (Taniguchi *et al.* 1996), is such a common accompaniment to oval cell proliferation (Enomoto *et al.* 1978; Brill *et al.* 1993).

9. THE MULTIPOTENTIALITY OF OVAL CELLS

In rodents, oval cells are certainly not restricted to differentiation along the biliary and hepatocyte cell lineages, and appear additionally to have the lineage potential of uncommitted gastrointestinal stem cells. For instance, they can differentiate into intestinal absorptive cells (Tatematsu *et al.* 1985), goblet cells (Tatematsu *et al.* 1985; Golding *et al.* 1995) and endocrine and Paneth cells (Karaki *et al.* 1991; Elmore & Sirica 1993). In humans, endocrine differentiation can occur in normal bile ducts (Kurumaya *et al.* 1989), in ductular hyperplasia accompanying regeneration after submassive necrosis (Roskams *et al.* 1991) and in cholangiocarcinomas (Roskams *et al.* 1993a,b).

Pancreatic differentiation/metaplasia has been observed in liver associated with oval cell proliferation (Thorgeirsson *et al.* 1993), and the human peribiliary glands, which develop from the biliary ductal plate after birth, contain serous and mucus acini that can develop into ectopic exocrine pancreas (Terada & Nakanuma 1993). Conversely, hepatocyte (Rao *et al.* 1984, 1989) and goblet

cell differentiation (Zalatnai & Schally 1990) can occur in pancreatic ductular and/or periductular cells in rodents, and pancreatic oval cells in culture exhibit a ductular albumin-expressing phenotype (Ide *et al.* 1993). Similarly, cultured pancreatic ductal epithelial cells from rats can undergo hepatocytic differentiation when transplanted subcutaneously or intraperitoneally into syngeneic rats (Chen *et al.* 1995). Furthermore, pancreatic acinar cell carcinomas expressing a combination of hepatocyte, intestinal and neuroendocrine markers have been found in humans (Shinagawa *et al.* 1995), and hepatocyte differentiation can occur in adenocarcinomas of the renal pelvis (Ishikura *et al.* 1991) and stomach (Ishikura *et al.* 1986). The hepatoid stomach adenocarcinomas express a number of hepatocyte-specific markers, and in one case even bile secretion was evident (Ishikura *et al.* 1986). Interestingly, these tumours were always associated with intestinal metaplasia, a further example of deregulated gene expression.

This plasticity of cellular phenotypes is perhaps not surprising considering the liver, pancreas and intestine have a common embryological origin, and all three are connected via a continuous epithelium. Indeed, culture and transplantation of stem-like cells from normal liver has shown that these cells share epitopes with cells in the small ducts of the liver, pancreas, intestinal crypts and tracheobronchial epithelium (Grisham 1994). In fact, the default lineage commitment of the dorsal endoderm of the primitive gut (i.e. that which is not destined to become liver epithelium), appears to be the hepatic programme. AFP is expressed throughout the foetal gut villous epithelium (Tyner *et al.* 1990), but is later inhibited by the suppressive effects of adjacent endodermal components, which direct differentiation towards non-hepatic tissues of the gastrointestinal tract (Gualdi *et al.* 1996). It is interesting to note that transcription of the AFP gene is retained in a small number of enteroendocrine cells of the adult murine intestine (Tyner *et al.* 1990).

Under normal conditions, remarkable similarities exist between the organs of the digestive system. For instance, like the intestine, goblet cells and brush border cells are found in the larger ducts of the pancreas, as are endocrine cells, which unlike those of the intestinal crypt, consist of cell foci (the islets of Langerhans). During embryological development the islet precursors appear as individual cells 'embedded' within the ducts (Slack 1995), and during situations that provoke chronic inflammation, not unlike those seen accompanying oval cell proliferation, new islets are observed budding from ductules in conjunction with ductular hyperplasia and goblet cell metaplasia. After selective ablation of the pancreatic acinar component, the remaining ducts will rapidly regenerate to replenish the lost tissue. In fact, the regenerative powers of the pancreas appear to be as great as those of the liver in that even after 90% of the organ is removed, the animal survives with subsequent regeneration of both exocrine and endocrine components (Slack 1995); evidently like liver stem cells, pancreatic stem cells also reside within the ductular network.

Are there markers on oval cells which are indicative of lineage commitment? Differential expression of a combination of AFP and CK14 may be a useful indicator (Thorgeirsson *et al.* 1993). CK14 is not normally associated

with the liver (Moll *et al.* 1982), but has been detected in a small population of biliary-associated cells (Blouin *et al.* 1992). During the oval cell response using the AAF/PH model (Thorgeirsson *et al.* 1993), the newborn cells initially express CK14, and these cells give rise to two expanding oval cell populations consisting of cells expressing CK14 alone, or CK14 in combination with AFP. The latter phenotype appears to be restricted to differentiation along the hepatocyte lineage, while CK14 expression alone may be a marker of multipotential progenitor cells.

Oval cells clearly give rise to many different cell types, but it is uncertain as to whether there is a common multipotential stem cell, as in the bone marrow, which gives rise to committed unipotent stem cells, or if there are lineage-specific stem cells scattered about within the biliary tree.

10. IN VITRO STUDIES

Most cells isolated from normal or carcinogen-fed rat livers soon die in primary culture and cultures become dominated by rapidly growing colonies of small non-parenchymal epithelial cells traditionally called 'liver epithelial cells' (LECs) or 'rat liver epithelial' (RLE) cells (Tsao *et al.* 1984, 1985; Marceau *et al.* 1986; Blouin *et al.* 1992). Support for the existence of liver stem cells has received significant input from studies of cells derived from liver tissue from foetal and adult rats (Grisham 1980; Tsao *et al.* 1984; Grisham *et al.* 1993). These cells can share cholangiocyte and hepatocyte features, and closely resemble many of the cell lines that have been derived from oval cell-activated rodent livers (Hayner *et al.* 1984; Tsao *et al.* 1984; Marceau *et al.* 1986).

Other groups have propagated long-term cultures of small hepatocytes that generally do not express characteristics of biliary cells such as CK7 and CK19 expression (Mitaka *et al.* 1993; Tatenos & Yoshizato 1996). Cultured in the presence of nicotinamide and defined growth factor-containing medium, particularly EGF, these cells displayed appropriate hepatocyte traits: expression of CK8 and CK18, and production of albumin and transferrin. Such cells were considered to be committed progenitor cells rather than bipotential oval cells.

Cultured oval cells can certainly differentiate into hepatocytes when returned to their *in vivo* origins. Oval cells isolated from Long-Evans Cinnamon (LEC) rats will differentiate into albumin-producing hepatocytes when returned to the livers of LEC/Nagase analbuminemic, double-mutant rats (Yasui *et al.* 1997).

11. OVAL CELLS AND CANCER

Stem cells are important from the viewpoint of being the most likely targets for carcinogenic agents, because their longevity assures a continued presence during the long latency between exposure and cancer development. However the role of stem/oval cells in the histogenesis of primary tumours in the liver is still unresolved, and tumours could arise from the 'maturation arrest' of the differentiating stem cell progeny (Marceau 1990; Sell 1993a,b; Sell & Pierce 1994), or by dedifferentiation of mature liver cells. This question is still unresolved and will only be briefly covered here, but has been substantially

reviewed elsewhere (Sell & Dunsford 1989; Aterman 1992; Sell 1993b).

Of relevance here is the large body of *in vitro* experimental evidence supporting the stem cell 'maturation arrest' hypothesis, in that cultured, malignantly transformed oval cells are able to give rise to tumours with a whole range of phenotypes, highly suggestive of the multipotential nature of the target cells (Tsao *et al.* 1985; Tsao & Grisham 1987; Garfield *et al.* 1988; Steinberg *et al.* 1994). Similarly, it is possible to use extrinsic agents to switch a human cholangiocarcinoma cell line to one that has hepatocytic features (Enjoji *et al.* 1997), presumably reflecting the likely histogenesis of these two tumour types from a common precursor cell. In a similar vein, transplantation of transformed oval cells into syngeneic recipients can result in either hepatoblastomas, hepatocellular-carcinomas, cholangiocarcinomas, cholangiocarcinomas with intestinal features (Steinberg *et al.* 1994), intestinal adenocarcinomas (Marceau 1990), anaplastic tumours (Steinberg *et al.* 1994) and even sarcomas (Marceau 1990). Furthermore, the lineage commitment of oval cells can be altered by a single transforming oncogene, and the range of tumours derived from oval cells may reflect mutations in different combinations of cellular proto-oncogenes (Garfield *et al.* 1988).

12. CONCLUSIONS

Reactive biliary epithelium, usually organized into ductules and collectively known as oval cells, appear to be derived from facultative stem cells located in the cholangioles and small interlobular bile ducts. Both *in vitro* and *in vivo* studies have clearly demonstrated that oval cells have the potential to differentiate along many divergent pathways, and furthermore, after transformation and transplantation, they can similarly give rise to a range of tumour types, although their role in human hepatocarcinogenesis is unclear. Oval cell proliferation and differentiation can be readily induced in experimental animals, although the controlling mechanisms for these processes are poorly understood.

The selective harvesting and culture of both rodent (Alpini *et al.* 1997) and biopsied human (Strain *et al.* 1995) biliary epithelial cells is now possible, hopefully paving the way for *ex vivo* gene therapy of liver stem cell progeny for the correction of metabolic liver disorders. In addition, selective *in vivo* gene transfer to the cholangiocytes of temporarily ligated, rodent bile ducts has been performed (Cabrera *et al.* 1996), raising the potential for the application of a similar strategy to the human setting for the amelioration of the defective Cl⁻ and water secretion characteristic of cystic fibrosis.

REFERENCES

- Alison, M. R. 1986 Regulation of liver growth. *Physiol. Rev.* **66**, 499–541.
- Alison, M. R., Nasim, M., Anilkumar, T. V. & Sarraf, C. E. 1993a Transforming growth factor- α immunoreactivity in a variety of epithelial tissues. *Cell Prolif.* **26**, 449–460.
- Alison, M. R., Poulosom, R., Jeffery, R., Anilkumar, T. V., Jagoe, R. & Sarraf, C. E. 1993b Expression of hepatocyte growth factor mRNA during oval cell activation in the rat liver. *J. Pathol.* **171**, 291–299.
- Alison, M. R., Golding, M., Sarraf, C. E., Edwards, R. J. & Lalani, E. N. 1996 Liver damage in the rat induces hepatocyte stem cells from biliary epithelial cells. *Gastroenterology* **110**, 1182–1190.
- Alison, M., Golding, M., Lalani, E. N., Nagy, P., Thorgeirsson, S. & Sarraf, C. 1997 Wholesale hepatocytic differentiation in the rat from ductular oval cells, the progeny of biliary stem cells. *J. Hepatol.* **26**, 343–352.
- Alpini, G., Aragona, E., Dabeva, M., Salvi, R., Shafritz, D. A. & Tavoloni, N. 1992 Distribution of albumin and alpha-fetoprotein mRNAs in normal, hyperplastic, and preneoplastic rat liver. *Am. J. Pathol.* **141**, 623–632.
- Alpini, G., Roberts, S., Kuntz, S. M., Ueno, Y., Gubba, S., Podila, P. V., LeSage, G. & LaRusso, N. F. 1996 Morphological, molecular, and functional heterogeneity of cholangiocytes from normal rat liver. *Gastroenterology* **110**, 1636–1643.
- Alpini, G., Glaser, S., Robertson, W., Rodgers, R. E. D., Phinizz, J. L., Lasater, J. & Lesage, G. D. 1997 Large but not small intrahepatic bile ducts are involved in secretin-regulated ductal bile secretion. *Am. J. Physiol.* **272**, G1064–G1074.
- Anilkumar, T. V., Golding, M., Edwards, R. J., Lalani, E.-N., Sarraf, C. E. & Alison, M. R. 1995 The resistant hepatocyte model of carcinogenesis in the rat: the apparent independent development of oval cell proliferation and early nodules. *Carcinogenesis* **16**, 845–853.
- Arthur, M. J. & Iredale, J. P. 1994 Hepatic lipocytes, TIMP-1 and liver fibrosis. *J. R. Coll. Physicians, Lond.* **28**, 200–208.
- Arthur, M. J., Friedman, S. L., Roll, F. J. & Bissell, D. M. 1989 Lipocytes from normal rat liver release a neutral metalloproteinase that degrades basement membrane (type IV) collagen. *J. Clin. Invest.* **84**, 1076–1085.
- Aterman, K. 1992 The stem cells of the liver—a selective review [editorial]. *J. Cancer Res. Clin. Oncol.* **118**, 87–115.
- Barros, E. J., Santos, O. F., Matsumoto, K., Nakamura, T. & Nigam, S. K. 1995 Differential tubulogenic and branching morphogenetic activities of growth factors: implications for epithelial tissue development. *Proc. Natn. Acad. Sci. USA* **92**, 4412–4416.
- Betto, H., Kaneda, K., Yamamoto, T., Kojima, A. & Sakurai, M. 1996 Development of intralobular bile ductules after spontaneous hepatitis in Long-Evans mutant rats. *Lab. Invest.* **75**, 43–53.
- Bhathal, P. S. & Christie, G. S. 1969 A fluorescence microscopic study of bile duct proliferation induced in guinea pigs by alpha-naphthyl isothiocyanate. *Lab. Invest.* **20**, 480–487.
- Bisgaard, H. C., Nagy, P., Ton, P. T., Hu, Z. & Thorgeirsson, S. S. 1994 Modulation of keratin 14 and alpha-fetoprotein expression during hepatic oval cell proliferation and liver regeneration. *J. Cell Physiol.* **159**, 475–484.
- Bisgaard, H. C., Nagy, P., Santonirugiu, E. & Thorgeirsson, S. S. 1996 Proliferation, apoptosis, and induction of hepatic transcription factors are characteristics of the early response of biliary epithelial (oval) cells to chemical carcinogens. *Hepatology* **23**, 62–70.
- Blouin, R., Blouin, M. J., Royal, I., Grenier, A., Roop, D. R., Loranger, A. & Marceau, N. 1992 Cytokeratin 14 expression in rat liver cells in culture and localization in vivo. *Differentiation* **52**, 45–54.
- Blouin, M. J., Lamy, I., Loranger, A., Noel, M., Corlu, A., Guguen-Guillouzo, C. & Marceau, N. 1995 Specialisation switch in differentiating embryonic rat liver progenitor cells in response to sodium butyrate. *Exp. Cell Res.* **217**, 22–30.
- Bralet, M. P., Branchereau, S., Brechot, C. & Ferry, N. 1994 Cell lineage study in the liver using retroviral mediated gene transfer. Evidence against the streaming of hepatocytes in normal liver. *Am. J. Pathol.* **144**, 896–905.
- Brill, S., Holst, P., Sigal, S., Zvibel, I., Fiorino, A., Ochs, A., Somasundaran, U. & Reid, L. M. 1993 Hepatic progenitor populations in embryonic, neonatal, and adult liver. *Proc. Soc. Exp. Biol. Med.* **204**, 261–269.

- Brinkmann, V., Foroutan, H., Sachs, M., Weidner, K. & Birchmeier, W. 1995 Hepatocyte growth factor/scatter factor induces a variety of tissue specific morphogenic programs in epithelial cells. *J. Cell Biol.* **131**, 1573–1586.
- Burr, A. W., Carpenter, M. R., Hines, J. E., Gullick, W. J. & Burt, A. D. 1993 Intrahepatic distribution of transforming growth factor- α (TGF α) during liver regeneration following carbon tetrachloride-induced necrosis. *J. Pathol.* **170**, 95–100.
- Burt, A. D. & MacSween, R. N. 1993 Bile duct proliferation—its true significance? *Histopathology* **23**, 599–602.
- Cabrera, J. A., Wilson, J. M. & Raper, S. E. 1996 Targeted retroviral gene transfer into the rat biliary tract. *Somatic Cell Molec. Genet.* **22**, 21–29.
- Calnek, D. & Quaroni, A. 1993 Differential localization by in situ hybridization of distinct keratin mRNA species during intestinal epithelial cell development and differentiation. *Differentiation* **53**, 95–104.
- Carthew, P., Edwards, R. E., Hill, R. J. & Evans, J. G. 1989 Cytokeratin expression in cells of the rodent bile duct developing under normal and pathological conditions. *Br. J. Exp. Pathol.* **70**, 717–725.
- Chen, J.-R., Tsao, M.-S. & Duguid, W. P. 1995 Hepatocyte differentiation of cultured rat pancreatic ductal epithelial cells after *in vivo* implantation. *Am. J. Pathol.* **147**, 707–717.
- Chu, Y.-W., Seftor, E., Romer, L. H. & Hendrix, M. J. C. 1996 Experimental coexpression of vimentin and keratin intermediate filaments in human melanoma cells augments motility. *Am. J. Pathol.* **148**, 63–69.
- Cocjin, J., Rosenthal, P., Buslon, V., Luk, L., Barajas, L., Geller, S. A., Ruebner, B. & French, S. 1996 Bile ductule formation in fetal, neonatal, and infant livers compared with extrahepatic biliary atresia. *Hepatology* **24**, 568–574.
- Coleman, W. B., Wennerberg, A. E., Smith, G. J. & Grisham, J. W. 1993 Regulation of the differentiation of diploid and some aneuploid rat liver epithelial (stemlike) cells by the hepatic microenvironment. *Am. J. Pathol.* **142**, 1373–1382.
- Correspondence 1995 Do livers 'stream'? *Am. J. Pathol.* **146**, 772–776.
- Dabeva, M. D. & Shafritz, D. A. 1993 Activation, proliferation, and differentiation of progenitor cells into hepatocytes in the D-galactosamine model of liver regeneration. *Am. J. Pathol.* **143**, 1606–1620.
- Dabeva, M. D., Alpini, G., Hurston, E. & Shafritz, D. A. 1993 Models for hepatic progenitor cell activation. *Proc. Soc. Exp. Biol. Med.* **204**, 242–252.
- De-Vos, R. & Desmet, V. 1992 Ultrastructural characteristics of novel epithelial cell types identified in human pathologic liver specimens with chronic ductular reaction. *Am. J. Pathol.* **140**, 1441–1450.
- Delladetsima, J. K., Kyriakou, V., Vafiadis, I., Karakitsos, P., Smyrnof, T. & Tassopoulos, N. C. 1995 Ductular structures in acute hepatitis. *J. Pathol.* **175**, 69–76.
- Dunsford, H. A., Maset, R., Salman, J. & Sell, S. 1985 Connection of ductlike structures induced by a chemical hepatocarcinogen to portal bile ducts in the rat liver detected by injection of bile ducts with a pigmented barium gelatin medium. *Am. J. Pathol.* **118**, 218–224.
- Elmore, L. W. & Sirica, A. E. 1991 Phenotypic characterization of metaplastic intestinal glands and ductular hepatocytes in cholangiofibrotic lesions rapidly induced in the caudate liver lobe of rats treated with furan. *Cancer Res.* **51**, 5752–5759.
- Elmore, L. W. & Sirica, A. E. 1992 Sequential appearance of intestinal mucosal cell types in the right and caudate liver lobes of furan-treated rats. *Hepatology* **16**, 1220–1226.
- Elmore, L. W. & Sirica, A. E. 1993 'Intestinal-type' of adenocarcinoma preferentially induced in right/caudate liver lobes of rats treated with furan. *Cancer Res.* **53**, 254–259.
- Engelhardt, N. V., Baranov, V. N., Lazareva, M. N. & Goussev, A. I. 1984 Ultrastructural localisation of alpha-fetoprotein (AFP) in regenerating mouse liver poisoned with CCL4. 1. Re-expression of AFP in differentiated hepatocytes. *Histochemistry* **80**, 401–407.
- Engelhardt, N. V., Factor, V. M., Yasova, A. K., Poltoranina, V. S., Baranov, V. N. & Lasareva, M. N. 1990 Common antigens of mouse oval and biliary epithelial cells. Expression on newly formed hepatocytes. *Differentiation* **45**, 29–37.
- Enjoji, M., Nakashima, M., Honda, M., Sakai, H. & Nawata, H. 1997 Hepatocytic phenotypes induced in sarcomatous cholangiocarcinoma cells treated with 5-azacytidine. *Hepatology* **26**, 288–294.
- Enomoto, K., Dempo, K., Mori, M. & Onoe, T. 1978 Histopathological and ultrastructural study on extramedullary hematopoietic foci in early stage of 3'-methyl-4-(dimethylamino) azobenzene hepatocarcinogenesis. *Gann* **69**, 249–254.
- Evarts, R. P., Nagy, P., Marsden, E. & Thorgeirsson, S. S. 1987 A precursor-product relationship exists between oval cells and hepatocytes in rat liver. *Carcinogenesis* **8**, 1737–1740.
- Evarts, R. P., Nagy, P., Nakatsukasa, H., Marsden, E. & Thorgeirsson, S. S. 1989 *In vivo* differentiation of rat liver oval cells into hepatocytes. *Cancer Res.* **49**, 1541–1547.
- Evarts, R. P., Nakatsukasa, H., Marsden, E. R., Hsia, C. C., Dunsford, H. A. & Thorgeirsson, S. S. 1990 Cellular and molecular changes in the early stages of chemical hepatocarcinogenesis in the rat. *Cancer Res.* **50**, 3439–3444.
- Evarts, R. P., Nakatsukasa, H., Marsden, E. R., Hu, Z. & Thorgeirsson, S. S. 1992 Expression of transforming growth factor- α in regenerating liver and during hepatic differentiation. *Molec. Carcinog.* **5**, 25–31.
- Evarts, R. P., Hu, Z., Fujio, K., Marsden, E. R. & Thorgeirsson, S. S. 1993 Activation of hepatic stem cell compartment in the rat: role of transforming growth factor α , hepatocyte growth factor, and acidic fibroblast growth factor in early proliferation. *Cell Growth Diff.* **4**, 555–561.
- Fabrikant, J. I. 1968 The kinetics of cellular proliferation in regenerating liver. *J. Cell Biol.* **36**, 551–565.
- Factor, V. M., Radaeva, S. A. & Thorgeirsson, S. S. 1994 Origin and fate of oval cells in dipin-induced hepatocarcinogenesis in the mouse. *Am. J. Pathol.* **145**, 409–422.
- Farber, E. 1955 Similarities in the sequences of early histological changes induced in the liver of the rat by ethionine, 2-acetylaminofluorene, and 3-methyl-4-dimethylaminoazobenzene. *Cancer Res.* **16**, 142–148.
- Fausto, N., Lemire, J. M. & Shiojiri, N. 1993 Cell lineages in hepatic development and the identification of progenitor cells in normal and injured liver. *Proc. Soc. Exp. Biol. Med.* **204**, 237–241.
- Flint, N., Pemberton, P. W., Loble, R. W. & Evans, G. S. 1994 Cytokeratin expression in epithelial cells isolated from the crypt and villus regions of the rodent small intestine. *Epithelial Cell Biol.* **3**, 16–23.
- Fujio, K., Evarts, R. P., Hu, Z., Marsden, E. R. & Thorgeirsson, S. S. 1994 Expression of stem cell factor and its receptor, c-kit, during liver regeneration from putative stem cells in adult rat. *Lab. Invest.* **70**, 511–516.
- Galimi, F., Bagnara, G., Bonsi, L., Cottone, E., Follenzi, A., Simeone, A. & Comoglio, P. 1994 Hepatocyte growth factor induces proliferation and differentiation of multipotent and erythroid hemopoietic progenitors. *J. Cell Biol.* **127**, 1743–1754.
- Garfield, S., Huber, B. E., Nagy, P., Cordingley, M. G. & Thorgeirsson, S. S. 1988 Neoplastic transformation and lineage switching of rat liver epithelial cells by retrovirus-associated oncogenes. *Molec. Carcinog.* **1**, 189–195.
- Gerlach, C., Sakkab, D. Y., Scholzen, T., Dassler, R., Alison, M. R. & Gerdes, J. 1997 Ki-67 expression during rat liver regeneration after partial hepatectomy. *Hepatology* **26**, 573–578.

- Germain, L., Goyette, R. & Marceau, N. 1985 Differential cytokeratin and alpha-fetoprotein expression in morphologically distinct epithelial cells emerging at the early stage of rat hepatocarcinogenesis. *Cancer Res.* **45**, 673–681.
- Germain, L., Blouin, M. J. & Marceau, N. 1988a Biliary epithelial and hepatocytic cell lineage relationships in embryonic rat liver as determined by the differential expression of cytokeratins, alpha-fetoprotein, albumin, and cell surface-exposed components. *Cancer Res.* **48**, 4909–4918.
- Germain, L., Noel, M., Gourdeau, H. & Marceau, N. 1988b Promotion of growth and differentiation of rat ductular oval cells in primary culture. *Cancer Res.* **48**, 368–378.
- Ghoshal, A. K., Mullen, B., Medline, A. & Farber, E. 1983 Sequential analysis of hepatic carcinogenesis. Regeneration of liver after carbon tetrachloride-induced liver necrosis when hepatocyte proliferation is inhibited by 2-acetylaminofluorene. *Lab. Invest.* **48**, 224–230.
- Gilles, C., Polette, M., Piette, J., Delvigne, A. C., Thompson, E. W., Foidart, J. M. & Birembaut, P. 1996 Vimentin expression in cervical carcinomas: association with invasive and migratory potential. *J. Pathol.* **180**, 175–180.
- Golding, M., Sarraf, C. E., Lalani, E.-N., Anilkumar, T. V., Edwards, R. J., Nagy, P., Thorgeirsson, S. S. & Alison, M. R. 1995 Oval cell differentiation into hepatocytes in the acetylaminofluorene-treated regenerating rat liver. *Hepatology* **22**, 1243–1253.
- Griffiths, M. R., Shepherd, M., Ferrier, R., Schuppan, D., James, O. F. W. & Burt, A. D. 1992 Light microscopic and ultrastructural distribution of type VI collagen in human liver: alterations in chronic biliary disease. *Histopathology* **21**, 335–344.
- Grisham, J. W. 1962 A morphologic study of deoxyribonucleic acid synthesis and cell proliferation in regenerating rat liver; autoradiography with thymidine H-3. *Cancer Res.* **22**, 842–849.
- Grisham, J. W. 1980 Cell types in long-term propagable cultures of rat liver. *Ann. NY Acad. Sci.* **349**, 128–137.
- Grisham, J. W. 1994 Migration of hepatocytes along hepatic plates and stem cell-fed hepatocyte lineages. *Am. J. Pathol.* **144**, 849–854.
- Grisham, J. W. & Porta, E. A. 1964 Origin and fate of proliferated hepatic ductal cells in the rat: electron-microscopic and autoradiographic studies. *Exptl Molec. Pathol.* **2**, 242–261.
- Grisham, J. W., Coleman, W. B. & Smith, G. J. 1993 Isolation, culture, and transplantation of rat hepatocytic precursor (stem-like) cells. *Proc. Soc. Exptl Biol. Med.* **204**, 270–279.
- Gualdi, R., Bossard, P., Zheng, M., Hamada, Y., Coleman, J. R. & Zaret, K. S. 1996 Hepatic specification of the gut endoderm in vitro: cell signaling and transcriptional control. *Genes Dev.* **10**, 1670–1682.
- Haque, S., Haruna, Y., Saito, K., Nalesnik, M. A., Atillasoy, E., Thung, S. N. & Gerber, M. A. 1996 Identification of bipotential progenitor cells in human liver regeneration. *Lab. Invest.* **75**, 699–705.
- Haruna, Y., Saito, K., Spaulding, S., Nalesnik, M. A. & Gerber, M. A. 1996 Identification of bipotential progenitor cells in human liver development. *Hepatology* **23**, 476–481.
- Hayner, N. T., Braun, L., Yaswen, P., Brooks, M. & Fausto, N. 1984 Isozyme profiles of oval cells, parenchymal cells, and biliary cells isolated by centrifugal elutriation from normal and preneoplastic livers. *Cancer Res.* **44**, 332–338.
- He, X. Y., Smith, G. J., Enno, A. & Nicholson, R. C. 1994 Short-term diethylnitrosamine-induced oval cell responses in three strains of mice. *Pathology* **26**, 154–160.
- Hendrix, M. J. C., Seftor, E. A., Seftor, R. E. B. & Trevor, K. T. 1997 Experimental co-expression of vimentin and keratin intermediate filaments in human breast cancer cells results in phenotypic interconversion and increased invasive behavior. *Am. J. Pathol.* **150**, 483–495.
- Hiruma, M., Seki, A., Ono, A. & Kurihara, T. 1993 Early hepatic lesions with marked glandular structures induced in rats by 0.1% ethionine in a choline deficient diet. *Jikken Dobutsu* **42**, 197–201.
- Hofer, D. & Drenckhahn, D. 1996 Cytoskeletal markers allowing discrimination between brush cells and other epithelial cells of the gut including enteroendocrine cells. *Histochem. Cell Biol.* **105**, 405–412.
- Hsia, C. C., Evarts, R. P., Nakatsukasa, H., Marsden, E. R. & Thorgeirsson, S. S. 1992 Occurrence of oval-type cells in hepatitis B virus-associated human hepatocarcinogenesis. *Hepatology* **16**, 1327–1333.
- Hu, Z., Evarts, R. P., Fujio, K., Marsden, E. R. & Thorgeirsson, S. S. 1993 Expression of hepatocyte growth factor and c-met genes during hepatic differentiation and liver development in the rat. *Am. J. Pathol.* **142**, 1823–1830.
- Huitfeldt, H. S., Brandtzaeg, P. & Poirier, M. C. 1990 Reduced DNA adduct formation in replicating liver cells during continuous feeding of a chemical carcinogen. *Proc. Natn. Acad. Sci. USA* **87**, 5955–5958.
- Huitfeldt, H. S., Brandtzaeg, P. & Poirier, M. C. 1991 Carcinogen-induced alterations in rat liver DNA adduct formation determined by computerized fluorescent image analysis. *Lab. Invest.* **64**, 207–214.
- Ide, H., Subbarao, V., Reddy, J. K. & Rao, M. S. 1993 Formation of ductular structures *in vitro* by rat pancreatic epithelial oval cells. *Exptl Cell Res.* **209**, 38–44.
- Ishikura, H., Kirimoto, K., Shamoto, M., Miyamoto, Y., Yamagiwa, H., Itoh, T. & Aizawa, M. 1986 Hepatoid adenocarcinomas of the stomach. An analysis of seven cases. *Cancer* **58**, 119–126.
- Ishikura, H., Ishiguro, T., Enatsu, C., Fujii, H., Kakuta, Y., Kanda, M. & Yoshiki, T. 1991 Hepatoid adenocarcinoma of the renal pelvis producing alpha-fetoprotein of hepatic type and bile pigment. *Cancer* **67**, 3051–3056.
- Jamison, K. C., Larson, J. L., Butterworth, B. E., Harden, R., Skinner, B. L. & Wolf, D. C. 1996 A non-bile duct origin for intestinal crypt-like ducts with periductular fibrosis induced in livers of F344 rats by chloroform inhalation. *Carcinogenesis* **17**, 675–682.
- Johnson, S. J., Hines, J. E. & Burt, A. D. 1992 Macrophage and perisinusoidal cell kinetics in acute liver injury. *J. Pathol.* **166**, 351–358.
- Joplin, R., Hishida, T., Tsubouchi, H., Daikuhara, Y., Ayres, R., Neuberger, J. M. & Strain, A. J. 1992 Human intrahepatic biliary epithelial cells proliferate *in vitro* in response to human hepatocyte growth factor. *J. Clin. Invest.* **90**, 1284–1289.
- Karaki, Y., Munakata, S., Saeki, T., Hirota, S. & Fujimaki, M. 1991 Appearance of a carcinoid-like pattern in rat hepatic tumors induced by 3'-methyl-4-dimethyl-aminoazobenzene. *Jap. J. Cancer Res.* **82**, 397–402.
- Kennedy, S., Rettinger, S., Flye, M. W. & Parker-Ponder, K. 1995 Experiments in transgenic mice show that hepatocytes are the source for postnatal liver growth and do not stream. *Hepatology* **22**, 160–168.
- Keshet, E., Lyman, S. D., Williams, D. E., Anderson, D. M., Jenkins, N. A., Copeland, N. G. & Parada, L. F. 1991 Embryonic RNA expression patterns of the c-kit receptor and its cognate ligand suggests multiple functional roles in mouse development. *EMBO J.* **10**, 2425–2435.
- Koukoulis, G., Rayner, A., Tan, K. C., Williams, R. & Portmann, B. 1992 Immunolocalization of regenerating cells after submassive liver necrosis using PCNA staining. *J. Pathol.* **166**, 359–368.
- Kroese, E. D., Bannenberg, G., Dogterom, P., Noach, A. B., Nagelkerke, J. F. & Meerman, J. H. 1990 Lipid peroxidation and protein thiol depletion are not involved in the cytotoxicity of N-hydroxy-2-acetylaminofluorene in isolated rat hepatocytes. *Biochem. Pharmacol.* **40**, 1885–1892.

- Kurumaya, H., Ohta, G. & Nakanuma, Y. 1989 Endocrine cells in the intrahepatic biliary tree in normal livers and hepatolithiasis. *Arch. Pathol. Lab. Med.* **113**, 143–147.
- Ledda, G. M., Sells, M. A., Yokoyama, S. & Lombardi, B. 1983 Metabolic properties of isolated rat liver cell preparations enriched in epithelial cells other than hepatocytes. *Int. J. Cancer* **31**, 231–237.
- Lee, J. H., Rim, H. J. & Sell, S. 1997 Heterogeneity of the oval-cell response in the hamster liver during cholangiocarcinogenesis following *Clonorchis sinensis* infection and dimethylnitrosamine treatment. *J. Hepatol.* **26**, 1313–1323.
- Lemire, J. M. & Fausto, N. 1991 Multiple alpha-fetoprotein RNAs in adult rat liver: cell type-specific expression and differential regulation. *Cancer Res.* **51**, 4656–4664.
- Lemire, J. M., Shiojiri, N. & Fausto, N. 1991 Oval cell proliferation and the origin of small hepatocytes in liver injury induced by D-galactosamine. *Am. J. Pathol.* **139**, 535–552.
- Lenzi, R., Liu, M. H., Tarsetti, F., Slott, P. A., Alpini, G., Zhai, W. R., Paronetto, F., Lenzen, R. & Tavoloni, N. 1992 Histogenesis of bile duct-like cells proliferating during ethionine hepatocarcinogenesis. Evidence for a biliary epithelial nature of oval cells. *Lab. Invest.* **66**, 390–402.
- Makino, Y., Yamamoto, K. & Tsuji, T. 1988 Three-dimensional arrangement of ductular structures formed by oval cells during hepatocarcinogenesis. *Acta Med. Okayama* **42**, 143–150.
- Marceau, N. 1990 Cell lineages and differentiation programs in epidermal, urothelial and hepatic tissues and their neoplasms. *Lab. Invest.* **63**, 4–20.
- Marceau, N. 1994 Epithelial cell lineages in developing, restoring, and transforming liver: evidence for the existence of a 'differentiation window'. *Gut* **35**, 294–296.
- Marceau, N., Germain, L., Goyette, R., Noel, M. & Gourdeau, H. 1986 Cell of origin of distinct cultured rat liver epithelial cells, as typed by cytokeratin and surface component selective expression. *Biochem. Cell Biol.* **64**, 788–802.
- Marsden, E. R., Hu, Z., Fujio, K., Nakatsukasa, H., Thorgeirsson, S. S. & Evarts, R. P. 1992 Expression of acidic fibroblast growth factor in regenerating liver and during hepatic differentiation. *Lab. Invest.* **67**, 427–433.
- Martinez-Hernandez, A. & Amenta, P. 1995 The extracellular matrix in hepatic regeneration. *FASEB J.* **9**, 1401–1410.
- Mathis, G. A., Walls, S. A., D'Amico, P., Gengo, T. F. & Sirica, A. E. 1989 Enzyme profile of rat bile ductular epithelial cells in reference to the resistance phenotype in hepatocarcinogenesis. *Hepatology* **9**, 477–485.
- Meybehm, M., Fischer, H. P. & Pfeifer, U. 1993 Expression of HBs- and HBc-antigen in neoductular epithelium in chronic active hepatitis B. A further support for hepato-ductular metaplasia. *Virchows Arch. B Cell Pathol.* **63**, 167–172.
- Miettinen, P. J., Ebner, R., Lopez, A. R. & Derynck, R. 1994 TGF-beta induced transdifferentiation of mammary epithelial cells to mesenchymal cells: involvement of type I receptors. *J. Cell Biol.* **127**, 2021–2036.
- Milani, S., Herbst, H., Schuppan, D., Niedobitek, G., Kim, K. Y. & Stein, H. 1989 Vimentin expression of newly formed rat bile duct epithelial cells in secondary biliary fibrosis. *Virchows Arch. A Pathol. Anat. Histopathol.* **415**, 237–242.
- Milani, S., Herbst, H., Schuppan, D., Stein, H. & Surrenti, C. 1991 Transforming growth factors beta 1 and beta 2 are differentially expressed in fibrotic liver disease. *Am. J. Pathol.* **139**, 1221–1229.
- Millward-Sadler, G. H. & Jezequel, A.-M. 1992 Normal histology and ultrastructure. In *Wright's liver and biliary disease*, vol. 1 (ed. G. H. Millward-Sadler, R. Wright & M. J. P. Arthur), pp. 12–42. London: W. B. Saunders Co. Ltd.
- Mitaka, T., Norioka, K., Nakamura, T. & Mochizuki, Y. 1993 Effects of mitogens and co-mitogens on the formation of small-cell colonies in primary cultures of rat hepatocytes. *J. Cell Physiol.* **157**, 461–468.
- Moll, R., Franke, W. W., Schiller, D. L., Geiger, B. & Krepler, R. 1982 The catalog of human cytokeratins: patterns of expression in normal epithelia, tumors and cultured cells. *Cell* **31**, 11–24.
- Nagy, P., Evarts, R. P., McMahon, J. B. & Thorgeirsson, S. S. 1989 Role of TGF-beta in normal differentiation and oncogenesis in rat liver. *Molec. Carcinog.* **2**, 345–354.
- Nagy, P., Bisgaard, H. C. & Thorgeirsson, S. S. 1994 Expression of hepatic transcription factors during liver development and oval cell differentiation. *J. Cell Biol.* **126**, 223–233.
- Nomoto, M., Uchikosi, Y., Kajikazawa, N., Tanaka, Y. & Asakura, H. 1992 Appearance of hepatocytelike cells in the interlobular bile ducts of human liver in various liver disease states. *Hepatology* **16**, 1199–1205.
- Novikoff, P. M., Ikeda, T., Hixson, D. C. & Yam, A. 1991 Characterizations of and interactions between bile ductule cells and hepatocytes in early stages of rat hepatocarcinogenesis induced by ethionine. *Am. J. Pathol.* **139**, 1351–1368.
- Novikoff, P. M., Yam, A. & Oikawa, I. 1996 Blast-like cell compartment in carcinogen-induced proliferating bile ductules. *Am. J. Pathol.* **148**, 1473–1492.
- Omori, M., Evarts, R. P., Omori, N., Hu, Z., Marsden, E. R. & Thorgeirsson, S. S. 1997 Expression of alpha-fetoprotein and stem cell factor/c-kit system in bile duct ligated young rats. *Hepatology* **25**, 1115–1122.
- Onoe, T., Kaneko, A., Dempo, K., Ogawa, K. & Minase, T. 1975 Alpha-fetoprotein and early histological changes of hepatic tissue in DAB-hepatocarcinogenesis. *Ann. NY Acad. Sci.* **259**, 168–180.
- Opie, E. I. 1944 The pathogenesis of tumours of the liver produced by butter yellow. *J. Expl. Med.* **80**, 231–246.
- Pagan, R., Llobera, M. & Vilaro, S. 1995 Epithelial-mesenchymal transition in cultured neonatal hepatocytes. *Hepatology* **21**, 820–831.
- Pagan, R., Martin, I., Llobera, M. & Vilaro, S. 1997 Epithelial-mesenchymal transition of cultured rat neonatal hepatocytes is differentially regulated in response to epidermal growth factor and dimethyl sulfoxide. *Hepatology* **25**, 598–606.
- Park, C. M., Cha, I. H., Chung, K. B., Suh, W. H., Lee, C. H., Choi, S. Y. & Chae, Y. S. 1991 Hepatocellular carcinoma in extrahepatic bile ducts. *Acta Radiol.* **32**, 34–36.
- Potten, C. S. & Loeffler, M. 1990 Stem cells: attributes, cycles, spirals, pitfalls and uncertainties. Lessons for and from the crypt. *Development* **110**, 1001–1020.
- Rao, K. N., Shinozuka, H., Kunz, H. W. & Gill, T. D. 1984 Enhanced susceptibility to a chemical carcinogen in rats carrying MHC-linked genes influencing development (GRC). *Int. J. Cancer* **34**, 113–120.
- Rao, M. S. (and 10 others) 1989 Role of periductal and ductular epithelial cells of the adult rat pancreas in pancreatic hepatocyte lineage. A change in the differentiation commitment. *Am. J. Pathol.* **134**, 1069–1086.
- Ray, M. B., Mendenhall, C. L., French, S. W. & Gartside, P. S. 1993 Bile duct changes in alcoholic liver disease. The Veterans Administration Cooperative Study Group. *Liver* **13**, 36–45.
- Rescan, P. Y., Loreal, O., Hassell, J. R., Yamada, Y., Guillouzo, A. & Clement, B. 1993 Distribution and origin of the basement membrane component perlecan in rat liver and primary hepatocyte culture. *Am. J. Pathol.* **142**, 199–208.
- Rogler, L. E. 1997 Selective bipotential differentiation of mouse embryonic hepatoblasts *in vitro*. *Am. J. Pathol.* **150**, 591–602.
- Rosen, E. M., Nigam, S. K. & Goldberg, I. D. 1994 Scatter factor and the c-met receptor: a paradigm for mesenchymal/epithelial interaction. *J. Cell Biol.* **127**, 1783–1787.
- Roskams, T., De Vos, R., van den Ord, J. & Desmet, V. 1991 Cells with neuroendocrine features in regenerating human liver. *Apmis Suppl.* **23**, 32–39.

- Roskams, T., Campos, R. V., Drucker, D. J. & Desmet, V. J. 1993a Reactive human bile ductules express parathyroid hormone-related peptide. *Histopathology* **23**, 11–19.
- Roskams, T., Willems, M., Campos, R. V., Drucker, D. J., Yap, S. H. & Desmet, V. J. 1993b Parathyroid hormone-related peptide expression in primary and metastatic liver tumours. *Histopathology* **23**, 519–525.
- Roskams, T., Moshage, H., Depla, E., Willems, M., Desmet, V. & Yap, P. 1995 Parathyroid hormone-related peptide is expressed and rapidly inducible in human liver cell cultures that have a bile duct phenotype. *J. Hepatol.* **23**, 160–165.
- Sarraf, C. E., Lalani, E.-N., Golding, M., Anilkumar, T. V., Poulsom, R. & Alison, M. R. 1994 Cell behaviour in the acetylaminofluorene-treated regenerating rat liver. Light and microscopic observations. *Am. J. Pathol.* **145**, 1114–1126.
- Schirmacher, P., Geerts, A., Pietrangelo, A., Dienes, H. P. & Rogler, C. E. 1992 Hepatocyte growth factor/hepatopoietin A is expressed in fat-storing cells from rat liver but not myofibroblast-like cells derived from fat-storing cells. *Hepatology* **15**, 5–11.
- Schmitt-Graf, A., Druger, S., Bochar, F., Gabbiani, G. & Denk, H. 1991 Modulation of alpha smooth muscle actin and desmin expression in perisinusoidal cells of normal and diseased human livers. *Am. J. Pathol.* **138**, 1233–1242.
- Sell, S. 1980 Heterogeneity of alpha-fetoprotein (AFP) and albumin containing cells in normal and pathological permissive states for AFP production: AFP containing cells induced in adult rats recapitulate the appearance of AFP containing hepatocytes in fetal rats. *Oncodev. Biol. Med.* **1**, 93–105.
- Sell, S. 1983 Comparison of oval cells induced in rat liver by feeding N-2-fluorenylacetylamide in a choline-devoid diet and bile duct cells induced by feeding 4,4'-diaminodiphenylmethane. *Cancer Res.* **43**, 1761–1767.
- Sell, S. 1990 Is there a liver stem cell? *Cancer Res.* **50**, 3811–3815.
- Sell, S. 1993a Cellular origin of cancer: dedifferentiation or stem cell maturation arrest? *Environ. Health Perspect.* **5**, 15–26.
- Sell, S. 1993b The role of determined stem-cells in the cellular lineage of hepatocellular carcinoma. *Int. J. Dev. Biol.* **37**, 189–201.
- Sell, S. 1994 Liver stem cells. *Mod. Pathol.* **7**, 105–112.
- Sell, S. & Dunsford, H. A. 1989 Evidence for the stem cell origin of hepatocellular carcinoma and cholangiocarcinoma. *Am. J. Pathol.* **134**, 1347–1363.
- Sell, S. & Pierce, G. B. 1994 Maturation arrest of stem cell differentiation is a common pathway for the cellular origin of teratocarcinomas and epithelial cancers. *Lab. Invest.* **70**, 6–22.
- Sell, S. & Salman, J. 1984 Light- and electron-microscopic autoradiographic analysis of proliferating cells during the early stages of chemical hepatocarcinogenesis in the rat induced by feeding N-2-fluorenylacetylamide in a choline-deficient diet. *Am. J. Pathol.* **114**, 287–300.
- Sell, S., Leffert, H. L., Shinozuka, H., Lombardi, B. & Gochman, N. 1981 Rapid development of large numbers of alpha-fetoprotein-containing 'oval' cells in the liver of rats fed N-2-fluorenylacetylamide in a choline-devoid diet. *Gann* **72**, 479–487.
- Sell, S., Hunt, J. M., Dunsford, H. A. & Chisari, F. V. 1991 Synergy between hepatitis B virus expression and chemical hepatocarcinogens in transgenic mice. *Cancer Res.* **51**, 1278–1285.
- Shah, K. D. & Gerber, M. A. 1990 Development of intrahepatic bile ducts in humans. Possible role of laminin. *Arch. Pathol. Lab. Med.* **114**, 597–600.
- Shinagawa, T., Tadokoro, M., Maeyama, S., Maeda, C., Yamaguchi, S., Morohoshi, T. & Ishikawa, E. 1995 Alpha fetoprotein-producing acinar cell carcinoma of the pancreas showing multiple lines of differentiation. *Virchows Arch.* **426**, 419–423.
- Shiojiri, N., Lemire, J. M. & Fausto, N. 1991 Cell lineages and oval cell progenitors in rat liver development. *Cancer Res.* **51**, 2611–2620.
- Sigal, S. H., Brill, S., Fiorino, A. S. & Reid, L. M. 1992 The liver as a stem cell and lineage system. *Am. J. Physiol.* **263**, G139–G148.
- Sirica, A. E., Sattler, C. A. & Cihla, H.-P. 1985 Characterisation of a primary bile ductular cell cultures from the livers of rats during experimental cholestasis. *Am. J. Pathol.* **120**, 670–678.
- Sirica, A. E., Mathis, G. A., Sano, N. & Elmore, L. W. 1990 Isolation, culture, and transplantation of intrahepatic biliary epithelial cells and oval cells. *Pathobiology* **58**, 44–64.
- Sirica, A. E., Gaaney, T. W. & Mumaw, V. R. 1994 Ductular hepatocytes. Evidence for a bile ductular cell origin in furan-treated rats. *Am. J. Pathol.* **145**, 375–383.
- Slack, J. 1995 Developmental biology of the pancreas. *Development* **121**, 1569–1580.
- Smith, P. G., Tee, L. B. & Yeoh, G. C. 1996 Appearance of oval cells in the liver of rats after long-term exposure to ethanol. *Hepatology* **23**, 145–154.
- Stamatoglou, S. C. & Hughes, R. C. 1994 Cell adhesion molecules in liver function and pattern formation. *FASEB J.* **8**, 420–427.
- Steinberg, P., Steinbrecher, R., Radaeva, S., Schirmacher, P., Dienes, H. P., Oesch, F. & Bannasch, P. 1994 Oval cell lines OC/CDE 6 and OC/CDE 22 give rise to cholangio-cellular and undifferentiated carcinomas after transformation. *Lab. Invest.* **71**, 700–709.
- Steiner, J. & Carruthers, J. 1961 Studies on the fine structure of the terminal branches of the biliary tree. *Am. J. Pathol.* **38**, 639–661.
- Stoker, M. & Perryman, M. 1985 An epithelial scatter factor released by embryo fibroblasts. *J. Cell Sci.* **77**, 209–213.
- Strain, A. J., Ismail, T., Tsubouchi, H., Arakaki, N., Hishida, T., Kitamura, N., Daikuhara, Y. & McMaster, P. 1991 Native and recombinant human hepatocyte growth factors are highly potent promoters of DNA synthesis in both human and rat hepatocytes. *J. Clin. Invest.* **87**, 1853–1857.
- Strain, A. J., Wallace, L., Joplin, R., Daikuhara, Y., Ishii, T., Kelly, D. A. & Neuberger, J. M. 1995 Characterization of biliary epithelial cells isolated from needle biopsies of human liver in the presence of hepatocyte growth factor. *Am. J. Pathol.* **146**, 537–545.
- Suzuki, Y., Ishizuka, H., Kaneda, H. & Taniguchi, N. 1987 Gamma-glutamyl transpeptidase in rat liver during 3'-Me-DAB hepatocarcinogenesis: immunohistochemical and enzyme histochemical study. *J. Histochem. Cytochem.* **35**, 3–7.
- Taniguchi, H., Toyoshima, T., Fukao, K. & Nakauchi, H. 1996 Presence of hematopoietic stem cells in the adult liver. *Nat. Med.* **2**, 198–203.
- Tarsetti, F., Lenzi, R., Salvi, R., Schuler, E., Rijhsinghani, K., Lenzen, R. & Tavoloni, N. 1993 Liver carcinogenesis associated with feeding of ethionine in a choline-free diet: evidence against a role of oval cells in the emergence of hepatocellular carcinoma. *Hepatology* **18**, 596–603.
- Tatematsu, M., Kaku, T., Medline, A. & Farber, E. 1985 Intestinal metaplasia as a common option of oval cells in relation to cholangiofibrosis in liver of rats exposed to 2-acetylaminofluorene. *Lab. Invest.* **52**, 354–362.
- Tateno, C. & Yoshizato, K. 1996 Long term cultivation of adult rat hepatocytes that undergo multiple cell divisions and express normal parenchymal phenotypes. *Am. J. Pathol.* **148**, 383–392.
- Tee, L. B., Kirilak, Y., Huang, W. H., Morgan, R. H. & Yeoh, G. C. 1994 Differentiation of oval cells into duct-like cells in preneoplastic liver of rats placed on a choline-deficient diet supplemented with ethionine. *Carcinogenesis* **15**, 2747–2756.
- Terada, T. & Nakanuma, Y. 1993 Development of human intrahepatic peribiliary glands. Histological, keratin immunohistochemical, and mucus histochemical analyses. *Lab. Invest.* **68**, 261–269.
- Terada, T., Okada, Y. & Nakanuma, Y. 1995 Expression of matrix proteinases during human intrahepatic bile duct development. A possible role in biliary cell migration. *Am. J. Pathol.* **147**, 1207–1213.

- Thorgeirsson, S. S., Evarts, R. P., Bisgaard, H. C., Fujio, K. & Hu, Z. 1993 Hepatic stem cell compartment: activation and lineage commitment. *Proc. Soc. Expl Biol. Med.* **204**, 253–260.
- Tsao, M. S. & Grisham, J. W. 1987 Hepatocarcinomas, cholangiocarcinomas, and hepatoblastomas produced by chemically transformed cultured rat liver epithelial cells. A light- and electron-microscopic analysis. *Am. J. Pathol.* **127**, 168–181.
- Tsao, M. S., Smith, J. D., Nelson, K. G. & Grisham, J. W. 1984 A diploid epithelial cell line from normal adult rat liver with phenotypic properties of 'oval' cells. *Expl Cell Res.* **154**, 38–52.
- Tsao, M. S., Grisham, J. W., Nelson, K. G. & Smith, J. D. 1985 Phenotypic and karyotypic changes induced in cultured rat hepatic epithelial cells that express the 'oval' cell phenotype by exposure to N-methyl-N'-nitro-N-nitrosoguanidine. *Am. J. Pathol.* **118**, 306–315.
- Tyner, A. L., Godbout, R., Compton, R. S. & Tilghman, S. M. 1990 The ontogeny of alpha-fetoprotein gene expression in the mouse gastrointestinal tract. *J. Cell Biol.* **110**, 915–927.
- Uchida, T. & Peters, R. L. 1983 The nature and origin of proliferated bile ductules in alcoholic liver disease. *Am. J. Clin. Pathol.* **79**, 326–333.
- Van-Eyken, P., Sciot, R., Callea, F., van der Steen, K., Moerman, P. & Desmet, V. J. 1988a The development of the intrahepatic bile ducts in man: a keratin-immunohistochemical study. *Hepatology* **8**, 1586–1595.
- Van-Eyken, P., Sciot, R. & Desmet, V. J. 1988b A cytokeratin immunohistochemical study of alcoholic liver disease: evidence that hepatocytes can express 'bile duct-type' cytokeratins. *Histopathology* **13**, 605–617.
- Van-Eyken, P., Sciot, R. & Desmet, V. J. 1989 A cytokeratin immunohistochemical study of cholestatic liver disease: evidence that hepatocytes can express 'bile duct-type' cytokeratins. *Histopathology* **15**, 125–135.
- Vandersteenhoven, A. M., Burchette, J. & Michalopoulos, G. 1990 Characterization of ductular hepatocytes in end-stage cirrhosis. *Arch. Pathol. Lab. Med.* **114**, 403–406.
- Wilson, J. & Leduc, E. H. 1958 Role of cholangioles in restoration of the liver of the mouse after dietary injury. *J. Path. Bact.* **76**, 441–449.
- Wright, N. & Alison, M. (eds) 1984 The conditional renewal populations: the liver. In *The biology of epithelial cell populations*, vol. 2, pp. 873–980. Oxford: Clarendon.
- Wu, H., Wade, M., Krall, L., Grisham, J., Xiong, Y. & Vandyke, T. 1996 Targeted *in vivo* expression of the cyclin dependent kinase inhibitor p21 halts hepatocyte cell cycle progression, postnatal liver development, and regeneration. *Genes Dev.* **10**, 245–260.
- Yang, L., Faris, R. A. & Hixson, D. C. 1993 Phenotypic heterogeneity within clonogenic ductal cell populations isolated from normal adult rat liver. *Proc. Soc. Expl Biol. Med.* **204**, 280–288.
- Yasui, O., Miura, N., Terada, K., Kawarada, Y., Koyama, K. & Sugiyama, T. 1997 Isolation of oval cells from Long-Evans Cinnamon rats and their transformation into hepatocytes *in vivo* in the rat liver. *Hepatology* **25**, 329–334.
- Yaswen, P., Goyette, M., Shank, P. R. & Fausto, N. 1985 Expression of c-Ki-ras, c-Ha-ras, and c-myc in specific cell types during hepatocarcinogenesis. *Molec. Cell Biol.* **5**, 780–786.
- Yavorkovsky, L., Lai, E., Ilic, Z. & Sell, S. 1995 Participation of small intraportal stem cells in the restitutive response of the liver to periportal necrosis induced by allyl alcohol. *Hepatology* **21**, 1702–1712.
- Zajicek, G., Oren, J. & Weinreb, M. 1985 The streaming liver. *Liver* **5**, 293–300.
- Zalatnai, A. & Schally, A. V. 1990 Hepatic lesions in Syrian golden hamsters with pancreatic carcinoma induced by N-nitrosobis(2-oxopropyl)amine (BOP). *Acta. Morphol. Hung.* **38**, 119–130.