How much will Herceptin really cost?

Balancing local and national priorities is not new

EDITOR—Herceptin for early stage breast cancer is a cost effective treatment, and it is right that the NHS should make it available. The challenge of funding new treatments and balancing local and national priorities is one the NHS has faced since its inception.¹ The process the National Institute for Health and Clinical Excellence (NICE) follows for evaluating new medicines is the most rigorous anywhere in the world, and local NHS organisations can have confidence that implementing our recommendations is an effective use of their resources.

NICE's independent advisory committees do not rubber stamp decisions. They make them based on an objective and impartial interpretation of the evidence. Those who suggest otherwise should come forward with their evidence for doing so.

It is wrong to suggest that a single recommendation from NICE will necessarily cause other services to be reduced and that these decisions are made by clinicians only. Many demands are made on available funds in any hospital at any one time and it is the responsibility of local NHS management, working with their clinical teams, to make the best use of these resources. This difficult job is made easier by NICE guidance and the support tools we offer, such as our commissioning guides and our forthcoming advice on shifting spending from less to more effective treatments.

Andrew Dillon *chief executive*
lucy.betterton@nice.org.uk

Peter Littlejohns *clinical and public health director*
National Institute for Health and Clinical Excellence, London WC1V 6NA

Competing interests: None declared.

¹ Barrett A, Roques T, Small M, Smith RD. How much will Herceptin really cost? *BMJ* 2006;333:1118-20. (25 November.)

The money is already there

EDITOR—Barrett et al state that “the real cost of Herceptin is in the other patients not treated, whether they are patients with cancer or those with other conditions”—the opportunity cost.¹ The competing demands that medical professionals across multiple disciplines place on limited resources and their responsibility to ensure the judicious

use of publicly funded resources should also have been acknowledged.

It is too easy to blame the National Institute for Health and Clinical Excellence (NICE) when healthcare professionals can enable judicious use of what has already been provided. Moon and Bogle highlighted savings of £1.1 bn (€1.6bn; \$2.1bn) over five years by promoting the generic substitution of statins for the primary and secondary prevention of cardiovascular disease,² and Barrett et al's strategic health authority estimates savings of £20m this way.³

Clinicians currently have little incentive to seek prescribing savings, particularly if savings are then used to stem deficits. Devolving prescribing costs into locally held unified budgets for clinical service delivery across multiple diseases would promote negotiation and priority setting. Oncologists, cardiologists, and general practitioners would have to talk to each other and ensure the sensible use of public funds.

Introducing a system of reference pricing in which drug reimbursement is pegged to the price of a therapeutically equivalent generic drug could do much to resolve these types of issues at source. Clinicians could decide whether to pay a premium for a non-generic drug from regional budgets or fund new technologies such as Herceptin.

External pressures that should not determine who is treated and what they are treated with can affect prescribing decisions. Specialist professional groups that adopt a narrow viewpoint can be such a pressure. Their guidelines are usually uncoded and promote recommendations that do not consider the impact across other specialties. Uncoded recommendations may be expensive to implement in both mortality and financial terms.

Disinvestment will promote efficient allocation, but as clinicians we must also disinvest ourselves of a myopic clinical mindset that results in “silo medicine.” Expecting NICE to provide all the answers is to be blind to our own faults and only shifts the blame.

Rubin Minhas *general practitioner*
Sunlight Medical Centre, Gillingham, Kent
ME7 1LX
anything@rubin.icom43.net

P. MARAZZI/ISPL

Competing interests: Over the past 12 months RM has attended advisory boards for Novartis and Pfizer. Over the past 10 years RM has attended educational meetings, received travel grants, honorariums for lectures and advisory boards from a number of pharmaceutical companies, including AstraZeneca, Bayer, Fournier, GlaxoSmithKline, Pfizer, Merck, MSD, and Sanofi-Aventis. RM is a member of the Primary Care Cardiovascular Society and the South Asian Health Foundation. RM is an appraisal committee member in NICE and a participant in NICE guideline development groups.

- 1 Barrett A, Roques T, Small M, Smith RD. How much will Herceptin really cost? *BMJ* 2006;333:1118-20. (25 November.)
- 2 Moon JC, Bogle RG. Switching statins. *BMJ* 2006;332:1344-5. (10 June.)
- 3 Gill K. Statins—the SHA perspective. Achieving clinical efficiency. A practical case study. www.heart.nhs.uk/scripts/default.asp?site_id=23&id=28050 (accessed 24 Nov 2006).

bmj.com

Summary of other responses

The 19 responses to the article by Barrett et al express concerns about the independence of the National Institute for Health and Clinical Excellence (NICE), its remit, and its role in relation to primary care trusts.¹

People are divided on whether NICE should decide which services to cut to fund new technologies (opportunity costs). But without an implementing body it is unclear how to react to NICE's recommendations. NICE's remit is not the equitable distribution of a limited healthcare budget, but its guidance should provide healthcare professionals and political bodies with a means to make informed decisions and communicate these to the public. The public in turn needs to accept that health care is a finite resource, and that the decision is between rationing and paying higher taxes.

The role of primary care trusts as strong local commissioners is emphasised by several correspondents—postcode prescribing, rather than being a lottery, is their *raison d'être*, and as such they are in the best position to judge local needs. They might also be expected to put their case for bigger budgets to the politicians who control national budgets and to the public. They are in a position to support clinicians in facing up to media, politicians, and pressure groups.

Two correspondents provide examples of how NICE guidance is already restricting treatment options—in palliative care and in chemotherapy. And one correspondent pragmatically suggests a budget split into “historical” and “innovative” treatments.

Yet others question whether cost effectiveness is the right prioritisation tool. And,

given that breast cancer is not a disease of ageing any more, these kinds of prescribing dilemma may in future be avoided altogether if the focus were on health and prevention.

The role of the media is touched on by several correspondents, most notably a doctor and former patient with breast cancer, Jane Cooper. She got caught up in the media circus surrounding Herceptin and, after discussions with her oncologist, decided against taking it. Benjamin White, a senior house officer in critical care, sums it up nicely. Either "the profession lays itself open to blame for all deficiencies in service or we face the possibility of health care for those who shout loudest. Clearly the public needs to be properly informed about what to expect from their health service, but we cannot risk the debate being hijacked by those groups who are most vocal or well funded."

Birte Twisselmann assistant editor, (web)
BMJ

Competing interests: None declared.

1 Rapid responses. How much will Herceptin really cost?
www.bmj.com/cgi/eletters/333/7578/1118.

Influenza vaccine uptake among British Muslims attending Hajj, 2005 and 2006

See Editorial by Gabrad *et al*

EDITOR—Every year 1% of British Muslims travel to Saudi Arabia on the Hajj pilgrimage. High rates of influenza have been reported among pilgrims,^{1,2} and the Saudi Arabian Ministry of Health recommends that all pilgrims should receive influenza vaccination before travelling. We determined the uptake of influenza vaccination among British Hajj pilgrims who attended the British Hajj Delegation Clinic in Mecca and mobile clinics set up by us in Mina in 2005 and 2006.

In 2005, 196 pilgrims were recruited and in 2006, 146 (age range 2 to 83; 313 men); 104 were from London. Ninety two pilgrims (27%) were categorised as being at risk of influenza, one fifth (19) of them solely because of being 65 or over and the rest because of having chronic diseases regardless of age.

A total of 108 pilgrims (32%) were vaccinated, all but two by their general practitioner. Vaccine coverage among pilgrims at risk was 55% (51/92) and among those not at risk 23% (57/250) ($P < 0.001$). Coverage among over-65s was 63% (19/30) and among under-65s regardless of underlying chronic disease 52% (32/62).

Immunisation uptake by all pilgrims in 2005 was 27% (53/196) and 38% (55/146) in 2006 ($P = 0.036$), and the coverage among those at risk was 51% (29/57) and 63% (22/35) respectively ($P = 0.26$). Uptake by all pilgrims who lived in London in these two years was 20% (21/104) compared with 37% (87/238) in other centres ($P = 0.003$). Uptake by "at risk" residents of London was

38% (10/26) compared with 62% (41/66) elsewhere ($P = 0.04$).

The rate of vaccination among Muslims at risk is worryingly lower (55%) than the nationwide uptake rate among the target population (70%),³ and even lower among the residents of London. Uptake could be improved by advertising, phone calls, patient and staff education, standing orders, targeting people with high risk behaviours at their meeting venues, and offering home visits by nurses.⁴ Clinics set up, in partnership with community leaders, travel agents, and imams for pilgrims planning to go on Hajj might further improve uptake.

Shuja Shafi consultant microbiologist
shuja.shafi@nwlh.nhs.uk

Harunor Rashid
Kamal Ali
Haitham El Bashir
Elizabeth Haworth
Ziad A Memish

Robert Booy
Department of Microbiology and HPA
Collaborating Laboratory, Northwick Park
Hospital, Harrow, Middlesex HA1 3UJ

Funding: This work was supported from a Department of Health grant. We thank the British Hajj Delegation team, European Hajj Mission and the GPs for help with the data collection and staff at HPA laboratory, London & Respiratory Virus Unit, Health Protection Agency, Centre for Infections, London, for performing the PCR tests. Ethical Approval: Multicentre Research Ethics Committee, UK (MREC 02/2/12).

Competing interests Roche supplied free courses of Tamiflu for treatment of pilgrims with influenza infection on near-patient testing. SS served as Chair, Health and Medical Committee of the Muslim Council of Britain (until August 2006)

1 El Bashir H, Haworth E, Zambon M, Shafi S, Zuckerman J, Booy R. Influenza among UK pilgrims to Hajj, 2003. *Emerg Infect Dis* 2004;10:882-3.

2 Balkhy HH, Memish ZA, Bafaqeer S, Almuneef MA. Influenza a common viral infection among Hajj pilgrims: time for routine surveillance and vaccination. *J Travel Med* 2004;11:82-6.

3 Muller D, Nguyen-Van-Tam JS, Szucs TD. Influenza vaccination coverage rates in the UK: A comparison of two monitoring methods during the 2002-2003 and 2003-2004 seasons. *Public Health* 2006;120:1074-80.

4 Ompad DC, Galea S, Vlahov D. Distribution of influenza vaccine to high-risk groups. *Epidemiol Rev* 2006; 28:54-70.

How many whole time equivalents rather than heads in the NHS?

EDITOR—At about the time that the government announced the target for the increased numbers of consultants, general practitioners, and nurses in the NHS, plans were developed for more flexible working. This

was to enable those with domestic responsibilities to work part time and to allow flexible retirement. Increasing numbers of consultants are retiring and then coming back to work up to half time. This is good for all concerned, especially the Department of Health, which counts each part time post in the same way as a whole time post.

When talking about the increase in staff numbers,¹ Patricia Hewitt and her team very rarely, if ever, mention whole time equivalents. No one ever seems to ask her about these either.

Apples need to be compared with apples. We should know the increase in whole time equivalents, not the increased head count. The overall head count may have increased, but the whole time equivalent, which better represents the hours available to treat patients, may even have gone down. We just don't know.

Anthea J Sherwood consultant histopathologist
Derriford Hospital, Plymouth PL6 8DH
anthea.sherwood@phnt.swest.nhs.uk

Competing interests: AJS is a member of the Central Consultants and Specialists Committee (CCSC).

1 Eaton L. NHS employs too many hospital doctors, minister says. *BMJ* 2006;333:1086-a. (25 November.)

Fever epidemic moves into Sri Lanka

EDITOR—Patients with chikungunya viral infections also spread by the aedes mosquito are being reported from Sri Lanka. Refugees from Indian cities where this disease is epidemic have returned to some cities¹ and may have carried the virus across.

Over the past months, many cases of chikungunya viral infection have been reported from India, some islands in the Indian Ocean, Malaysia, and other South East Asian countries.^{2,3} According to the World Health Organization, over 1 million suspected cases have been reported since the beginning of this year. Imported cases have been reported from Europe, the United States, and Australia.⁴ In the United Kingdom, 93 cases have been reported by the Health Protection Agency in the first eight months of this year among travellers returning from regions where the disease is endemic.⁵ In the US, cases were reported in Louisiana and Maryland among returning travellers, prompting the Centers for Disease Control and Prevention to advise clinicians to be alert to the appearance of more cases.

Effective mosquito control measures are paramount in keeping these diseases under control. Both political will and community participation must come together to make effective strides in protecting people's health from these different mosquito borne diseases. If not, people from these regions will continue to be held to ransom by these different viruses, protozoa, and other microorganisms using the mosquito as the messenger service.

Suranjith L Seneviratne *consultant clinical immunologist*

Department of Clinical Immunology, Manchester Royal Infirmary, Manchester M13 9WL
suran200@yahoo.co.uk

Jennifer Perera *professor of microbiology*
University of Colombo, Sri Lanka

Competing interests: None declared.

- 1 Mudur G. Failure to control mosquitoes has led to two fever epidemics in India. *BMJ* 2006;333:773-c. (14 October.)
- 2 Hermon YE. Virological investigations of arbovirus infections in Ceylon, with special reference to the recent Chikungunya fever epidemic. *Ceylon Med J* 1967;12:81-92.
- 3 Higgs S. The 2005-2006 Chikungunya epidemic in the Indian Ocean. *Vector Borne Zoonotic Dis* 2006;6:115-6.
- 4 Nau JY. [Chikungunya, when politicians are interested in arbovirus infections (I)]. *Rev Med Suisse* 2006;2:617.
- 5 Pfeffer M, Loscher T. Cases of chikungunya imported into Europe. *Euro Surveill* 2006;11:E060316 2.

Getting knowledge into clinical practice

EDITOR—Six years ago the Nicaraguan government asked us to implement treatment guidelines for the most prevalent health problems of the country in six months. After some discussion, a stepwise development of a treatment guideline for postpartum haemorrhage was selected as a training example.

Several Nicaraguan institutions had already published treatment guidelines for postpartum haemorrhage, but the real management of delivery differs from their recommendations. The causes seemed simple: treatment guidelines are written far from the clinical reality and are usually imposed, which translates into low adherence. Thus a crucial step is to get clinicians involved with the elaboration of treatment guidelines. We used the results of a drug use study as the stimulus to realising what is happening in real practice, to ask why things happen, and to look for answers, adapting evidence into treatment guidelines.

Postpartum haemorrhage has well known and affordable management, and its high incidence in less developed countries is unjustifiable.¹ Furthermore, the inefficacy of ergometrine in extreme heat and humidity conditions when appropriate conservation cannot be ensured was described more than 20 years ago.² Despite the published evidence, this information seems not to have reached the target professionals, as it continues to be widely used in hospitals where it is stored at room temperature.³

In this century of information, oversaturation is easy. Health professionals should be reading 19 scientific articles each day of the year to keep up to date.⁴ The gap between scientific knowledge and health professionals' knowledge increasingly widens: lack of time is combined with difficult access to information and, sometimes, idiomatic barriers.

Some professionals ignore the fact that a prescription habit is obsolete or wrong; and not all relevant clinical evidence reaches prescribers. Without questioning one's own prescribing habits, it is difficult to identify practices that need to be updated. Despite being observational, drug use studies help

clarify doubts in clinical settings, introduce the idea of conducting research, and stimulate the information seeking process to reduce the gap between knowing and doing.

Independent digested information and continuous medical education further improve prescription. The latter, although not universally considered a worthwhile investment by health authorities, at least seems better than leaving it to drug manufacturers.⁵

Albert Figueras *manager, international cooperation area*

Fundació Institut Català de Farmacologia, U Autònoma de Barcelona, 08035 Barcelona, Spain
afs@icf.uab.es

Edgar Narváez *manager, technical unit of medical evaluation*

Instituto Nicaragüense del Seguro Social, Managua, Nicaragua

A grant from the Agència Catalana de Cooperació al Desenvolupament is helping to develop a Latin-American Network Against Maternal Mortality (GIRMMAH, www.girmmah.net) an intervention based on the discussed considerations.

Competing interests: None declared.

- 1 United Nations Population Fund. State of the World Population 2004. *The Cairo consensus at ten: population, reproductive health, and global effort to end poverty*. UNFPA, New York, 2005. www.unfpa.org/upload/lib_pub_file/327_filename_en_swp04.pdf (accessed 28 Sep 2006).
- 2 Walker GJ, Hogerzeil HV. Potency of ergometrine in tropical countries. *Lancet* 1988;2:393.
- 3 Figueras A, Narváez E, Aguilera C, Laporte JR. Securing reproductive rights. *Lancet* 2004;363:989.
- 4 Rawlins M. St. Paul International Health Care Annual Lecture on Sept. 7th, 1999 <http://www.nice.org.uk/page.aspx?o=27856>.
- 5 Drug companies influence on medical education in USA. *Lancet* 2000;356:781.

Mental Health Act: why not learn from devolution?

EDITOR—The UK government's third attempt to introduce an amended Mental Health Act is already meeting the same resistance as its previous attempts.¹ The reasonable concerns of ministers in the Home Office about protecting citizens from violent people, who are likely to offend, have been interpreted in such a way in the proposed bill as to achieve a degree of unity in opposition from those in the mental health field that would have seemed improbable 10 years ago. Moreover they have done so not once or twice but now three times.

At the same time, the Scottish government, when I was deputy justice minister, successfully introduced a new act that not only protects our citizens but also met with broad agreement and now supports a similar alliance in Scotland that condemns the English government's approach. The two previous acts in the 20th century were passed first in the UK parliament applying to England and Wales and were subsequently amended marginally to become law in Scotland.

My question is a simple one. Why not use the benefits of devolution and adopt the Scottish Act amended to suit English needs?² Psychiatrists in the former Soviet Union were condemned for acting on behalf of the state to incarcerate citizens who did

not suffer from a treatable illness. English and Welsh psychiatrists should not be forced to put themselves in a similar position. Are we to be the only signatory out of 45 nations to reserve the right not to comply with the Council of Europe's Recommendation³ on protecting the human rights and dignity of people with mental disorders?

As Dyer has put it so concisely, if people who do not have a treatable mental illness are deemed a danger to others then criminal proceedings need to be implemented, if appropriate.

Richard Simpson *consultant psychiatrist*

Lothian NHS, Livingston, West Lothian EH54 6QF
richard.simpson@hushmail.com

Competing interests: None declared.

- 1 Dyer C. Third time lucky? *BMJ* 2006;333:1090. (25 November.)
- 2 Mental Health (Care and Treatment) [Scotland] Act 2003.
- 3 Council of Europe, Committee of Ministers, Recommendation No REC (2004)10 [concerning the protection of the human rights and dignity of persons with mental disorders].

WHO obesity charter seems more like a mantra

EDITOR—Every time the World Health Organization publishes a nutrition related strategy, whether for cardiovascular disease, obesity, or general health, there are calls to reduce fat, sugar, and salt. As far as I am aware, there is no relation between salt and the aetiology, prevention, and treatment of obesity. This makes me wonder whether the ideas in the new obesity charter are evidence based or just another mantra.¹

As usual, many regulatory initiatives are aimed at the multinational food companies and few at the day to day drivers of our obesogenic lifestyle—such as fast food outlets (most of which are not chain stores), burger or chip vans (particularly those parked outside schools every lunchtime), out of town shopping, low cost home entertainment, lack of cooking skills among young people, unsafe open spaces for play and recreation, long working hours, lack of showers and bicycle parking, unimaginative physical education at school, low cost and readily available alcohol, insufficient NHS obesity services (particularly for children).

It is good that European governments agree on obesity action. However, I am concerned that European charters focus too much attention on headline grabbing initiatives aimed at industry—an easy target since many health professionals and public interest groups feel uncomfortable about profit—and too little attention on all the other drivers of obesity, especially those experienced by lower income people who are most at risk of obesity.

Carrie H Ruxton *freelance dietitian*

Cupar, Fife KY15 4EA
carrie@nutrition-communications.com

Competing interests: CHR consults to the NHS (writes obesity strategies) and to the food industry.

- 1 Groves T. Pandemic obesity in Europe. *BMJ* 2006;333:1081-2. (25 November.)

Shark cartilage in the water

One size does not fit all

EDITOR—I agree with many of Waxman's points, but one size does not fit all.¹ The efficacy and potential physical harm of someone self administering coffee enemas is different from that of someone receiving aromatherapy for improvement of general wellbeing. A distinction needs to be made between "complementary" and "alternative" therapies. The psychological and spiritual domains are as important to many patients as their objective response to treatments.

Rather than being a "battle," for many patients their experience of cancer treatment is a helplessly passive one. Accessing complementary therapies can be an empowering experience regardless of the anticipated benefits. If 80% of patients are seeking out additional or alternative therapies, we need to equip ourselves with a knowledge and evidence base from which to advise patients. The websites often warn patients that doctors will disparage the use of their product, and we have a responsibility not to push our patients towards seeking alternatives by giving reflex responses when other therapies are brought up in clinic.

The use of complementary therapies and alternative cures is commonplace in palliative care. Many take the view that they have "nothing to lose," and it is here where the patient may be most vulnerable to therapies that claim a cure. Such claims can inhibit realistic goal setting and future planning. In addition, complex bereavement is often worsened by unrealistic goals that were set by the use of alternative cures. The internet can be an anarchic library when it comes to cancer cures and, regardless of what legislation is developed locally, this will not stop people accessing information and medicines from elsewhere. We need to be able to advise patients with credibility and empathy and not throw the baby out with the bathwater when it comes to therapies that are there purely to support and edify the individual.

Simon I R Noble *senior lecturer in palliative medicine*
Cardiff University and Royal Gwent Hospital
NP20 2UB
simon.noble@gwent.wales.nhs.uk

Competing interests: None declared.

¹ Waxman J. Shark cartilage in the water. *BMJ* 2006;333:1129. (25 November.)

Effective legislation is already in place but is not being properly used

EDITOR—Waxman has identified the need for effective legislation to protect vulnerable cancer patients from unproved and/or harmful cancer remedies.¹

The UK 1939 Cancer Act in section 4.1 prohibits the publication of any advertisement containing an offer to treat any person for cancer.

I have now made successful use of this legislation in relation to patients who have used cancer therapies that have been adver-

tised in both the complementary health sector and by registered medical practitioners associated with a registered pharmacy.^{2 3}

If doctors identify complementary practitioners, commercial companies, or even registered health professionals who are advertising cancer remedies they should contact their local trading standards department, who have a responsibility to enforce the act. In the case of registered medical practitioners, the General Medical Council should also be notified.

Charles B Shepherd *physician*
Stroud, Gloucestershire GL6 8EH
charlesbshepherd@lineone.net

Competing interests: None declared.

¹ Waxman J. Shark cartilage in the water. *BMJ* 2006;333:1129. (25 November.)

² Dyer O. British company fined for falling foul of Cancer Act. *BMJ* 1994;308:158.

³ Campbell D. "Quackery" outlawed in registered pharmacy. *BMJ* 1997;315:625.

Ignorance based cancer treatment

EDITOR—I write with reference to Waxman's personal view.¹ A reflexologist believes that the organs of the body reappear symbolically on the soles of the feet and that disease of these organs can be treated by massaging that part of the foot. Anyone who believes that is either woefully ignorant of human biology and seriously misinformed, or barking mad.

I spent most of a year, from November 2002 to October 2003, in a cancer ward, being treated for the complications of two different types of lymphoma. During that time the nurses repeatedly offered me the services of a reflexologist.

After two months in bed with septicaemia, my toenails had grown too long for me to wear my slippers comfortably. I could not cut them myself and a nurse told me that she could not, as it was against some kind of rule.

I am all for hospitals offering non-medical treatments to improve the happiness and well-being of patients. Hairdressing—for those of us with hair—and skin and nail care would be welcome. The same hospital trust now offers cancer patients the services of Reiki therapists, and has a homoeopathy hospital in its group.

It is unsurprising that a high proportion of cancer patients use improbable, potentially hazardous, non-evidence-based therapies. They come with the apparent endorsement of their NHS doctors.

Caroline Richmond *obituary writer*
London SW3 5AQ
c@roline.demon.co.uk

Competing interests: CR is a sometime cancer patient.

¹ Waxman J. Shark cartilage in the water. *BMJ* 2006;333:1129. (25 November.)

Miracle hiccough cure gets the attention it deserves

EDITOR—Reluctantly, I must thank Roberts for bringing to greater attention Lam's

miraculous hiccough cure (plugging both ears tightly, pushing both right and left tragus, and drinking the entire glass of water through the straw without pause, without releasing the pressure over the ears).¹ It certainly works. Crushingly, I was under the misapprehension that the cure was mine alone. I thought that the first professorship of hiccough medicine may well be mine also.

Firstly, hiccoughs or hiccups? Neither: the rarely used but proper term is singultus. The Latin word singult means catching one's breath while sobbing.

I was a late convert to the miracle method: I was formerly a "drinking from the wrong side of the glass" man. In contrast to the description given, I never use a straw. Slow drinking of even 20 to 30 millilitres of fluid is often sufficient. Keeping the pressure until the last swallow is complete seems beneficial. If no liquid is available, the Valsalva manoeuvre would suffice. Most cures focus on interruption of vagal or phrenic nerve irritation (which is part of the reflex arc and commonly a cause). The rise in intrathoracic pressure is enough to break that arc.

After a few successes (and from personal experience, the method is nearly infallible when applied correctly) you get an intuition about whether the cure has worked. This allows you to step back (as a true magician would) while the recipient waits indefinitely for the next spasm. This is a particularly satisfying way to bond with a patient, or make friends in the pub.

Before dashing for an antipsychotic, attempt a miracle cure. If the idea of putting your fingers in the ears of a hiccoughing friend or patient is distasteful to you, it is more socially acceptable than some cures. Digital rectal massage² and intercourse to orgasm³ (although never suggested synchronously) have both been described.

Rhys H Thomas *medical senior house officer*
Rhys.Thomas@doctors.org.uk
Gloucestershire Royal Hospital, Gloucester
GL1 3NN

Naomi J P Thomas *paediatric senior house officer*
Worcestershire Royal Hospital, Worcester
WR5 1DD

Competing interests: RHT was hoping to make his fortune from the hiccough cure. NJPT frequently benefits from the miracle cure.

¹ Roberts H. Bloodletting and miraculous cures. *BMJ* 2006;333:1127-a (25 November.)

² Fesmire F. Termination of intractable hiccups by digital rectal massage. *Ann Emerg Med* 1988;17:872.

³ Peleg R, Peleg A. Case report: sexual intercourse as potential treatment for intractable hiccups. *Can Fam Physicians* 2000;46:1631-2.

bmj.com

We select the letters for these pages from the rapid responses posted on bmj.com, favouring those received within five days of publication of the article to which they refer.

Letters are thus an early selection of rapid responses on a particular topic. Readers should consult the website for the full list of responses and any authors' replies, which usually arrive after our selection.