STUDIES ON PHAGOCYTOSIS. DETERMINATION OF BLOOD OPSONIN FOR BRUCELLA

JOSEPH VICTOR, ABOU D. POLLACK, R. RAYMOND, AND
JEANNE R. VALLIANT

Biological Laboratories, Chemical Corps, Camp Detrick, Frederick, Maryland

Received for publication January 7, 1952

The bactericidal properties of dog blood for *Brucella suis* have been described previously (Victor, Wells *et al.*, 1952). Because leucocytes were essential for bactericidal activity toward this strain, the relation between bactericidal and phagocytic activity of dog blood was investigated.

In order to measure phagocytosis satisfactorily a method must be employed yielding reproducible results. The opsonocytophagic test of Huddleson, Johnson, and Hamaan (1933) was inadequate for our purposes because it employs 0.8 per cent sodium citrate which inhibits the bactericidal action of dog blood (Victor, Wells *et al.*, 1952). Many investigators have found that sodium citrate also inhibited phagocytosis of brucellae by normal blood, presumably without interfering with the phagocytic activity of leucocytes of individuals with brucellosis. Others have suggested that heparin, which did not interfere with phagocytosis, be employed as an anticoagulant. The test devised by Huddleson *et al.* (1933) is done in tubes only intermittently agitated even though the influence of continuous agitation on phagocytosis had been demonstrated (Fenn, 1923). Furthermore, evaluation of the intensity of phagocytosis by counting the number of brucellae within individual leucocytes was a highly subjective one. Finally, the opsonocytophagic test of Huddleson *et al.* (1933) did not distinguish between opsonin titer and the phagocytic activity of leucocytes.

Experiments were performed to define factors which influence phagocytosis of *Brucella suis* by dog leucocytes; namely, phagocytic power of various types of blood leucocytes, activity in heparinized and citrated blood, effect of continuous and of intermittent agitation, effect of temperature, effect of time of agitation, comparison of phagocytosis of living and of formalin killed bacteria, variability of a standardized test procedure, and effect on phagocytes of repeated washing in an isotonic solution of electrolytes and gelatin. From these experiments a method for determination of opsonin titer in plasma was developed.

METHODS

Two-tenths ml of 20 per cent sodium citrate solution in 0.9 per cent saline was added to 5 ml of blood. The final concentration was 0.8 per cent sodium citrate, the same as that employed by Huddleson *et al.* (1933). Heparin or "liquaemin" (1 per cent solutions) was added in 0.1 ml volume to 10 ml of fresh blood. All tests were made with dog blood unless otherwise noted and initiated within 2 hours after bleeding.

Dilution of plasma and washing of blood cells were performed with a mixture

of gelatin and the Krebs-Henseleit solution as previously described (Victor, Wells *et al.*, 1952).

A smooth virulent strain (PSIII) of *B. suis* was grown for 24 hours on tryptose agar slants. It was harvested in tryptose saline solution and made up to contain a suspension of 1×10^{10} organisms per ml by matching against a cholesterol standard. Later experiments were conducted with bacterial suspensions prepared by matching in a photoelectric colorimeter. In each instance the number of organisms was checked by plate counts after diluting the original suspension. This concentration of cells of *Brucella* was the same as that employed by Huddleson *et al.* (1933) and greatly exceeded critical ranges shown by others (Fenn, 1923).

Phagocytosis was determined in mixtures of 0.1 ml of blood with 0.1 ml of organisms suspended in a Krebs gelatin solution. The test tubes were mounted in a rotator (Robertson and Sia, 1924) and rotated 24 or 36 times a minute. Experiments were conducted at 37 C in an air incubator or at room temperature (19 to 20 C).

Phagocytosis was demonstrated in smears prepared on glass slides warmed to facilitate spreading. After drying, the smears were stained with the Jenner-Giemsa stain. Wherever possible, 100 consecutive leucocytes were examined. Tallies were made of those that contained bacteria. Phagocytosis for each type of leucocyte was recorded as the percentage that contained bacteria.

RESULTS

Phagocytic power of various types of blood leucocytes in heparinized and citrated blood. Lucké *et al.* (1933) found no difference in phagocytic activity of *Mycobacterium tuberculosis* by mononuclear and polymorphonuclear leucocytes from rabbit exudates. Others have observed different phagocytic responses with different organisms in various leucocytes. With brucellae Huddleson *et al.* (1933) noted greater phagocytosis by neutrophils than by mononuclear leucocytes, and Dickey and Forbus (1945) reported chemotaxis as well as phagocytosis by neutrophils. It is noteworthy that recent editions of textbooks of histology, namely, Maximow and Bloom (1948) and Bailey's textbook by Smith and Copenhaver (1948), state that eosinophilic leucocytes have no phagocytic activity.

It is generally agreed that heparin does not impair phagocytosis. Although 0.8 per cent sodium citrate has long been known to inhibit phagocytosis (Otani, 1918), Castaneda *et al.* (1942) and Tovar (1944) found that 1.6 per cent sodium citrate was necessary to prevent phagocytosis of brucellae in "normal" blood. Such disagreements about the conditions for normal phagocytosis of brucellae made it seem desirable to study further the role of the anticoagulant.

Table 1 records the results of 20 of 44 experiments, with heparinized and with citrated blood, obtained simultaneously from the same source. The blood sample was mixed with bacteria for 30 minutes at 37 C. In heparinized blood, phagocytosis was most frequent by polymorphonuclear neutrophils. Monocytes had phagocytic activity intermediate between that of neutrophilic and of eosino-

philic leucocytes. Lymphocytes had no phagocytic activity. The percentage of polymorphonuclear neutrophils containing bacteria was greater in heparinized than in citrated blood in 36 of 44 tests. In only 8 cases did the percentage in citrated equal or exceed that in heparinized blood.

Effect of continuous and intermittent agitation on phagocytosis. Agitation is essential for phagocytic tests (Fenn, 1923; Robertson and Sia, 1924). Too little or too much agitation interferes with optimal phagocytosis. The opsonocytaphagic test of Huddleson *et al.* (1933) required citrated blood in tubes shaken intermittently by hand during 30 minutes' incubation. In the present tests with

TABLE 1
Comparison of phagocytic activity of various types of leucocytes

PERCENTAGE OF LEUCOCYTES CONTAINING BRUCELLA SUIIS						DIFFERENTIAL COUNT		
Heparin			Citrate			Neutrophile	Mononuclear	Eosinophile
Neutrophile	Mononuclear	Eosinophile	Neutrophile	Mononuclear	Eosinophile			
100	28	0	14	16	0	54	7	1
91	62	0	17	50	0	54	14	0
95	100	100	18	28	0	59	7	3
98	79	100	18	0	0	52	14	1
100	100	100	18	25	0	45	7	1
92	92	50	2	0	0	49	12	2
5	25	0	1	10	0	54	4	3
30	50	0	2	0	0	62	4	3
38	20	33	1	0	0	55	5	3
25	50	0	1	0	0	69	8	5
45	50	20	2	0	50	69	4	5
36	33	0	13	0	0	55	6	7
100	0	0	15	0	0	40	0	0
100	0	0	19	0	0	38	1	0
100	0	0	18	0	0	60	1	2
100	100	0	29	0	0	42	2	1
95	0	33	5	0	0	37	2	22
90	0	0	6	14	0	39	2	2
100	100	75	1	0	0	40	3	4
93	0	62	7	0	0	42	0	8

heparinized blood intermittently agitated the mean percentages of neutrophils containing brucellae in 4 to 12 tests for each dog were 96, 94.5, 99, and 34, whereas continuous agitation resulted in means of 100, 98.4, 99.5, 97, and 53, respectively. Ranges of variation for the former were 9, 16, 17, 3, and 51 and for the latter 0, 6, 2, 7, and 7, respectively. Phagocytosis was usually greater and variation less in rotated mixtures.

Effect of temperature on phagocytosis. Certain observers (Harmon *et al.*, 1946; Zarafonetis *et al.*, 1947) have claimed that opsonization was less at 37 C than at 22 C while phagocytosis was enhanced. However, these experiments were made in mixtures violently agitated, i.e., shaken 560 times a minute for 5 min-

utes, and it is questionable whether such conditions are suitable for quantitative demonstration of phagocytic activity (Fenn, 1923).

Observations were obtained from twelve aliquots of blood samples of three dogs, six aliquots at each temperature. The mean percentage of neutrophiles which contained bacteria was 98.4 at 37 C and 93.5 at 20 C. In the three series at 37 C the variations were 0, 4, and 8 per cent, while in the same subjects at 20 C the corresponding variations were 7, 17, and 7 per cent.

Effect of duration of agitation. The duration of phagocytic tests has varied in different investigations (Cottingham and Mills, 1943; Berry *et al.*, 1945) from 4 minutes to 60 minutes. Berry and Haller (1947) changed the 4 minute to a 10 minute period of agitation (method of Cottingham and Mills, 1943). Most observers have employed a 30 minute period.

It seemed necessary to establish the optimal time required for mixing. This optimal period would be the shortest one in which variation of phagocytosis was minimal. Maximal variations between 4 to 12 reduplicated observations at 30, 15, and 10 minute periods at 37 C were 7, 15, and 54 per cent, respectively.

Comparison between phagocytosis of living and formalin killed bacteria. Blood samples of 3 dogs were divided; 6 aliquots of each sample were tested with living and 6 with formalin killed bacteria. The bacteria were killed by treating with 1 per cent formaldehyde for 24 hours, then washed with sterile saline and stored at 5 C in vaccine flasks. The mean percentage of neutrophiles phagocytizing formalin killed brucellae was 98.6 in 18 tests, essentially the same as that phagocytizing living organisms, 97.6.

Tovar (1944) recently suggested using formalin killed cells of brucellae. The present experiments confirm those of Tovar with respect to phagocytosis of formalin killed brucellae.

Variation under standardized conditions. A series of experiments was performed to ascertain the variability under optimal conditions. Each of 10 samples of blood was tested six times. These samples were selected because at least 1 of 6 aliquots showed bacteria in 100 per cent of neutrophiles. The results are recorded in table 2. The mean percentage of neutrophiles that contained bacteria was 98.7. A statistical analysis of these data by Dr. Clifford Maloney is recorded in table 3. This analysis indicated that when 50 polymorphonuclear neutrophiles were examined in each test, bacteria would be found in 100 per cent of the neutrophiles in 53 per cent of the tests, in 98 to 100 per cent of the neutrophiles in 87 per cent of the tests, in 96 to 100 per cent of the neutrophiles in 97 per cent of the tests, and in 94 to 100 per cent of the neutrophiles in over 99 per cent of the tests. In other words, less than 1 per cent of the tests with bacteria in less than 94 per cent of the neutrophiles would have arisen from a population in which the mean was 98.7 per cent.

When the number of cells studied was below 33, small changes in the number of neutrophiles free of bacteria would produce large possible errors.

End points other than 94 to 100 per cent were more variable and required many more experiments for definition. Examples of such variability may be found in data obtained from the following observations. Blood sample 1 had 11

aliquots in which counts ranged from 94 to 100 per cent for neutrophiles which contained bacteria. Sample 2 had 10 observations ranging from 83 to 92 per cent, and sample 3 had 12 observations ranging from 30 to 57 per cent. Such

TABLE 2
*Phagocytosis of Brucella suis by polymorphonuclear neutrophiles**

DOG	TOTAL NEUTROPHILES EXAMINED	PER CENT WITH BACTERIA	DOG	TOTAL NEUTROPHILES EXAMINED	PER CENT WITH BACTERIA
1	37	92	2	61	97
	40	98		59	97
	53	100		72	99
	27	96		47	100
	24	96		49	98
	33	94		63	94
2	44	100	3	70	100
	65	100		58	98
	58	100		76	99
	56	96		54	100
	66	100		51	100
	77	100		74	100
3	65	100	1	48	96
	20	100		74	100
	29	100		75	99
	32	100		25	96
	28	100		43	98
	53	100		14	93
4	47	100	2	59	98
	66	100		58	100
	69	100		55	98
	68	100		42	100
	52	98		49	100
	68	100		48	98
5	62	100	6	53	96
	57	100		50	96
	68	100		77	96
	68	100		61	98
	85	100		59	98
	74	99		67	100

* Six tests with each sample

data show that variability was inversely related to the percentage of neutrophiles that phagocytosed brucellae.

Effect on phagocytosis of repeated washing of blood cells in an isotonic solution of electrolytes and gelatin. In attempting to distinguish between serum and cellular factors concerned with phagocytosis, it was necessary to obtain cells in a medium free of plasma constituents. Removal of plasma from cells required a

fluid suitable for washing without damaging blood cells. It is well recognized that isotonic electrolyte or crystalloid solutions are hemolytic when mixed with small numbers of cells. Addition of colloids greatly diminishes the hemolytic action of salt solutions. The purified gelatin (Knox P-20) was tested and found useful for this purpose. Krebs gelatin was prepared as described elsewhere (Victor, Wells *et al.*, 1952). The blood cells were washed 6 times, each time with a 10-fold volume of Krebs gelatin solution. After washing, the blood cells were resuspended in their own plasma. Phagocytic activity of washed blood cells was compared with that of untreated blood. Washing of dog blood cells 20 times with Krebs gelatin solution did not interfere with phagocytosis of living *B. suis*.

A method for quantitative determination of opsonin titer in plasma. The data derived from cited experiments formed the basis for a method measuring the opsonin titer of blood. Heparinized blood cells were separated from plasma by

TABLE 3

Probable incidence of tests with various cell counts and percentages of phagocytic neutrophils when the population had a mean of 98.7 per cent phagocytic neutrophils

NUMBER OF NEUTROPHILES WITHOUT BACTERIA	NUMBER OF NEUTROPHILES EXAMINED							
	14		25		33		50	
	Probable incidence of tests and phagocytic neutrophils							
	Test incidence	Per cent phago- cytosis	Test incidence	Per cent phago- cytosis	Test incidence	Per cent phago- cytosis	Test incidence	Per cent phago- cytosis
0	0.83410	100	0.72465	100	0.65268	100	0.52512	100
1	0.98569	93	0.95954	96	0.93338	97	0.86555	98
2	0.99846	86	0.99609	92	0.99140	94	0.97369	96
3	0.99911	79	0.99972	88	0.99917	91	0.99612	94
4			0.99998	84	0.99993	88	0.99954	92
5					0.99999	85	0.99995	90

centrifugation. The sedimented blood cells were washed 6 times in 10 volumes of Krebs gelatin solution. Plasma was diluted with Krebs gelatin solution. Washed blood cells were mixed with equal volumes of diluted plasma. *B. suis*, grown on tryptose agar slants for 24 hours, was treated at room temperature with 1 per cent formaldehyde in saline for 24 hours and then suspended in a concentration of 1×10^{10} per ml in Krebs gelatin solution. For the test, 0.1 ml of the blood cell plasma mixture was added to 0.1 ml of the bacterial suspension. Test tubes with 8 mm inside diameter were employed. A separate tube was employed for each concentration of plasma. The test tubes containing the mixtures of blood cells, plasma, and bacteria were agitated by rotation 24 or 36 times a minute at 37 C for 30 minutes. Smears of the mixture were prepared on glass slides. Since the viscosity of gelatin at room temperature is high, the preparation of smears was facilitated by using warmed slides. The extent of phagocytosis was determined as the percentage of 50 polymorphonuclear neutrophils containing bacteria. Since the variability of measurement was mini-

num in cases where the percentage of neutrophiles containing bacteria was 94 to 100, this range was selected as the end point for determining the opsonin titer. A unit of opsonin was defined as the quantity of opsonin causing phagocytosis by 94 to 100 per cent of neutrophiles under the previous conditions. The opsonin titer was expressed as the number of opsonin units in 0.1 ml of blood. Opsonin titer was determined from the greatest dilution of plasma in which 94 per cent or more of the neutrophiles contained bacteria under the conditions of the test.

In other experiments it was found that phagocytosis by dog leucocytes of *Brucella abortus*, strain 19, was the same as that of *B. suis*, strain PSIII.

Opsonin titer of normal dog plasma. The method was applied to the blood of 11 dogs. The results are recorded in table 4. The table gives percentages of neutrophiles which phagocytosed *B. abortus*, strain 19, in different concentrations

TABLE 4
Effect of plasma concentration upon phagocytosis

DOG	PER CENT PLASMA CONCENTRATION							OPSONIN TITER
	100	50	23	17	10	1	0	
	<i>Per cent phagocytic neutrophiles</i>							
1	100	71	40	6			5	1
2	100	86	73	22			13	1
3	92	72	19	16			12	0
4	96	81	30	9			30	1
5	5	11		3			13	0
6	100		84		10		10	1
7	91		37	30	11		6	0
8	47	23	6	15	9		0	0
9	52	7	7	7			8	0
10	38	13	8	22	21		1	0
11	96	71	50	6			6	1

of plasma as well as in Krebs gelatin solution without plasma. Five dogs showed 1 unit of opsonin in their blood. The blood of the other 6 dogs had less than 1 unit of opsonin in 0.1 ml of blood although all, except dog 5, showed more phagocytosis with than without plasma. Whether this effect of plasma was due to opsonin was not ascertained in these animals. It is noteworthy that the dogs with opsonin titers of 1 had been employed in the laboratory for repeated experiments for a year before these tests were performed. The six other dogs had been in the laboratory only on the single occasion when blood was obtained in the present test.

The application of this test to the blood of man and its relation to contact with brucellae and to brucellosis will be described in a subsequent report.

DISCUSSION

The experiments described previously indicated a variety of factors that influenced *in vitro* phagocytosis of *B. suis*, strain PSIII. These factors have been

investigated only with blood leucocytes and may not be the same with exudate leucocytes. The various factors that influenced phagocytosis have been combined in a method for measuring opsonins that lends itself to experimental and clinical application.

Attempts to count brucellae within leucocytes were unsuccessful. The organisms were so small that it was impossible to keep track of their position within the cell. In many instances the bacteria could not be distinguished from cytoplasmic granules. This was found with a large variety of stains including Hastings, various toluidine blue strains, Giemsa, Wright, carbol fuchsin, alkaline methylene blue, and finally, the Jenner-Giemsa stain. The latter stain was the most satisfactory for defining the organisms. Yet the best preparations were unsuited for accurate counts. Even in enlarged photomicrographs obtained with the phase microscope, it was impossible to distinguish some cytoplasmic granules from intracellular bacteria. For these reasons it was decided to quantitate phagocytosis from the percentage of cells that had phagocytosed the organisms. When 94 to 100 per cent of the neutrophils contained bacilli this method yielded unequivocal information from a population of neutrophils in which phagocytosis was intense and plainly evident.

Standardized conditions lead to uniform results in a sensitive system. With the present method, it was possible to measure opsonin in both normal and abnormal blood. In determining the opsonin level of plasma, especially in blood with high titers, it was mandatory to wash the cells free of plasma. Washing was readily accomplished with Krebs gelatin solution without impairing phagocytic activity of neutrophils. Ecker, Weisberger, and Pillemer (1942) employed 0.9 per cent NaCl solution for washing blood cells. This was adequate for five repeated washings. In the present studies, when it was desired to wash blood cells 20 times, hemolysis frequently occurred after three washings in Krebs solution. Incorporating 3 per cent gelatin in the medium prevented hemolysis.

The method developed from the present studies had certain features of those employed by Jersild (1941), Hald *et al.* (1943), and Ecker, Weisberger, and Pillemer (1942) in that the opsonin titer was determined from the phagocytic activity in various concentrations of plasma or serum. These methods had the same objective, namely, to obtain leucocytes free of opsonin. In other methods, opsonin titer was determined with phagocytes of one subject and serum or plasma of another. In the present method the phagocytes and plasma were from the same blood. Jersild (1941) and Hald *et al.* (1943) opsonized bacteria in serum and observed phagocytosis with citrated blood. In this way, they avoided the effect of citrate on opsonization but did not prevent its possible effect on phagocytosis. Ecker *et al.* (1942), Ecker, Pillemer, and Kuehn (1942), and Ecker and Lopez-Castro (1947) washed cells of defibrinated blood with saline solution. The phagocytic test of Jersild (1941) and Hald *et al.* (1943) was conducted with stationary mixtures, while Ecker's tests and our own were performed with agitation of the mixtures. Bacterial counts within leucocytes and percentage of phagocytic leucocytes were determined in the other tests, while only the latter was recorded in the present experiments, for reasons just dis-

cussed. This technique has been applied to human brucellosis (Victor, Pollack *et al.*, 1952).

These experiments were initiated to explain the bactericidal activity of dog blood, which has been found to depend on the combined activity of leucocytes and plasma (Victor, Wells *et al.*, 1952). Investigations on the relationship between phagocytic and bactericidal activity will be described in a later report.

SUMMARY

Factors influencing phagocytosis of *Brucella suis*, strain PSIII, by normal dog blood leucocytes *in vitro* were investigated. Neutrophils were more active than mononuclear or eosinophilic leucocytes. Citrate, unlike heparin, inhibits phagocytosis. Uniform phagocytosis occurred with rotation at 37 C for 30 minutes. Formalin killed organisms were phagocytosed to the same degree as living ones. Repeated washing of blood cells as frequently as 20 times in a modified Krebs solution containing 3 per cent gelatin did not impair phagocytic activity. On the basis of these findings, a method was developed for titrating opsonin using cells and plasma of the same blood.

REFERENCES

- BERRY, L. J., AND HALLER, E. C. 1947 Influence of anemia on phagocytic functions in rats. *Blood*, Special Issue, **1**, 108-116.
- BERRY, L. J., DAVIS, J., AND SPIES, T. D. 1945 Phagocytic activity of neutrophils in anemias. *J. Lab. Clin. Med.*, **30**, 910-916.
- CASTANEDA, M. R., TOVAR, R., AND VELEZ, R. 1942 Studies on brucellosis in Mexico. Comparative study of various diagnostic tests and classification of isolated bacteria. *J. Infectious Diseases*, **70**, 97-102.
- COTTINGHAM, E., AND MILLS, C. A. 1943 Influence of environmental temperature and vitamin-deficiency upon phagocytic functions. *J. Immunol.*, **47**, 493-502.
- DICKEY, J. W., JR., AND FORBUS, W. D. 1945 Chemotactic properties of *Brucella suis*; study of phagocytosis of *Brucella* in vitro by normal, nonimmune human leukocytes. *Am. J. Path.*, **21**, 195-203.
- ECKER, E. E., AND LOPEZ-CASTRO, G. 1947 Complement and opsonic activities of fresh human sera. *J. Immunol.*, **55**, 169-181.
- ECKER, E. E., PILLEMER, L., AND KUEHN, A. O. 1942 Opsonins of normal and immune sera; opsonins of sera of different species, role of complement in opsonic activity and combination of immune serum and normal serum as influencing opsonization. *J. Immunol.*, **43**, 245-258.
- ECKER, E. E., WEISBERGER, A. J., AND PILLEMER, L. 1942 Opsonins of normal and immune sera; methods. Comparison of effects of normal and immune opsonins on *Staphylococcus aureus*. *J. Immunol.*, **43**, 227-243.
- FENN, W. O. 1923 Phagocytosis of solid particles; carbon and quartz in solutions of varying activity. *J. Gen. Physiol.*, **5**, 311-325.
- HALD, A., JERSILD, M., AND RASCH, G. 1943 On determination of phagocytic power of leucocytes; study in economical design of experiments. *Acta path. et microbiol. Scand.*, **20**, 64.
- HARMON, D. R., ZARAFONETIS, C., AND CLARK, P. F. 1946 Temperature relations in phagocytosis. *J. Bact.*, **52**, 337-347.
- HUDDLESON, I. F., JOHNSON, H. W., AND HAMAAN, E. E. 1933 Study of opsono-cytophagic power of blood and allergic skin reaction in *Brucella* infection and immunity in man. *Am. J. Pub. Health*, **23**, 917-929.

- JERSILD, M. 1941 Cytophagic reaction employed in diagnosis of brucella infection. *J. Infectious Diseases*, **68**, 16-19.
- LUCKÉ, B., STRUMIA, M., MUDD, S., McCUTCHEON, M., AND MUDD, E. B. H. 1933 On comparative phagocytic activity of macrophages and polymorphonuclear leucocytes; essential similarity of tropin action with respect to two types of phagocytes. *J. Immunol.*, **24**, 455-491.
- MAXIMOW, A. A., AND BLOOM, W. 1948 Textbook of Histology. W. O. Saunders Co., Philadelphia, 5th ed., p. 52.
- OTANI, M. 1918 Acceleration of phagocytosis in citrated blood and citrated blood-plasma. *Kitasato Arch. Exptl. Med.*, **2**, 147-218.
- ROBERTSON, O. H., AND SIA, R. H. P. 1924 Studies on pneumococcus growth inhibition; method for demonstrating growth-inhibitory and bactericidal action of normal serum-leucocyte mixtures. *J. Exptl. Med.*, **39**, 219-244.
- SMITH, P. E., AND COPENHAVER, W. W. 1948 Bailey's Textbook of Histology. Williams and Wilkins, Co., Baltimore, 12th ed., p. 155.
- TOVAR, R. M. 1944 Opsonization test in brucellosis. *J. Immunol.*, **49**, 203-207.
- VICTOR, J., POLLACK, A. D., VALLIANT, J., AND RAYMOND, R. 1952 Human blood opsonin titers against *Brucella*. *J. Am. Med. Assoc.*, *In press*.
- VICTOR, J., WELLS, E. B., CLEMENTS, J., AND POLLACK, A. D. 1952 The bactericidal action of dog blood against *Brucella suis*. *J. Infectious Diseases*, *In press*.
- ZARAFONETIS, C., HARMON, D. R., AND CLARK, P. F. 1947 Influence of temperature upon opsonization and phagocytosis. *J. Bact.*, **53**, 343-349.