be rapidly restored by rewarming and inversion of the cylinder (Crawford *et al.*, 1967) or, more slowly, by storing the cylinders in a horizontal position at a temperature above 5° C. (Bracken *et al.*, 1968). If these precautions are taken the use of Entonox in the ambulance service is wholly justified as a simple and safe method of pain relief at a time when it is particularly important. The method of self-administration with a demand apparatus allows the patient freedom of choice and regulation of his relief. The use of the gas has raised morale in the ambulance service because, for the first time, ambulance personnel can do something specific to relieve pain. None of the men who have had experience in the use of Entonox would now want to be without it.

As a result of our experience in this survey the Gloucestershire County Health Committee has decided to extend the use of Entonox to the whole of the county ambulance fleet. We should also like to recommend that other ambulance authorities consider the use of Entonox.

We gladly record our thanks to Gloucestershire County Council for approving the survey and for providing the necessary equipment. We also have pleasure in paying tribute to Mr. A. Johnston,

the county ambulance officer, Mr. G. Turnbull, the deputy county ambulance officer, Mr. E. Lovell, the area superintendent, and to all the personnel at the Almondsbury, Wotton under Edge, Yate, and Soundwell Ambulance Stations. Appreciation is due to the casualty surgeons at the hospitals in Bristol and to Miss Gabrielle Routledge, who provided invaluable secretarial help. Without all their enthusiastic support and co-operation the survey would not have been possible.

REFERENCES

- Baskett, P. J. F., Hyland, J., Deane, M., and Wray, G. (1969). *British Journal of Anaesthesia*, **41**, 684.
 Bethune, D. W., and Collis, J. M. (1967). *Anaesthesia*, **22**, 43.
 Bracken, A., Broughton, G. B., and Hill, D. W. (1968). *British Medical Journal*, **3**, 715.
 Crawford, J. S., Ellis, D. B., Hill, D. W., and Payne, J. P. (1967). *British Medical Journal*, **2**, 138.
 Dundee, J. W., Brown, S. S., Hamilton, R. C., and McDowell, S. A. (1969). *Anaesthesia*, **24**, 52.
 Gale, C. W., Tunstall, M. E., and Wilton-Davies, C. C. (1964). *British Medical Journal*, **1**, 732.
 Lunn, J. N., and Kennedy, B. R. (1968). *British Medical Journal*, **2**, 828.
 MacGregor, W. G., (1967). *Nursing Mirror*, **124**, 23 June, p. i.
 Parbrook, G. D. (1967a). *British Journal of Anaesthesia*, **39**, 730.
 Parbrook, G. D. (1967b). *British Journal of Anaesthesia*, **39**, 974.
 Petrovsky, B. V., and Yefuni, S. N. (1965). *British Journal of Anaesthesia*, **37**, 42.

Mobile Unit for Screening for Cervical and Breast Cancer

MARY SEACOME,* M.A., B.M., B.CH.; HILDA K. PAINE†

British Medical Journal, 1970, **2**, 43-44

Summary: In 12 months' use of a mobile unit for cervical and breast cancer screening in Gloucestershire 3,211 women attended at an average of five sessions a week. Clinic sessions were organized and the running costs of the service met by a voluntary organization. The keeping of records, provision of laboratory facilities, and the follow-up of patients were carried out in close co-operation with the county health department.

Introduction

The mobile staywell clinic which took to the roads of Gloucestershire in October 1968 was provided and is maintained by a voluntary organization, the Gloucester City and County Appeal for the Prevention of Cervical Cancer. This article reports our experience in the first year of using the clinic in a programme of screening for cancer of the cervix and breast.

The initial impetus for forming the organization came from small pressure-groups of women who were dissatisfied with the facilities then provided for the detection of cervical and breast cancer. Initially the organization comprised representatives of the various women's organizations operating in this county. The major aspect of the organization's work comprises the general education of the public about cancer, emphasizing in particular the need for taking cervical smears and raising funds to provide this service. The latter aspect was stimulated by the suggestion that a mobile unit could serve the entire area and thus take the service to those who might otherwise find it inaccessible. From the outset the voluntary organization worked in close co-operation with the local authorities, and in particular with doctors from the health departments. Thus it was possible to plan a course of action which was satisfying to the volunteers and acceptable to the medical profession, and which also fulfilled a vital role in the total scheme for providing a cervical cytology service.

Planning of the Service

The general principle of any screening service for cervical cancer is that it must be comprehensive. For this reason, plans for providing facilities for taking smears, examination of slides, general-practitioner follow-up, gynaecological advice and treatment, and an alerting system for repeat investigations were laid well ahead to ensure that this total service should be available. General practitioners were told about the plans through the local medical committee and gynaecologists were consulted about various points. It is evidence of the good will of the pathologists that each of the pathological laboratories serving the area agreed to undertake the extra cytology work when the clinic went on the road. Though co-operation from medical colleagues was readily forthcoming, many were sceptical about whether women would be willing to use such a unit.

During this time public meetings were held over a period of about 12 months. At these a representative of the voluntary organization explained its aims and a medical officer of the local authority spoke about the medical aspects of screening for cervical cancer and about cancer in general. Local groups were also formed for the purpose of raising funds, disseminating information, seeking out volunteers, and helping with the organization of sessions.

Organization of Sessions

The keynote of the organization is to provide an efficient service on a sound economic basis. For this reason the unit visits a location only when there are enough women wishing to attend to make a full session of 20 patients. The local committees make the service known to the residents of their area, compile a list of patients, and inform the central secretary when a visit is needed.

The work of each session is carried out by a volunteer clerk and nurse, and a doctor who is either employed by the voluntary organization or seconded by the county health depart-

* Senior Medical Officer, Gloucestershire County Council.

† Chairman, Gloucester City and County Appeal for the Prevention of Cervical Cancer.

ment on a sessional basis. As well as taking the smear a full pelvic examination is carried out, the breasts are examined, and instruction is given in self-examination of the breasts. Where necessary patients are referred to their own doctors for further advice or treatment. All medical records are handled and filed by the county health department, a medical officer accepting the responsibility for liaison with general practitioners when further action is recommended by the pathologist. Follow-up smear examinations are carried out either by the general practitioner or by a local authority clinic, patients who leave a self-addressed envelope being informed of the results. Where further action is requested the patient is told about the result after consultation with the general practitioner, the health visitor often being asked to undertake this task.

Plan of the Unit

The unit consists of a 22 ft. (6.7 m.) caravan mounted on a close-coupled twin-wheeled chassis towed by a standard Land-Rover. It is divided into three compartments: a waiting-room, nurse's room, and medical room (see Fig.). The

Layout of mobile unit. Sliding doors ensure privacy in each section without loss of space.

waiting-room accommodates six people in comfort, which is ample space, as patients arrive by appointment. Record completion and sterilization of instruments take place in the middle compartment. The medical room measures about 8 by 6 ft. (2.4 by 1.8 m.), and once the doctor and nurse have become used to working together in a small space they find the layout practical. Power is provided for the unit by a generator carried in the Land-Rover. This supplies the lighting circuit and the sterilizer and also heats the water, of which the unit carries about 40 gallons (180 litres), an ample supply for most sessions. The effluent water is conducted by a pipe to the nearest drain. The caravan is heated by a Webasto oil-fired system. Where there is a convenient source of mains electricity, however, the unit can be plugged in and this used as an alternative. Nevertheless, the fact that the unit is independent of the mains supply makes it possible to visit isolated places.

Costs

A private firm donated the Land-Rover and trailer to the organization for conversion into a mobile cytology unit. For this reason it is impossible to quote actual costs for providing such a unit but only conversion costs. The sum spent on adapting the trailer from a shell on a chassis to a three-roomed clinic was £1,500, plus another £170 for modifications to the heating system. A generator was purchased for £113, and the cost of basic non-expendable medical equipment was £250 10s. The average running costs per five-session week are £67. This sum includes payments to doctors, driver, and central organization secretary and the payment of tax, insurance, petrol and oil for the Land-Rover, telephone, postage, etc., and disposable medical equipment. Depreciation of the unit is not included. The low costing is partly due to the fact that the clerks and nurses give their services voluntarily. All these running-costs are met by the voluntary organization, which

continues with the daunting task of fund-raising as well as organizing the clinic sessions.

Results of Screening

During the year 1 October 1968 to 1 October 1969 the unit unfortunately had to be taken off the road for two periods owing to teething troubles with the heating system, so the figures represent less than a full 12 months. The average number of sessions a week was five and a total of 3,211 women attended, of whom 1,578 were aged between 30 and 35 and 582 were under 30. The results of screening over the 12-month period are given in Table I. The patients with negative results are asked to reattend after an interval of three years.

TABLE I.—Results of Screening over 12-month Period

	No.
Total attendances	3,211
Smears taken	3,197
Breast examinations only	14
Positive smears showing malignancy	14
Repeat smears requested (excluding above)	164
Breast abnormalities referred to general practitioner	24
Other gynaecological conditions referred to general practitioner	148

Comment

The members of the voluntary organization are well aware of the need to encourage multiparous women and those of the lower socio-economic groups to attend for cervical screening. It is always easier to ensure the co-operation of the higher social groups and those with fewer commitments in these schemes. Nevertheless, the fact that the percentages of those attending according to social class are roughly comparable to the overall figures for the county is probably a measure of success in encouraging the lower socio-economic groups to attend (Table II). To make the services available to those in

TABLE II.—Analysis of Attendances of 3,211 Women by Class and Parity

	No.	%	% Resident in County (Working Males 1966 Census)
Social class I and II	895	28	22
" " III	1,430	45	48
" " IV and V	850	27	27
" " not recorded	36		3
Born after 1939 (most of this group were multiparous)	582		
No. with 3 children or more	974		

the priority groups local committees have offered both transport and baby-sitting services to would-be attenders. The unit has also been taken to several factories where the co-operation of the employers has made possible the screening of many women who otherwise would not have had the time or possibly the inclination to attend.

An unrecorded aspect of the work is the number of discussions about marital problems and contraceptive methods which have taken place. The atmosphere of the staywell clinic and the nature of the work conducted within it are conducive to the discussion of topics which many women normally hesitate to mention. Normally, patients with difficulties are advised to see their general practitioner or to visit the family-planning clinic. It would, however, be an advantage to have a more clearly developed follow-up system than this, since there is not time in the normal screening session to pay full attention to these incidental problems.

Though the unit has so far had medical officers of the local authority or doctors paid by the voluntary organization in attendance, it is hoped that eventually general practitioners will conduct sessions for their own patients in isolated areas or localities without a surgery or branch surgery.

It has been interesting to observe the changing attitude towards this unit from scepticism to recognition of the part that such an enterprise can play within the structure of the medical services. In an area that has always had reason to value the good relations existing between statutory and voluntary bodies it is exciting to discover a new field which can be explored.