of the area was performed, and a fistula was found communicating with the terminal ileum. She recovered from her gas-gangrene infection and one month later had a right hemicolectomy performed for Crohn's disease, which was confirmed histologically in 25 cm. of the terminal ileum. She has since made a good recovery and has had no more fistulae.

A 26-year-old woman had been suffering with Crohn's disease confined to the terminal ileum for three years when seen in 1965. She became pregnant, and at four months developed pain in the right hypochondrium radiating round her right lower ribs to the back. She was tender in this area but no other abnormality was found. The pain continued to trouble her throughout her pregnancy. At term a healthy child was born. One month later the pain suddenly became more severe and was associated with sweating and rigors. She was treated with antibiotics and a mass appeared in the right hypochondrium. Seven weeks after delivery a liver abscess was drained. The pus was bacteriologically sterile. Subsequently she made a good recovery and remains well, apart from residual thickening of the terminal ileum.

The clinical course of Crohn's disease is often difficult to forecast. This also applies equally to the disease during pregnancy. Crohn¹ took the view that if the onset of the disease occurred during pregnancy then its subsequent course was severe. Our first case suffering from gas gangrene as a result of Crohn's disease would support this view. The second case, who was under steroid therapy for the whole duration of her Crohn's disease, demonstrates that complications occurring during pregnancy can be exacerbated during the puerperium.

We wish to thank Mr. H. M. Goldberg, under whose care these patients were, for permission to report these cases.—We are, etc.,

B. HEYWORTH.
S. BASU.
J. CLEGG.

Crumpsall Hospital,
Manchester.

REFERENCE

- ¹ Crohn, B. B., Yarnis, H., and Korelitz, B. I., *Gastroenterology*, 1956, 31, 615.

SIR,—Dr. J. F. Fielding and Dr. W. T. Cooke (11 April, p. 76) in their interesting article on pregnancy and Crohn's disease point out that pregnancy has little effect on the course of the disease but observe a decrease in fertility in patients with Crohn's disease. From observations in patients attending the Cleveland Clinic, Ohio, we would agree with these two points, but our experience with regard to problems with the fetus has not been as happy as that of the Birmingham workers. In our series there appears to be an increased risk to the fetus. It is difficult to be certain of all early abortions so we have included only pregnancies going beyond the first trimester.

Number of pregnancies in Crohn's disease	34
Stillbirths or spontaneous abortion after the first trimester	8
Premature births	2
Abnormal fetus	2

There is a marked difference between Drs. Fielding and Cooke's patients and those which we are reporting here in that the majority of the former had small bowel Crohn's disease and the majority of the latter had large bowel Crohn's disease.

However, we feel that the figure of 12 abnormalities in 34 well-established pregnancies is sufficiently great to sound a note of caution. Incidentally, we have observed only two stillbirths in 31 pregnancies in patients with total colonic involvement with mucosal ulcerative colitis.

We would speculate that the severity of the illness is a factor which may lead to problems with the fetus, and the difference in distribution of the lesions in these two reports of pregnancy in Crohn's disease may well be a significant factor accounting for the observed differences.—We are, etc.,

PHILIP F. SCHOFIELD.

General Hospital,
Ashton under Lyne, Lancs.

R. B. TURNBULL.
W. A. HAWK.

Cleveland Clinic,
Ohio, U.S.A.

Campbell de Morgan Spots

SIR,—Dr. H. P. Ferrer in his letter (7 March, p. 628) suggests that the outbreak of Campbell de Morgan spots was of food-borne origin. This was considered very carefully and ultimately discounted when it became apparent that controls in Liverpool had the same incidence of lesions, and no foods common to all patients could be found. The outbreak was certainly not confined to institutions, as he suggests, for our outpatients here and in Westmorland were similarly affected. A leading article in the *Lancet* (18 April, p. 824) mentions an autumn outbreak in Cumberland, but this also followed an increase in ambient temperature.

One of our reasons for describing the outbreak was that others, knowing its benign nature, might then be spared alarmist press publicity.—We are, etc.,

R. H. SEVILLE.
G. BIRCHALL.

Beaumont Hospital,
Lancaster.

General-practitioner Anaesthetists

SIR,—I have been most interested in the "Unheard Voices" series, especially that of the consultant anaesthetist (4 April, p. 45). As such a one of 18 years standing, 12 years in the North and subsequently in the South, I must most vehemently disagree with Dr. Cadwallader's adamant rejection of part-time help from general practitioners.

I am motivated to write this rebuttal by my deep indebtedness to the general-practitioner anaesthetists without whose help down the years life as a provincial consultant would have been the stuff of nightmares. Many of the doctors in question reached registrar standard as full-time anaesthetists before taking up general practice. One subsequently reversed the process, took his Fellowship, and became full-time consultant.

Those personally trained required close supervision over many months. The eventual result is a safe and efficient team with

a permanent career interest in hospital work. No one is exploited, and the heavy emergency load, inevitable in the specialty both during the week and over the weekend, is spread sufficiently thinly to be painless.—I am, etc.,

D. F. REES.

Ventnor,
Isle of Wight.

Drug Neutropenia

SIR,—Mohan¹ reported a patient who bled severely because of platelet deficiency due to the drug Septrin. During the past three months I have seen two women and one man with dangerous blood dyscrasias closely related in time to taking Septrin (trimethoprim and sulphamethoxazole).

The first patient, aged 61, was seen at home on 28 December, 1969 with bronchopneumonia which had been treated with Septrin 1 tab. t.d.s. for four days, and ampicillin for one day. He was deteriorating rapidly with a fever of 104.6° F. (40° C.), and on admission to hospital was found to have only 1,800 W.B.C./cu.mm. and two days later 1,600 W.B.C./cu.mm. and platelets 70,000/cu.mm. By 6 January his W.B.C. and platelets had risen to the lower limits of normal, and his pneumonia had responded to large doses of penicillin and prednisone.

The second patient aged 63, had received two courses of Septrin tabs. 2 b.d. from 19-26 January, and 1-12 February for P.U.O. eventually diagnosed as subacute bacterial endocarditis. Four days after the end of the second course she was admitted to hospital because she was less well and was found to have no granulocytes at all, whereas the W.B.C. had been normal with a normal differential on 2 February. However, retrospectively, mild neutropenia had begun to show on 10 February with 54% of 4,200 cu.mm. Sternal puncture on 17 February showed a highly cellular marrow; myelocytes and metamyelocytes but no neutrophils—for example, a maturation defect. Serum folates were normal 340 µg./ml. She recovered, and the blood picture was normal by 27 February, eleven days after her agranulocytosis.

The third patient, aged 74, had received eight days Septrin tabs. 2 b.d. for a urinary tract infection when she developed extensive purpura on 5 April. Her platelet count was 18,000 cu.mm. and W.B.C. 4,500 cu.mm. Septrin was stopped, and the patient admitted to hospital. Fortunately no further bleeding occurred and her platelets returned to normal within eight days.

We are assured that these complications are rare with Septrin, but the quoted experience in so short a time leads one to question this, and to ask how often patients already debilitated are having to fight off pyogenic chest infections, or perhaps failing to do so, with a neutropenia of which no one is aware.—I am, etc.,

J. W. PAULLEY.

Ipswich and East Suffolk Hospital,
Ipswich.

REFERENCE

- ¹ Mothan, P., *Practitioner*, 1969, 202, 553.

Smoking and Bronchitis

SIR,—In your leading article on bronchitis (25 April, p. 190) you state that "preventive measures are unlikely to have much impact on the frequency of chronic bronchitis in adults unless ways and means are found of drastically reducing the cigarette-smoking habits of the population."

One obvious preventive measure is to try to stop children from smoking, and there is