Patient	Dosage of Clindamycin	Time of Specimens	Serum level	Sputum level
1 ...	300 mg.	2 hours	12.0 µg./ml.	4.025 µg./ml.
2 ...	"	"	7.6 µg./ml.	7.0 µg./ml.
3 ...	"	"	Not done	0.91 µg./ml.
4 ...	"	"	3.1 µg./ml.	2.1 µg./ml.
5 ...	"	"	Not done	2.2 µg./ml.

It would seem therefore that serum and sputum levels can be achieved sufficient to inhibit *H. influenzae*.—I am, etc.,

ARTHUR A. B. MITCHELL.
Department of Bacteriology,
Law Hospital,
Carlisle, Lanarks.

Another Paraquat Fatality

SIR,—We wish to record a further instance of fatal paraquat poisoning. So far as we can ascertain, this is the tenth reported fatality caused by this widely used herbicidal preparation.¹⁻⁴ In December 1965 a 42-year-old male accidentally drank a small quantity of "weed-killer" from an unlabelled stout bottle. It was later estimated that the quantity of fluid consumed was about three-quarters of a teaspoonful. The nature of the substance could not be immediately ascertained, but it was subsequently identified as Gramaxone 19% w/w paraquat ion) by the laboratories of I.C.I. Ltd. On admission to hospital a few hours after taking the poison, the patient complained of a burning sensation in the throat and repeated vomiting. Despite immediate gastric lavage, prophylactic penicillin, and abundant oral and intravenous fluids, there was progressive clinical deterioration. At three days the tongue was severely ulcerated and two days later haemoptysis occurred. On the eighth day jaundice, pyrexia, tachycardia, and a blood urea of 375 mg./100 ml. were recorded.

The patient died some hours later. The outstanding necropsy findings included focal myocardial necrosis, severe pulmonary haemorrhage and oedema, dense eosinophilic alveolar membrane, and proliferation of fibroblast-like cells in the alveolar septa. It is apparent from this and other reports that the lethal dose of paraquat may be very small. Further fatalities will only be avoided by proper care in the storage and use of this valuable but potentially dangerous compound. A detailed and illustrated clinico-pathological report is being prepared for publication.—We are, etc.,

Department of Pathology, J. G. MASTERSON.
University College,
Dublin 2.
St. Columcille's Hospital, W. J. ROCHE.
Loughlinstown,
Co. Dublin.

REFERENCES

- 1 Almog, Ch., and Tal, E., *British Medical Journal*, 1967, 3, 721.
- 2 Bullivant, C. M., *British Medical Journal*, 1966, 1, 1272.
- 3 Campbell, S., *Lancet*, 1968, 1, 144.
- 4 Oreopoulos, D. G., *et al.*, *British Medical Journal*, 1968, 1, 749.

Drugs in Depressive Illness

SIR,—I was interested to read the article by Dr. A. M. Porter (28 March, p. 773). In his discussion he concluded that his work provided evidence to suggest that the milder forms of depression may be effectively treated with support and a placebo. One of the difficulties he mentioned in carrying out a drug trial with placebo was the easy breach of the cipher when drugs like imipramine were used because of their side-effects, and suggested the possibility of adding atropine to the placebo tablets to avoid this difficulty. This possibility has been discussed previously in a critical view of antidepressant drug trials.¹ This review also mentioned that most depressed patients get better anyway, and the patients who improve after being prescribed tablets have done *post hoc*, but not necessarily *propter hoc*.

In a double-blind drug trial carried out in this hospital for which preliminary results are now available, 54 patients entered the trial and of these 15 were treated with a tablet containing amitriptyline and perphenazine (Triptafen), 20 with perphenazine (Fentazin), and 19 with a placebo containing atropine. Each of the patients was randomly allocated to one of the treatments and assessed on three occasions in terms of 15 symptoms plus an overall rating. The trial and its analysis are still continuing.

Degree of Improvement	Triptafen	Fentazin	Placebo
Worse	2	1	5
Unchanged	2	4	3
Improved	0	0	4
Much Improved	1	4	5
Recovered	10	11	2

χ^2 (2 d.f.) = 1.52

χ^2 was evaluated after pooling the "worse" and "unchanged" categories.

The Table shows the distribution of patients according to their degree of improvement at the final assessment. A statistical analysis shows that $\chi^2 = 1.52$, implying a lack of significant difference between the various treatments. Although these patients were from hospital practice, the results are certainly very similar to those of Dr. Porter, and of Hunter *et al.*², and would support the view that antidepressant drugs tend to become established in clinical practice on inadequate evidence.—I am, etc.,

ZULFIQUAR HUSSAIN.

Cherry Knowle Hospital,
Sunderland,
Co. Durham.

REFERENCES

- 1 Leyburn, P., *Lancet*, 1967, 2, 1135.
- 2 Hunter, J. W., Owen, S., Eksi, A., and Scott, J. F., *British Journal of Psychiatry*, 1967, 113, 667.

Blue Light and Jaundice

SIR,—Inasmuch as I am an "enthusiastic" supporter of phototherapy, I feel that I should comment on the ultra-conservative editorial comment, "Blue Light and Jaundice" (4 April, p. 5).

Workers in this field^{1,2} now recognize that bilirubin can produce learning difficulties in humans at lower serum concentrations than were previously realized. Previous studies have not detected these defects because of inadequate testing. It is for this reason that a safe and simple method to replace exchange transfusion is needed. The statement that phototherapy is "unlikely to be adequate treatment" in preventing low bilirubin kernicterus is an opinion with which I certainly do not agree. Preliminary data suggest that this is the type of infant who will benefit. Phototherapy is also now widely being used to control mild haemolytic disease. There has been no convincing evidence published to support the original suggestion by Franklin³ that the photochemical products might be toxic. There is considerable evidence that these products are not toxic either in vivo, in vitro, and in human infants.⁴ I would ask that the author of this leading article and others read the Bilirubin Symposium⁴ quoted, and not just the summary as has apparently been done, for more data on this important point. I would also refer the reader to two recent articles,⁵ for more reassuring evidence in human infants. Published human clinical experience for several countries with phototherapy now numbers over 5,000 cases, and no serious toxic effects have been reported to date.⁷ More recent studies in two human infants with Crigler-Najjar syndrome employing ¹⁴C labelled bilirubin and phototherapy indicate that the breakdown products are rapidly and safely excreted.⁸

Early feeding may have limited the incidence of hyperbilirubinemia of prematurity in one nursery in England to less than 1%, but this has certainly not been the experience in the U.S.A. Where this problem has been studied carefully, one usually finds that from 10-25% of premature infants can be expected to develop serum bilirubin concentrations of over 15 mg./100 ml.⁷

The leader writer also cites a reference⁹ referring to a hypothesis that light might cause retrolental fibroplasia. This curious theory is certainly not widely held by many paediatricians.

I, and others, have not used "blue" light, but have employed broad spectrum light. One should not of course use "blue light," especially if you want to judge skin colour. Phototherapy units are now designed so that they not only allow better observation of the infant but also free access to the infant. The leader writer's quaint reluctance to allow even a minor change in the present lighting conditions in the nursery suggests that he knows that these conditions are indeed optimal for newborn infants. I challenge this idea. I am afraid that few doctors actually have any idea about the present lighting conditions in their nursery. Outdoor light on a bright day supplies 10,000 foot candles of broad spectrum light. I have found nurseries with 25 foot candles to 2,000 foot candles in the same nursery. Phototherapy units supply only about 300-500 foot candles of light. "Phototherapy" is,

therefore, being carried out in many nurseries without physicians being aware of it. There are no data whatsoever to support the present "dim light" conditions present in many nurseries as being the best. Outdoor lighting is used to illuminate many nurseries and "long-term disadvantages" are not apparent.

I agree that we have much to learn about light and its effect on human infants. The present leading article implies, however, that we know the present highly artificial light conditions are good, so why change? I would suggest that we face our ignorance on this point, examine the currently available evidence without prejudice, and then make a judgement.—I am, etc.,

JEROLD LUCEY.

Department of Pediatrics,
University of Vermont,
Burlington, Vermont, U.S.A.

REFERENCES

- 1 Odell, G. B., Storey, G. N., and Rosenberg, L. A., *Journal of Pediatrics*, 1970, **76**, 12.
- 2 Johnson, L., *Proceedings of the Society for Pediatric Research*, 107 1970, Atlantic City N.J.
- 3 Franklin, A. W., 1958, *Lancet*, **1**, 1227.
- 4 Hsia, D. Y., editor, *Symposium on Bilirubin Metabolism*, Original Article Series—National Foundation, New York City, 1970, (in press).
- 5 Gorodischer, R., Levy, G., Krasner, J., and Yaffe, S. J., *New England Journal of Medicine* 1970, **282**, 375.
- 6 Karon, M., Imach, D., and Schwartz, A., *New England Journal of Medicine*, 1970, **282**, 377.
- 7 Lucey, J. F., *Symposium on Bilirubin Metabolism*, Original Article Series—National Foundation, 1970 (in press).
- 8 Callahan E., et al., *Proceedings of American Society for Study of Liver Disease Meeting*, November 1969, Chicago, Ill, U.S.A.
- 9 Riley, P. A., and Slater, T. F., *Lancet*, 1969, **2**, 265.

*We agree with Professor Lucey that the most important question in deciding the possible value of phototherapy is whether bilirubin may produce brain damage at lower concentrations than those normally considered safe. We briefly reviewed the available evidence which suggests that serum bilirubin levels below 20-25 mg./100 ml. do not cause brain damage. Professor Lucey dismisses these studies on the grounds that testing was "inadequate" yet in two of them the children were followed up to the age of seven years, and in two cognitive disorders or "minimal cerebral dysfunction" were specifically looked for. The question is intrinsically a difficult one to answer, because the premature babies who are most likely to be jaundiced are also those most likely to suffer other possibly brain damaging circumstances. Professor Lucey appears to discount any possible ill-effects on the grounds that phototherapy illuminates an infant less brightly than would outdoor light on a bright day — but the natural environment for a premature baby is presumably total darkness. Professor Lucey is surely right in pointing out that too little attention has been paid to illumination as an aspect of the premature infant's environment. Nevertheless whatever variations there may be in present practice in this respect, phototherapy must mean an increase in illumination: otherwise there would be no point in discussing it. One of the papers Professor Lucey quotes,⁵ and an accompanying editorial (in the same journal), discuss the biological potency of light in much the same terms as we did, and recommend caution in its use.

We agree with Drs. Behrman and Hsia¹

that "Pediatricians should consider phototherapy in the same cautious manner as they would in deciding upon the use of a new drug which has become available for the treatment of newborn infants." Conservatism in neonatal care is not altogether to be condemned, remembering the various forms of treatment which have been advocated and widely accepted for the newborn, and have subsequently proved harmful. The research now in progress will make it easier to assess the proper place of phototherapy in neonatal care. Meanwhile, we believe our recommendations are reasonable for the paediatrician concerned with routine care as opposed to research.—ED., B.M.J.

REFERENCE

- 1 Behrman, R. E., and Hsia, D. Y. Y., *Journal of Pediatrics*, 1969, **75**, 718.

Amblyopic Eyes

SIR,—I would like to compliment Mr. D. H. Nixseaman on his useful article on amblyopia (2 May, p. 279). I do feel, however, that he takes rather a pessimistic view of the liaison between ophthalmologist and practitioner.

In the area served by my hospital, it is very much the exception for a case of squint to be referred with a history of squint for longer than six weeks. In consequence of this very little occlusion is required in my orthoptic department, and the aim now is to obtain a binocular functional result rather than only a cosmetic result. I would also mention the Sheridan Gardiner test, using letter symbols, as a very useful way of obtaining an accurate visual acuity, and in fact I would expect to have an accurate assessment of acuity by this means in most of my patients by three years of age.

With this early referral of cases, even when amblyopia is present, conventional occlusion produces good results, and there is no recourse to pleoptics.—I am, etc.,

G. V. CATFORD.

St. George's Hospital,
London S.W.17.

Entonox as a Psychotherapeutic Aid

SIR,—Premixed 50% oxygen and 50% nitrous oxide originally introduced for obstetric use¹ has been found to have an application in many other fields where safe analgesia is required. Part of its function in obstetrics is probably due to its effect on the anxiety and tension factors in the vicious circle of pain in parturition. Drs. P. J. F. Baskett and A. Withnell (4 April, p. 41) refer to its anxiety-relieving properties

in a sample of patients with a number of different painful conditions conveyed by ambulance.

The disadvantage of most of the agents used as psychotherapeutic aids, whether administered intravenously or inhaled, is that their effects are not quickly reversible. Trichlorethylene using the Tekota apparatus² proved to be much less satisfactory than Entonox and had an obvious odour.

A trial with patients suffering from various psychoneurotic disorders has shown its usefulness. As might be expected its effects in low dosage were very variable and depended on the expectation of the patient, suggestion, and the doctor-patient relationship at that time. Even after inhaling the mixture to the point of unconsciousness the patients recovered very rapidly and were able to go home unaccompanied. It was found to be of particular value in those patients whose tension and inhibition made verbal psychotherapy very difficult. The patient was able to get used to the fear of loss of control by holding the mask himself and then allowing the psychotherapist to do so. It also had an application in desensitization techniques for phobic and obsessional patients.—I am, etc.,

IAN J. MACDONALD.

London W.1.

REFERENCES

- 1 Gale, C. W., Tunstall, M. E., and Wilton-Davies, C. C., *British Medical Journal*, 1964, **1**, 732.
- 2 Ellis, M. W., and Bryce-Smith, R., *British Medical Journal*, 1965, **2**, 1412.

Colour in Rhodesia

SIR,—Your correspondents from Rhodesia, Mr. W. Houston and Dr. R. F. Browne (9 May, p. 366), are devoted to the principle of the free choice of doctor. Would they, I wonder, allow patients to reject the house surgeon on the grounds that he had hazel eyes, or that he had no turnups on his trousers?

Sweet reason and a sense of proportion are essential to all organized activities of men, and hospitals are no exception. Those who consider that objection to a doctor on grounds of race comes within the bounds of reason, and reflects a legitimate exercise of free choice of doctor, must in some measure share the racial prejudices of the patients concerned.

Hospitals should be considerate to the legitimate and reasonable needs and wishes of the patients. But they have an important task to perform on the basis of teamwork, and the whims and quirks of the neurotic women described by Dr. Browne should be patiently overruled.—I am, etc.,

M. J. CLAY.

Bishop Auckland,
Co. Durham.

Extra Duty Allowances

SIR,—The hospital junior staff representatives on the negotiating subcommittee of the Central Committee for Hospital Medical Services will soon have the duty of examining the awaited report of the Review Body, and of deciding with our senior colleagues whether to recommend its acceptance. An integral part of these discussions will be a re-examination of the existing agreement on

extra duty and acting-up allowances. The "pricing" of units will not be known until the recommendations of the Review Body are published.

A good deal of concern has been expressed that extra duty allowances are unprofessional and should be rejected—reliance instead being placed upon a claim for a substantial increase in basic salary.