than to insulin deficiency. This view does not now seem to be a correct one. When the blood glucose is raised in such patients there may be an excessive secretion of insulin. Nevertheless, these levels are still less than those observable in normal people, in whom the blood glucose is raised to comparable heights.²⁸ In addition, the secretion of insulin in such patients may show a delay.

These patients show insulin antagonism, which seems to be directly associated with their obesity; thus insulin is less effective in non-diabetic obese subjects than in normal people. Similarly, animals made experimentally obese may show a resistance to the action of insulin.^{29, 30} The non-diabetic obese person is, however, able to compensate for this insulin resistance by secreting extra insulin; in the mild diabetic adequate compensation does not seem possible, presumably because of some β cell defect.

The nature of the insulin antagonism that accompanies obesity is the subject of dispute. Both circulating antagonists,³¹ as well as a tissue antagonism based on a deranged fatty acid metabolism,³² have been postulated to account for it. Carbohydrate restriction leading to weight reduction may restore both glucose tolerance and insulin secretion virtually to normal in these patients, emphasizing that in the absence of obesity the β cells of such patients may function adequately.

Other Hormones and the Islets of Langerhans

Both growth hormone and the structurally similar material derived from the placenta, placental lactogen, appear to affect the islets of Langerhans. In man diabetes quite commonly accompanies acromegaly. Excess growth hormone has long been known to induce diabetes in some animals,³³ partly through a direct effect on the islets and partly by increasing resistance to insulin. Nevertheless, smaller quantities of growth hormone may raise blood insulin, probably by a direct effect on the β cells. This effect which takes some hours to develop may involve a sensitization of the islets to the secre-

tory effects of glucose.³⁴ Growth hormone also affects the β cells themselves directly to increase insulin synthesis. The extra responsiveness of the β cells may be especially important during growth, since insulin is important in increasing protein synthesis at this time.

Similar changes may occur during pregnancy, probably as a result of the secretion of the hormone placental lactogen, which closely resembles growth hormone in structure and properties. The effects of this substance may underly the increased insulin secretion which is known to accompany pregnancy. The alterations in metabolism in pregnancy may produce a temporary diabetic state—gestational diabetes, which disappears after delivery.

Insulin Secretion in Potential Diabetes

In the nomenclature of the British Diabetic Association the term potential diabetic is restricted to those who are particularly liable to develop diabetes. Ideally this category includes the identical twins of a known diabetic. As a group, unaffected identical twins of diabetics secrete less insulin in response to glucose than do normal people, either as a result of an intravenous glucose tolerance test³⁵ or after oral glucose,³⁶ though there may be considerable individual variation.³⁷ The delay in insulin secretion in this group may be due to the ineffectiveness of a system involving cyclic AMP, which promotes insulin release.

In summary, diabetes is often associated with some defect within the β cell, leading to an impairment of insulin secretion. This may range from complete β cell destruction to relatively minor changes in secretory response. Factors such as obesity or pregnancy in which there is an increased need for insulin may accentuate the abnormality. The discovery of a low insulin response to glucose among close relatives of diabetics emphasizes the possibility of inherited defects in the secretory mechanism for insulin. It is, however, much too early to be certain of their true significance in the aetiology of most cases of clinical diabetes.

General Treatment of Diabetes

D. A. PYKE,* M.D., F.R.C.P.

British Medical Journal, 1970, 3, 268-269

In a month or two it will be 50 years since Banting and Best started their work leading to the discovery of insulin. It seemed then that the problem of treatment of diabetes had been solved; it has dawned on us gradually over the years that it has not. Indeed, not only have advances in treatment been slow, there is now even a suggestion that a well-tried oral agent—tolbutamide—might be harmful (the evidence presented to the American Diabetes Association on 14 June³⁸—but not yet published in a medical journal—that patients on tolbutamide show an increased late incidence of arterial disease will need to be carefully scrutinized).

New Insulins

Recently two new types of insulin³⁹ have been introduced. The first is a neutral soluble insulin (Nuso in the United Kingdom, Actrapid in Denmark). Ordinary soluble insulin is

strongly acid and when mixed with neutral insulins produces a change in pH which affects, to a variable degree, the time-effect of the insulin mixture. It has therefore been unsatisfactory to mix soluble with the lente insulins, but it is now possible in those patients on lente insulin who show day-time hyperglycaemia and glycosuria and night-time hypoglycaemia to mix Nuso or Actrapid with lente and often obtain good control. The presumed advantage of Nuso and Actrapid that, being neutral, they are more quickly absorbed, is not of practical significance.

The second new type, "biphasic" (Rapitard) insulin consists of a mixture of neutral soluble and Crystal II insulin. It was hoped that this insulin might provide effective 24-hour control of blood sugar with a single injection. In practice this is seldom the case and usually biphasic insulin has to be given twice daily. Actrapid insulin, being pork insulin, is useful in the rare cases of species insulin resistance.

In spite of these new insulins there has recently been little real change in insulin therapy. Indeed, there has been a turning back to two-injection regimens (soluble insulin alone, or

* Physician, King's College Hospital, London S.E.5.

soluble insulin and N.P.H. b.d.). If one could choose only one type of insulin for use in all circumstances soluble insulin, introduced to medicine nearly half a century ago, would certainly be the choice.

Oral Treatment

Sulphonylurea treatment has been used for fifteen years and there are now six compounds to choose from. They differ from each other in time of action, production of flushing after alcohol, and cost. Since these drugs can cause hypoglycaemia—especially in the elderly—it is often best to prescribe a short-acting drug for patients over the age of 60 who live alone. Chlorpropamide causes unpleasant and embarrassing facial flushing after alcohol in a few patients. The other sulphonylureas rarely do so and tolanae is a suitable alternative, since the dose and frequency of administration are similar.

A new sulphonylurea, glibenclamide (Daonil or Euglucon) has recently been introduced and great advantages claimed for it, since it is a much more potent hypoglycaemic agent than the others. Besides being effective in a daily dose of only 5 mg., it is said to be less toxic than the other preparations. As against this, all sulphonylureas are very safe drugs and glibenclamide has not yet been on the market for long enough to know whether it does in fact have any advantage in this respect. That the dose is small is irrelevant. The main criterion by which a new oral antidiabetic agent should be judged is whether it increases the *scope* of oral therapy. The main disadvantage of glibenclamide is its cost and I can see no present indication for its routine clinical use.

Biguanides.—Phenformin and metformin are, as far as can be judged, equally effective. The place of these drugs in treatment is often in conjunction with a sulphonylurea, when they are undoubtedly effective in some cases. Many physicians find them valuable aids to weight reduction in obese diabetics. The mechanism and extent of their weight-losing effect are uncertain, and, as with all remedies for obesity, they are most effective when used by the enthusiastic and determined physician on the acquiescent and persevering patient. Both of these biguanides cause alimentary side-effects in some patients, but fewer with the slow-release preparations.

Diet

The only important recent advance in the dietary treatment of diabetes is that it has now been widely agreed that diets should be reckoned in 10 g. carbohydrate portions and that, except in the obese, fat and protein need not be restricted. Control of diabetes, even in the obese, is often achieved by carbohydrate restriction alone, before there is any weight reduction. Though it may be a useful encouragement to a fat patient to lose weight that this will control his diabetes, this is in fact seldom essential; the important measure is carbohydrate restriction.

Diabetic Coma

In the past few years a form of diabetic coma has been recognized with increasing frequency—diabetic coma without ketoacidosis or ketotic diabetic coma.⁴⁰ Though first described nearly a century ago, only in the last decade have many cases been encountered and it is now a common clinical entity. Features are, firstly, a high average age; secondly, it often appears in a previously unknown diabetic; and, thirdly, there is an apparently increased susceptibility of West Indian immigrants. The patients are extremely dehydrated but not overbreathing; there is no smell of acetone on the breath and

little or no ketonuria. Plasma ketone levels are raised but they are much lower than in keto-acidotic coma; blood sugar values are even higher, often exceeding 1,000 mg. per 100 ml.; serum electrolyte concentrations are usually normal, though the sodium may be raised; bicarbonate is only moderately depressed; urea is raised to about 100 mg. per 100 ml., but may go much higher if serious oliguria develops; the mean plasma osmolality is about 370 mOsm per litre.

Treatment consists of rapid fluid replacement, using hypotonic solutions such as half or even one-fifth strength saline, and insulin in relatively small doses (say, about one-half those used in keto-acidotic coma). In spite of reports of high mortality, recovery is the rule, though the elderly patient who is unconscious is at considerable risk. After recovery at least half these patients can be treated without insulin, which is rare after keto-acidotic coma.

The *cause* of diabetic coma without keto-acidosis is not known. Presumably the very high blood glucose levels are related to the relative lack of ketosis, but why they should be so high is unclear. Unlike cases of keto-acidotic coma serum insulin is present, but the amount is low in relation to the blood glucose level.

Lactic Acidosis

In a few diabetics acidosis may develop not from an increase of ketoacids but from lactic acid.⁴¹ The condition is not confined to diabetics and is usually associated with some serious underlying illness, and the significance of diabetes in its pathogenesis is uncertain. Some degree of lactic acidosis is fairly common in diabetic keto-acidosis and it may increase during the first few hours of treatment, but is seldom of importance.

Cases have been described of lactic acidosis developing on phenformin treatment, usually when the patient also had another serious illness. The role of phenformin in causing lactic acidosis, if any, and the method by which it does so are unknown. It is certainly not a common event.

Long-term Complications

Eyes.—Pituitary ablation in the treatment of certain forms of diabetic retinopathy has now been in use for about 15 years, yet its place in therapy is still not fully established. The difficulty derives from lack of fully controlled long-term therapeutic trials and from the variable progress of diabetic eye disease. Retinopathy with new-vessel formation and vitreous and retinal haemorrhages may benefit, but retinal exudates and retinitis proliferans are not affected. Renal involvement and other complications may be contraindications. Patients are, of course, dependent on hormone replacement therapy for the rest of their lives.

Photocoagulation in the treatment of new retinal vessel formation has recently been widely practised. The procedure is simple and, properly applied, has no appreciable ill effects, which may explain why many ophthalmologists are enthusiastic about it. However, again because of the unpredictability of diabetic eye complications, the only way of assessing the value of photocoagulation is by controlled trial. A multicentre study has been organized by the medical and scientific section of the British Diabetic Association; each patient is his own control, only one eye being treated.

Kidneys.—Relatively few patients with diabetic kidney disease have been treated by dialysis. The patients are often unsuitable because they have other general complications, but if these are absent diabetics should be considered in the same way as other patients. It is worth remembering that diabetic kidney disease may progress very slowly and patients may live reasonably healthy lives for some years with blood urea levels of 100 mg. per 100 ml.