The 14-day Gemini and 18-day Soyuz missions dispel the dire prophecy that man's bones may fold up after three or four weeks in the weightless state. Artificial gravity may have an important part to play in the exploration of the solar system, particularly with regard to changes in cardiovascular function,⁷ but refrigeration is in the realm of science fiction rather than the logical development of aerospace physiology.—I am, etc.,

A. N. NICHOLSON.

Royal Air Force Institute
of Aviation Medicine,
Farnborough, Hants.

REFERENCES

- 1 Deitrick, J. E., Whedon, G. D., and Shorr, E., *American Journal of Medicine*, 1948, 4, 3.
- 2 Brannon, E. W., Rookwood, jun., C. A., and Potts, P., *Aerospace Medicine*, 1963, 34, 90.
- 3 Birkhead, N. C., et al., *Wright-Patterson Aeromedical Research Laboratory Report TDR-64-61*, Ohio, United States Air Force, 1964.
- 4 Vost, F. B., et al., *National Aeronautics and Space Administration Report CR-182*, Washington D.C., National Aeronautics and Space Administration, 1965.
- 5 Mack, P. B., Vose, G. P., Vogt, F. B., and LaChance, P. A., in *Gemini Mid-program Conference*, p. 407, Washington D.C., National Aeronautics and Space Administration, 1966.
- 6 Nicholson, A. N., *British Journal of Hospital Medicine*, 1970, 3, 836.
- 7 Nicholson, A. N., *The Times*, 29 June 1970, p. 9.

Problems of Tuberculosis in Immigrants

SIR,—In the report on the Annual Scientific Meeting (18 July, p. 155). Dr. D. K. Stevenson (Bradford) is quoted as follows: "Immigrants had to have a medical check in their own country, but this did not apply to dependants. The facilities on entry were inadequate to deal with these, so that the problem was passed to the local health authorities . . ."

I am aware, as I am sure every practising doctor in Britain is aware, that potential emigrants to Australia and other Commonwealth countries are required to have rather more than a "medical check" before leaving the United Kingdom, and that a normal radiological examination of the chest is obligatory for all intending immigrants to the United States.

In all the political humbug on immigration which appears daily in the lay press my medical colleagues must surely share my disgust at the absence of any reference to the "tuberculosis problem," the "V.D. problem," and even the "trachoma problem." Are we not entitled to know the average cost of diagnosis, hospitalization, and treatment for immigrants admitted to this country with active disease, and the measures which have been or perhaps have yet to be recommended for dealing with the inadequate facilities on entry?

Perhaps a significant clue to this national scandal is to be found in the opening sentence of the report of the Scientific Meeting for Thursday 2 July, which suggests this is a parochial problem since all the speakers were from Bradford. In reality, Bradford is just one of many local authorities faced with the tuberculosis problem. They must all reflect, somewhat ruefully, that the final chapter of this problem was forecast for the early 1970's.—I am, etc.,

J. R. A. HARVEY.

London W.1.

Acute Malaria in Newborn Infants

SIR,—Dr. N. E. Okeke's letter (11 July, p. 108) quoting that "more than one third" of 50 African women who were delivered some five years ago in a medical centre in Eastern Nigeria had "malaria parasites" in their blood is not unduly surprising; this is the common parasite rate in unprotected adult pregnant females in highly endemic areas of Africa.

Far more surprising is the statement that such a high parasite rate was found in women who allegedly were on a prophylactic antimalarial regimen. One would wish to know more about the actual drug given, its dosage, and its regularity. Even more puzzling is the assertion that in all cord bloods from these women, and in the peripheral blood of their newborn, malaria parasites were found.

This observation is contrary not only to that of Blacklock and Gordon¹ quoted by Dr. Okeke, but disagrees also with the views of Covell,² Bruce-Chwatt,³ and Garnham,⁴ and with several other experienced workers in Africa whose concerted opinion is that in women in endemic malarious areas the infections of the placenta may be frequent but the incidence of congenital malaria is exceedingly low (less than 0.1%). It would be of interest if Dr. Okeke's findings could be reported in detail to allow for a proper appraisal of this disclosure of an exceptionally high frequency of congenital malaria in Eastern Nigeria during the period 1965-7. The reference to 0.3-0.5 ml. of intramuscular chloroquine given to infants does not indicate the actual dosage in terms of chloroquine base, and it would be interesting to know something of the results of this treatment.—I am, etc.,

L. J. BRUCE-CHWATT.

London School of Hygiene
and Tropical Medicine,
London W.C.1.

REFERENCES

- 1 Blacklock, B., and Gordon, R. M., *Annals of Tropical Medicine and Parasitology*, 1925, 19, 37.
- 2 Covell, G., *Tropical Diseases Bulletin*, 1950, 47, 1147.
- 3 Bruce-Chwatt, L. J., *Annals of Tropical Medicine and Parasitology*, 1952, 46, 173.
- 4 Garnham, P. C. C., *Malaria Parasites*, Oxford, Blackwell, 1966.

Acute Malaria in Newborn Infants

SIR,—Dr. N. E. Okeke (11 July, p. 108) asks about the experience of other doctors working in the tropics of the incidence they have noted of acute malaria in newborn infants.

I have described¹ how we found in 700 cases examined in the Fort Hall district of Kenya two infants with subtertian malarial parasites in their blood, which was taken immediately after they were born. In the mothers we observed in specimens of placental blood one benign tertian, 30 subtertian, and 9 quartan malarial parasites.

Also it is of interest to observe that we found in one patient *Microfilaria bancrofti* and in another patient *Spirillum rossi* in the

placental blood. These figures are not as impressive as those of Mackay.²—I am, etc.,

P. G. PRESTON.

Colchester,
Essex.

REFERENCES

- 1 Preston, P. G., *East African Medical Journal*, 1942, 19, 223; 247.
- 2 Mackay, R., *East African Medical Journal*, 1933, 10, 246.

Cervico-oculo-acoustic Dysplasia

SIR,—In the reported case of the above syndrome in an adult female by Dr. D. A. A. Primrose (11 July, p. 108) he described the syndrome as rare. This apparent rarity is probably due to the several specialties involved in the management of its features (otologists, oculists, paediatricians, or psychiatrists). To establish a more accurate incidence of the syndrome each specialist would be required to search for the presence of often minimal abnormalities in another system outside his own specialty.

By survey of the school health service, audiology clinics, and area mental deficiency service we found seven cases of cervico-oculo-acoustic syndrome among school-age Glasgow children.¹ There are some 200,000 school children in Glasgow and 17,000 are surveyed by the School Health Service audiometricians annually. We did not, however, survey orthopaedic or ophthalmic clinics, and it is likely that additional cases are referred to one or other of these disciplines without cross-referral to audiology or mental deficiency.

We note that the case described by Dr. Primrose had an occipital meningocele. Two further adult females found by us similarly show occipital meningocele, and indeed Wildervanck^{2,3} in his original description includes rachischisis as a salient feature of the dysplasia.—We are, etc.,

W. I. FRASER.

Lynebank Hospital,
Dunfermline.

R. C. MACGILLIVRAY.

Lennox Castle,
Lennoxtown.

REFERENCES

- 1 Fraser, W. I., and MacGillivray, R. C., *Journal of Mental Deficiency Research*, 1968, 12, 322.
- 2 Wildervanck, L. S., *Nederlandsch Tijdschrift voor Geneeskunde*, 1960, 104, 2600.
- 3 Wildervanck, L. S., in *Proceedings of the Second International Course in Paedo-Audiology*, ed. H. C. Huizing, p. 90. Groningen, 1961.

Police Drug Searches

SIR,—The licence given to police to search citizens for illegal possession of drugs is open to much abuse.

Recently I have seen young people searched in the middle of Piccadilly Circus in a way that could hardly help to solve the drug problem, and seemed devoid of all human dignity. Also in the past few months I have had two patients, both with a psychiatric history, seized by the police and kept in a police station till I informed the police that the possession of tablets or a prescription for tablets was medically authorized.