facturer's smallest standard pack, or a multiple of this number. This will enable the pharmacist to order the exact quantity required, and will have the desired effect of removing the temptation to steal a large stock bottle from the pharmacist's shelf.—I am, etc.,

M. J. LEVERTON.

Millom,
Cumberland.

Surgery for Duodenal Ulcer

SIR,—Your leading article on results of vagotomy (15 August, p. 358) offers little advance on the conclusions of six years ago.¹ The preliminary results of my own cases of peptic ulcer may therefore be of interest.

Since 1967 I have treated, by pyloroplasty alone, 55 patients suffering from chronic peptic ulcer requiring surgery (38 duodenal, 17 gastric). Thirty-seven patients have been followed for more than one year and 15 of these for more than two years. None now suffers any dyspepsia; nor have there been any side effects of the pyloroplasty. A second pyloroplasty was necessary in five patients before the ulcer healed. The first operation did not produce a wide, incompetent pylorus, but after a second and effective pyloroplasty the ulcers healed and remain so 20 months (one duodenal), one year (one duodenal, one gastric), and less than one year (one duodenal and one gastric) later. In three cases, preoperative peak acid secretion was high but fell to within normal range post-operatively. In seven patients eating excessive fat caused symptoms which are fully controlled by a low fat intake.

Capper² showed that benign gastric ulcer healed following pyloroplasty alone "because a wide, incompetent pylorus does not permit a narrow, forceful regurgitation" of duodenal contents into the stomach. Capper's words—narrow, forceful—led me to apply his conclusions to duodenal ulcer as well. Farris and Smith³ accepted that cure of anatomical obstruction at the pylorus was the most important factor in cure of gastric ulcer but added vagotomy to pyloroplasty for fear that "duodenal ulcer diathesis might recur." Their fears appear to be unfounded. Martin and Burden⁴ in 1928 considered that failure of cure by pyloroplasty of any peptic ulcer was due to incomplete abolition of sphincter activity at the pylorus, or to post-operative narrowing of the pyloroplasty. Their views were neglected perhaps because although we can assess the time the stomach takes to empty of barium we cannot see whether it is "swishing like a bath of water"⁵ through a relaxed, wide-open pylorus, or is squirting as a narrow, forceful "jet" through a restricted pylorus (restricted by spasm, muscle hypertrophy, inflammation and fibrosis, or even pyloric mucosal diaphragm). Perhaps also, because it was possible to "measure" them, however inaccurately, acid levels attained spurious importance.

The concept underlying use of pyloroplasty alone is a simple one. If the pylorus does not relax fully the flow becomes narrow and forceful; acid gastric contents are squirted as a "jet" to impinge upon duodenal mucosa, possibly persistently on the same localized area, before mixing with and being neutralized by duodenal

contents. If the pylorus does not close completely during duodenal cap systole pyloric reflux occurs. Resistance to flow of gastric contents through the pylorus into the duodenum may stimulate the antrum and cause a rise in gastric acid secretion.^{6,7} Anything which diminishes the "jet" itself (for example, by interfering with gastric motility), or which alters pH, can modify the ulcerating effect. Pyloroplasty prevents the formation of a "jet."

Pyloroplasty must be adequate to prevent reflux from duodenal cap systole and to decompress the antrum. It is then followed by healing of duodenal and high and low gastric ulcers, high acid secretion levels may become normal, and there are no side effects of the operation. I have not found any kind of vagotomy or gastrectomy necessary to cure any chronic peptic ulcer during the last two-and-a-half years. I have also used effective pyloroplasty alone successfully to cure cases of failed vagotomy and pyloroplasty and failed vagotomy and gastroenterostomy.—I am, etc.,

PETER CHILDS.

Plymouth General Hospital,
Plymouth, Devon.

REFERENCES

- 1 *British Medical Journal*, 1964, 1, 448.
- 2 Capper, W. M., *Annals of the Royal College of Surgeons of England*, 1967, 40, 21.
- 3 Farris, J. M., and Smith, G. K., *Surgical Clinics of North America*, 1966, 46, 329.
- 4 Martin, E., and Burden, V. G., *Annals of Surgery*, 1928, 88, 565.
- 5 Johnson, H. D., *Proceedings of the Royal Society of Medicine*, 1961, 54, 938.
- 6 Harkins, H. N., et al., *Surgical Forum*, 1954, 5, 281.
- 7 Menzies, R., *Surgical Clinics of North America*, 1966, 46, 257.

Pseudo-obstruction of the Large Bowel

SIR,—I am surprised that Dr. Barbara F. Smith (20 June, p. 732) and Mr. J. A. C. Neely (27 June, p. 793) have objected to the use by Mr. P. K. Caves and Dr. H. A. Crockard (6 June, p. 583) of the term "pseudo-obstruction." Dr. Smith's objection is that "pseudo-obstruction" was used to describe another condition of apparent obstruction by Naish and his colleagues in 1960. I would draw her attention to the use of the term by Dudley *et al.* in 1958¹ to describe the condition reviewed by Mr. Caves and Dr. Crockard.

Mr. Neely states that the term ileus should be used rather than obstruction, but the word ileus comes from the Greek *eileos* which means "intestinal obstruction." He would qualify the ileus as "paralytic," yet refers to a condition with features not of paralysis but of disorganized or ineffective function (the presence of bowel sounds associated with colic). He believes that the obstruction is not "pseudo" but is real, yet agrees that the tube is not blocked, but has an inefficient propulsive mechanism, and so appears with most features of blockage or organic obstruction.

To me, the term "pseudo-obstruction" describes the circumstances distinctly and effectively. If adopted universally it would make for clarity in terminology, and even improvement in its understanding.—I am, etc.,

F. O. STEPHENS.

Sydney Hospital,
N.S.W., Australia.

REFERENCE

- 1 Dudley, H. A. F., et al., *Journal of the Royal College of Surgeons of Edinburgh*, 1958, 3, 206.

Polyuria after Cardiac Surgery

SIR,—We were interested to read the report by Dr. R. O. Robinson and Mr. K. M. Pagliero (1 August, p. 265) of a patient with polyuria after cardiac surgery. Two patients in the Liverpool Regional Cardio-thoracic Centre undergoing open heart surgery developed polyuria postoperatively, and in one there was a dramatic response to pitressin.

A 34-year-old housewife with a 27-year history of a cardiac murmur presented with an 18-month history of dyspnoea on effort and palpitations. Physical examination revealed an atrial septal defect and the E.C.G. showed a right bundle branch block. At cardiac catheterization, a large atrial septal defect was found with a 1.5:1 shunt.

At operation a 15 sq. cm. atrial septal defect was repaired using a Dacron patch on full normothermic cardiopulmonary bypass with a perfusion time of 45 minutes.

She returned to the intensive care unit fully conscious and co-operative with normal pulse, blood pressure, and central venous pressure. The urine output was adequate, glycosuria was present, but blood sugar was normal. The next day she was vague and had echolalia, but there were no other abnormal neurological features. A brisk diuresis started with a urine flow of 9 ml./min. for eight hours and then returned to normal. Twelve hours later it again increased to 8 ml./min., and after 1½ hours she had a major generalized convulsion. This persisted until she was curarized, intubated, and put on a ventilator. The urine flow then returned to normal. A cardiac arrest developed from unknown causes and following successful resuscitation she was found to have bilateral pyramidal damage and was comatose. Her cardiac, renal, and cerebral function did not alter during the next 10 days, when she succumbed to an overwhelming bronchopneumonia. Necropsy revealed petechial haemorrhages throughout the cerebral white matter but no softening or oedema.

A 39-year-old housewife, who had had a mitral valvotomy 14 years previously, was admitted to hospital for investigation of breathlessness, haemoptysis, and ankle swelling. Physical examination revealed a pure mitral stenosis and valvotomy was recommended. A left-sided empyema had complicated the original operation and a sinus had drained intermittently since.

An attempt to perform a mitral valvotomy from the right side failed, so the patient was put on to cardiopulmonary bypass and the mitral valve split to 3.5 cm. with a Tubbs's dilator. There was no valve clarification, no left atrial thrombus, and the aorta was cross-clamped during the period of left atriotomy. She did not regain consciousness after the operation and was ventilated. Neurological examination revealed bilateral pyramidal damage. She required an isoprenaline infusion for bradycardia and hypotension. In the 12-hour period postoperatively she passed 1.6 l. of urine in excess of input. The blood urea, calcium, potassium, and magnesium were all normal. The blood sugar, initially 325 mg./100 ml., fell to 124 mg. within 6 hours. The following day she became hypothermic and the diuresis persisted, urine S.G. 1000. Ten units of the short-acting Pitressin preparation were given subcutaneously and the urine flow fell from 7.0 ml./min. to 0.75 ml./min. The urine S.G. rose to 1010.

Fifteen hours later, with a normal urine output, the ventilator failed. When she was put back on another machine the tidal volume was smaller, and she became hypotensive with a tachycardia, and the urine flow ceased. She was found to be acidotic, hypoxic, and hypercapnoeic. Restoration of normal ventilatory function did not improve either cardiac or renal function and she died 120 hours after surgery. Necropsy was not performed.

In the second case diffuse cerebral damage was recognized before the onset of polyuria, and hypothermia was also present, suggesting a disturbance of hypothalamic

function to account for reduced antidiuretic hormone production. The cerebral dysfunction in this patient was thought to be due to hypotension at the time of operation. It is interesting that no hypothalamic disturbance was recognized by Javid *et al.*¹ in their review of neurological complications after open heart surgery.

There was no obvious cause for the cerebral dysfunction in the first case, but it is interesting that polyuria and echolalia began simultaneously and that fluctuations of urine flow coincided with alterations in cerebral function. To have been able to estimate levels of antidiuretic hormone at these times would have been helpful.

Polyuria after open heart surgery may be associated with diabetes mellitus, biochemical changes, such as hypokalaemia and as an exaggerated response to diuretics, and it can lead to peripheral circulatory failure. It is important to recognize polyuria due to diabetes insipidus as it is so amenable to Pitressin (vasopressin) therapy.—We are, etc.,

C. C. EVANS.

J. B. MEADE.

W. A. LITTLER.

Liverpool Cardio-thoracic
Surgical Centre,
Liverpool.

REFERENCE

- ¹ Javid, H., *et al.*, *Journal of Thoracic and Cardiovascular Surgery*, 1969, 58, 502.

Wrong Operations

SIR,—I was very pleased to read Mr. I. W. Payne's letter (5 September, p. 584), and fully agree with all that he says. It is of course of the utmost importance that every care should be taken to avoid operating on the wrong side, or even worse performing the wrong operation on the wrong patient. Like Mr. Payne I have never felt quite happy with regard to the marking of patients prior to surgery. On the one hand it is still possible, as he so rightly points out, to mark in error, and on the other hand when the patients are thus branded do they perhaps wonder if the surgeon has a full grasp of their illness, when apparently he has to depend on a mark to tell him what to do and where to make his cut?

My own custom is to see every patient before being anaesthetized in the anaesthetic room: to check on the notes, to check with the patient, and above all to reassure the patient—this last being of paramount importance. It may be argued that some patients owing to premedication may not be in a position to give a clear answer, but in my experience this is seldom the case. For many years I have established a routine whereby none of my patients are anaesthetized before I have had an opportunity of speaking to them, and apart from serving as a most valuable additional check I believe it to be much appreciated. I am against the practice of starting the next anaesthetic before the last case has been finally completed. This may save a little time with a long list, but surely adds to the chance of error, and as Mr. Payne says the surgeon is in constant need of vigilance and cannot afford to cut corners.

I should make the point I am not against the practice of marking patients, but making

a plea that this in itself is not sufficient. The personal approach cannot be replaced by the indelible mark.—I am, etc.,

ANDREW G. BUTTERS.

Barnsley,
Yorks.

Unusual Surgical Emergency in Pregnancy

SIR,—Inguinal hernia in pregnancy containing one horn of a bicornuate uterus and associated with a congenital solitary kidney is rare, so I would like to report the following case.

A 35-year-old woman in her fifth pregnancy was admitted as an emergency at 37 weeks gestation because a left indirect inguinal hernia had become irreducible. On admission apart from pain her general condition was satisfactory, and the pregnancy was normal. A large left inguinal hernia was present, and it was tender and irreducible. At operation the hernial sac was found to contain one horn of the uterus, about 8 × 6 cm. size, the left round ligament, and the left uterine tube with the ovary attached. The horn of the uterus and the adnexa were engorged but viable; they were reduced and a Bassini type of herniorrhaphy was performed. Her post-operative progress was satisfactory and she was delivered three weeks later of a live female baby weighing 3,430 g.

A hysterosalpingogram carried out six weeks post partum showed a uterus bicornis unicollis with dilatation of the right horn and free spill of dye into the peritoneal cavity. The left tube was not outlined (Fig.). An intravenous

pyelogram showed an enlarged left kidney with pelvicalyceal dilatation. There was no evidence of a kidney on the right side. The creatinine clearance was then 60 ml./min., and her blood urea 23 mg./100 ml.

Almost every abdominal organ has been described in the sac of an inguinal hernia.¹⁻³

A serious complication of a bicornuate uterus in labour is that the non-pregnant horn of the uterus can cause obstruction during delivery, but the case described above confirmed the clinical impression that the more complete the uterine duplication the less likely is dystocia.⁴ This case illustrates the fact that congenital malformations of the genital and urinary tracts are associated.^{5,6}—I am, etc.,

A. MAHMOOD.

Southlands Hospital,
Shoreham, Sussex.

REFERENCES

- ¹ Fusaro, W. J., *New York State Journal of Medicine*, 1946, 46, 1024.
- ² Tunney, R. B., and Hunter, R. E., *New England Journal of Medicine*, 1959, 242, 1011.
- ³ Sugarman, G. R., *Journal of the Newark Beth Israel Hospital*, 1961, 12, 160.
- ⁴ Chassar Moir, J., in Munro Kerr's *Operative Obstetrics*, 7th edn., London, Baillier, Tindall and Cox, 1964.
- ⁵ Woolf, R. B., and Allen, W. M., *Obstetrics and Gynecology*, 1953, 2, 236.
- ⁶ Chawla, S., Bery, K., and Indra, K. J., *British Medical Journal*, 1966, 1, 1398.

Value of Osteopathy

SIR,—In a review (6 December, 1969, p. 610) of the book *A Manual of Osteopathic Practice* by Alan Stoddard, Dr. D. A. H. Yates expressed his doubts on the validity of assessing leg length by palpating the iliac crests in the erect position. In a letter (7 February, p. 366) Dr. Yates reported some studies of observer error in measuring leg length by this method. He concluded that "with raises of $\frac{1}{2}$ in. (1.2 cm.) or less the observers were as often incorrect as correct as to which side, if either, was raised. With $\frac{3}{4}$ and 1 inch (1.8 and 2.5 cm.) all observations were correct."

The following experiment was carried out to test the validity of such an observation, and to find out with what degree of accuracy a known amount of shortening could be detected.

One subject was used. He stood with his back to a good light, his feet were obscured by a blanket, and five observations were made with varying raises under one or other foot: no raise, right foot raised $\frac{1}{2}$ in. (1.2 cm.), left foot raised $\frac{1}{2}$ in. (0.6 cm.), left foot raised $\frac{1}{2}$ in. (1.2 cm.), and finally left foot raised $\frac{1}{2}$ in. (0.3 cm.).

The observer stood behind and assessed the level of the pelvis by palpating the highest point of the iliac crest. The short leg was recorded or the pelvis stated to be level. Seventeen observers each made five observations. The observers were a group of doctors, mostly general practitioners, all with at least two years' experience of manipulative medicine, attending a postgraduate course run by the British Association of Manipulative Medicine.

Raise in Inches	Opinions of Observers		
	Level Pelvis	Short R. Leg	Short L. Leg
No raise	2	4	11
$\frac{1}{2}$ under	8	6	3
$\frac{1}{2}$ left	9	7	1
$\frac{1}{2}$ foot	2	14	0
$\frac{1}{2}$ under right foot	2	0	15

The results are seen in the Table. With the subject standing naturally, 64% of the observers found a short left leg. A raise of