Use of Google as a diagnostic aid

Bias your search

EDITOR—Although Google can be a valuable aid in reaching diagnoses in rare conditions,¹ the problem faced in a busy, fast-paced clinic can be the abundance of lesser quality material on the web. “Biasing” or directing your search by adding the name of an online clinical knowledge base such as Emedicine can lead you to quality information more quickly; you avoid sifting through dozens of unknown journals, personal webpages, blogs, or discussion forums containing your search terms.

For example, by typing “pigmented lesion+buccal” into Google you will get 55 000 hits with a vast array of very different material. However, typing “pigmented lesion+buccal+emedicine” provides only 106 hits (accessed 12 November 2006), the first being a comprehensive Emedicine review on Peutz-Jeghers syndrome.² You get similar results by adding “BMJ” to the end of your search; in this case the top hit is an ABC of oral health article.³

Mark Taubert *specialist registrar in palliative medicine*

Holme Tower Marie Curie Centre, Penarth, South Glamorgan CF64 3YR
mtaubert@hotmail.com

Competing interests: None declared.

1 Tang H, Ng JHK. Googling for a diagnosis—use of Google as a diagnostic aid: internet based study. *BMJ* 2006;333:1143-5. (2 December.)

2 www.emedicine.com/med/topic1807.htm

3 Scully C, Porter S. The ABC of oral health—Swellings and red, white and pigmented lesions. *BMJ* 2000;321:225-8.

Is Google like 10 000 monkeys?

EDITOR—The suggestion by Tang and Ng, that everything can be found on the web if only one knew the correct search terms is of course not surprising but invites some worrying conclusions¹:

- Lawyers can with ease and within minutes find cases of patients displaying certain symptoms which also include the correct diagnosis, which, they will suggest, a “negligent” clinician missed.
- The excuse, used since time immemorial, “I couldn’t possibly know,” and providing comfort to the ignorant, has lost some of its credibility.

One can find everything on Google, given time. Is Google perhaps the modern version of 10 000 monkeys sitting at 10 000 typewriters, who, given time, will eventually produce a true copy of *Hamlet*? Among the one correct version there will of course be

thousands with just one or two typos. So, can Mark Twain’s remark “Be careful about reading health books. You may die of a misprint”² be applied to Google?

Reinhard Wentz *retired*
Twickenham TW2 7PS
sleuthmedical@yahoo.com

Competing interests: RW uses Google often, when appropriate.

1 Tang H, Ng JHK. Googling for a diagnosis—use of Google as a diagnostic aid: internet based study. *BMJ* 2006;333:1143-5. (2 December.)

2 http://en.wikiquote.org/wiki/Mark_Twain.

Authors’ reply to responses

EDITOR—A common misinterpretation of our paper in the lay press was that doctors and patients could magically become diagnosticians simply by using web search engines.¹ The art of diagnostic reasoning is difficult, and doctors become competent at it only after years of training and seeing patients.²

Our intended message was that in cases of “mystery illnesses,” a web search may be fruitful in finding articles that may suggest a diagnosis that would not otherwise be considered because of the rarity of the illness and unfamiliarity with its symptoms and signs. Any article located from the web should be critically appraised in the usual manner. We are less concerned about the philosophical objections of how an article is located than with its usefulness in patient care.

To state the obvious, there is no danger of “Google misdiagnosing life threatening disease” as search engines cannot make diagnoses. Only doctors are capable of making diagnoses (and misdiagnoses). If the probability of a diagnosis exceeds the testing threshold,³ then tests would be performed to prove or disprove the diagnosis. However, the diagnosis has to be considered in the first place and search engines may act as a diagnostic reminder.

We do not think that our article downplays the role of medical librarians in the practice of evidence based medicine. We have benefited enormously from workshops and help from

librarians. Most questions in clinics can be answered using Pubmed, the *Cochrane Library*, journals, Uptodate, etc. However, if traditional sources of information have failed to answer these questions, then it would be unwise to ignore the web as a resource.

The sight of clinic patients clutching internet printouts fills some doctors with dread and may affect the doctor-patient relationship.⁴ Information found on the web is of variable quality but risk of harm to patients is probably small.⁵ Doctors in training should become proficient at using the internet because they can guide their patients to reliable websites. When used properly, the internet empowers both patients and doctors and may improve the quality of care.⁴

Hangwi Tang *respiratory and sleep physician*
hangwitang@yahoo.com

Jennifer H K Ng *consultant rheumatologist*
Princess Alexandra Hospital, Brisbane, QLD 4102, Australia

Competing interests: None declared.

1 Tang H, Ng JHK. Googling for a diagnosis—use of Google as a diagnostic aid: internet based study. *BMJ* 2006;333:1143-5. (2 December.)

2 Bowen JL. Educational strategies to promote clinical diagnostic reasoning. *N Engl J Med* 2006;355:2217-25.

3 Pauker SG, Kassirer JP. The threshold approach to clinical decision making. *N Engl J Med* 1980;302:1109-17.

4 Murray E, Lo B, Pollack L, Donelan K, Catania J, White M, et al. The impact of health information on the internet on the physician-patient relationship: patient perceptions. *Arch Intern Med* 2003;163:1727-34.

5 Crocco AG, Villasis-Keever M, Jadad AR. Analysis of cases of harm associated with use of health information on the internet. *JAMA* 2002;287:2869-71.

Summary of other responses

bmj.com

A few of the 40-odd respondents to the article by Hang and Ng think that the idea of the Google search engine being used as a diagnostic aid is downright laughable or dangerous, but most like this “new” facility, some even rating it quite highly.¹ Several doctors identify the need for training in using search engines; this seems to have been implemented in several teaching facilities but perhaps should become the gold standard everywhere. Several information specialists point out that doctors might be well advised to use librarians’ training and skills to conduct com-

plex searches, rather than try to do it themselves. That Google is forming partnerships with libraries is seen as a step in the right direction.

Others remind us that Google is not set up as a diagnostic decision support system—although it can be a useful aid to differential diagnosis once a diagnosis has been made—and especially that Google is not a content provider. And many respondents mention common conditions in their case for or against, rather than the rare conditions that Hang and Ng used in their searches. One or two correspondents praise the usefulness of Google and other search tools in particular specialties, for patients, and in veterinary medicine.

Many remind us that many other internet resources exist, from PubMed to specialty databases, which might be more specific and useful than Google, although Google is seen as a good first step. That Google can help doctors in remote areas or developing countries also finds mention in various responses, and one correspondent praises its usefulness in elucidating the geographical history of an illness.

The jury is still out on whether searching for symptoms on the internet is the way forward for doctors and consumers. But a sense of excitement at this new resource is palpable, even if it may be used for aspects of medical practice other than diagnosis.

Birte Twisselmann *assistant editor (web)*
BMJ

Competing interests: None declared.

1 Rapid responses. Googling for a diagnosis—use of Google as a diagnostic aid: internet based study. www.bmj.com/cgi/eletters/333/7579/1143.

Reconfiguring acute hospitals in England

Clinicians are fatigued by health policy and policy makers

EDITOR—Transferring “less sick” people into non-acute hospital settings doesn’t necessarily save money.¹ At least 15 schemes, vigorously supported and in some cases piloted by our acute hospital, have been rejected or discontinued by the primary care trust on financial grounds. The University of Birmingham has reviewed the evidence for the cost effectiveness of moving less sick patients into non-hospital settings and concludes that these changes will not automatically reduce costs.² Sadly this document also concludes that there is little evidence that this health policy will reduce reliance on acute hospital care, despite the premise that there is a reduced need for acute hospital beds.

We agree with Ham that the NHS should make it clear why change is needed and articulate a persuasive and reasoned case to support proposals that are bound to be controversial.¹ What we don’t need are further eloquent health policy editorials and overviews to “clarify.” As clinicians, we have been driving these changes in a timely and safe fashion as a consequence of clinician driven advances in medical care and the recognition of an ageing population whose

healthcare expectations are rising. Clinicians are not the dinosaurs resistant to change that the Department of Health would like the public to believe. What we require is the detail to show that the rushed and financially driven current health policies are safe for our patients. The BMA’s stance on hospital reconfiguration is that changes to our hospital services must be based on delivering the highest possible standard of care for patients. Reconfiguration to suit political needs or decisions based solely on cost would be completely unacceptable.³ Policies that are driven fundamentally by financial pressures are of course bound to be controversial, and Ham as professor of health policy and management, would do well to highlight this to those who can influence policy.

Richard Venn *consultant in intensive care*
richard.venn@wash.nhs.uk

Lui G Forni *consultant in renal medicine and intensive care*
Worthing Hospital, Worthing,
West Sussex BN11 2DH

Competing interests: None declared.

1 Ham C. Reconfiguring acute hospitals in England. *BMJ* 2006;333:1135-6. (2 December.)

2 University of Birmingham Health Services Management Centre. Making the shift: a review of the written evidence. www.hsmc.bham.ac.uk/news/MakingtheShift6881.pdf

3 Foster M. The great debate. *BMA News* 2006 Dec 2:7.

Try reconfiguring services to improve standards and meet patient preferences

EDITOR—Reconfiguration of services in Hexham has come with the replacement of the Nissen hut hospital with a private finance initiative development.¹ This, taken with the linking of services in Hexham with the hospital in Wansbeck, has resulted in some strange effects even if the objective standard of care has risen.

Patients fracturing their neck of femur in the Hexham catchment area will be taken initially to Hexham for diagnosis and stabilisation—this might entail an ambulance journey of up to 50 miles (80 km). Then they will be transferred to Wansbeck, another 50 mile journey, for operation—they are now 100 miles from home. On this second journey they drive past an even larger centre of excellence, Newcastle, which is a mere 30 miles from Hexham.

An audit that I conducted recently on patients in my practice who had had a baby within the last year revealed that 100% of them wanted Hexham to have a fully featured 24/7 obstetric provision instead of the present excellent midwife led unit. Most women who delivered in Newcastle commented adversely on the frantically busy environment there.

The point I seek to make with these two illustrations is that local services and patient preference can be delivered, and geographically appropriate provision made with a little more thought than the market and politics usually permit. While Hexham is destined to be a regional (?) centre of excellence for laparoscopic surgery there seems to me to be no compelling reason why it should not be a centre of excellence for

other high demand services too—sharing Newcastle’s, Carlisle’s, and Wansbeck’s burden more sensibly and equitably.

Looking to the future, transport is going to become cripplingly expensive within the lifetimes of the new units. Infection control is easier in smaller physically remote units because if one is closed the other(s) can continue. Local people want local services. Should we castigate them for being so hopelessly backward and having the temerity to have a preference and state it? Perhaps the putative variation in standards is a price they are willing to pay.

Steven Ford *general practitioner*
Haydon and Allen Valleys Medical Practice,
Hexham, Northumberland NE47 6LA
doctor.ford@virgin.net

Competing interests: None declared.

1 Ham C. Reconfiguring acute hospitals in England. *BMJ* 2006;333:1135-6. (2 December.)

A side effect of patient choice?

EDITOR—Ham illustrates the tension between the need for collaboration between hospitals to ensure appropriate location of specialist services and the opposing need to compete promoted by a pro-market policy agenda which undermines collaboration.¹

However, the editorial does not explore the effect that the monomaniac promotion of patient choice to the exclusion of any other considerations may have had in fuelling public opposition to reconfiguration.

Patients and the general public have repeatedly been assured that all aspects of reform in the NHS have been driven solely by the desire to give them more choice and by implication better services.

It is no surprise then that they find it difficult to understand why they cannot have an accident and emergency department on their doorstep if they so choose. A recent national survey suggested that one in three people believed the NHS should provide “all drugs and treatments, no matter what the cost,” while a further four out of 10 believed that the NHS should provide the “most effective treatment, no matter what the cost.” Against this backdrop, it is not surprising that the natural public resistance to change seems more resilient than ever on the subject of reconfiguration of services.

The rising cost of health care, the advent of newer and more expensive drugs, and the ageing population of the United Kingdom all indicate a need for more robust public debate on resource allocation and use within the NHS. The failure of policy makers to lead and promote this debate is in part responsible for the current difficulties in selling reconfiguration plans to the public.

Ike Anya *specialist registrar in public health medicine*
Department of Epidemiology and Public Health,
University College London WC1E 6BT
ikeanya@doctors.org.uk

Competing interests: None declared.

1 Ham C. Reconfiguring acute hospitals in England. *BMJ* 2006;333:1135-6. (2 December.)

Author's reply to review of *Bioethics and Armed Conflict*

EDITOR—As Nathanson says in her review of my book,¹ I argue that the use of “moderate physical pressure” might be defensible in some cases. “Defensible” is the right word. The acts may be excused but are not justified. However, I don’t defend interrogational torture with any great enthusiasm but simply admit that there might be some rare case where perpetrators might mount a successful defence. This is not because anyone rejects human rights but because even supremely important and inalienable rights can conflict. One person’s right to dignity collides with another person’s right to life, and in spite of every precaution and after exhausting every alternative you sometimes have to make a choice.

The weapons issue is more complicated. I make a case for the use of non-lethal chemical weapons. There are many instances in contemporary warfare where non-lethal weapons may cause considerably less harm than high explosives—and, yes, physicians should help build them. At the same time, I suggest that some nations, particularly those in the Middle East that have not signed the international treaties banning chemical and biological weapons, have grounds to develop these weapons for deterrent purposes. Doctors should help build these too if their expertise is necessary. But as anyone familiar with nuclear weapons knows, there is a huge and important difference between supporting the use of these weapons and supporting their development for deterrent purposes.

Although torture and, to a lesser extent, weapons development have attracted a great deal of attention, it is far more important to emphasise the role that medicine can play in conflict resolution, the protection of human rights, and the prevention of unjust war. But that’s the next chapter, the one that Nathanson does not review. It would be a shame if people get stuck on torture and weapons and don’t make it to peace.

Michael L Gross *professor*
Department of International Relations, University of Haifa, Haifa, Israel
mgross@poli.haifa.ac.il

Competing interests: MLG wrote the book that was reviewed.

¹ Nathanson V. Bioethics and armed conflict: moral dilemmas of medicine and war. *BMJ* 2006;333:1177-a. (2 December.)

Effective probiotic treatment is rarely cheap

EDITOR—In their review on the safety of probiotics Hammerman et al describe probiotic therapy as both cheap and effective¹; we agree that probiotic therapy seems efficacious; however probiotics with proven efficacy in clinical trials cannot be described as cheap. VSL#3 is probably one of the most studied probiotics; there is good evidence that it will maintain a remission in pouchitis. However, at the dose used in the three trials showing efficacy in

pouchitis,^{2,3} VSL#3 costs \$150 (£76; €114) per month from the company’s website.

Patients wishing to use this treatment will have to bear the cost themselves as currently UK practitioners cannot prescribe probiotic therapy on NHS prescriptions, furthermore it is our understanding that US health insurance companies will not pay for this treatment⁴; this may be a significant barrier to both use and compliance in clinical practice.

Probiotics from health food shops often do not contain what is stated on the label and may contain pathogenic bacteria,⁵ emphasising the fact that such treatments should be obtained from a reliable source.

Simon D McLaughlin *research fellow*
simon.mclaughlin@nhs.net

Sue K Clark *consultant colorectal surgeon*
Department of Gastroenterology, St Mark’s Hospital, Northwick Park, Harrow, Middlesex HA1 3UJ

R John Nicholls *visiting professor*

Paris P Tekkis *clinical senior lecturer in surgery*
Department of Surgical Oncology and Technology, 10th Floor QEOM Building, St Mary’s Hospital, London W1 2NY

Paul J Ciclitira *professor*

Department of Gastroenterology, St Thomas’ Hospital, London SE1 7EH

Competing interests: None declared.

- 1 Hammerman C, Bin-Nun A, Kaplan M. Safety of probiotics: comparison of two popular strains. *BMJ* 2006;333:1006-8. (11 November.)
- 2 Mimura T, Rizzello F, Helwig U, et al. Once daily high dose probiotic therapy (VSL#3) for maintaining remission in recurrent or refractory pouchitis. *Gut* 2004;53:108-14.
- 3 Gionchetti P, Rizzello F, Poggioli G, et al. Probiotic therapy to prevent pouchitis onset. *Dis Colon Rectum* 2005;48:1493-4.
- 4 Shen B, Brzezinski A, Fazio VW, et al. Maintenance therapy with a probiotic in antibiotic-dependent pouchitis: experience in clinical practice. *Aliment Pharmacol Ther* 2005;22:721-8.
- 5 Hamilton-Miller JM, Shah S, Winkler JT. Public health issues arising from microbiological and labelling quality of foods and supplements containing probiotic microorganisms. *Public Health Nutr* 1999;2:223-9.

Sexual relationships between doctors and former patients

Already guilty?

EDITOR—After reading Sheather’s editorial about sex with former patients, I was seriously troubled.¹ In 1971 I performed a minor surgical procedure on a young nurse. The following year we began a relationship that was not platonic. Now I find it to have been inappropriate. Should I own up to the General Medical Council? Shall I be struck off? What can I tell the children when it hits the tabloid press? Must I now divorce the victim of this heinous abuse of my professional position?

Before rushing to confess, I read the actual advice from the GMC. Much to my surprise, given the recent performance of that august body, it was extremely reasonable. Of course, the concept of discussion with a colleague or the council before embarking on a sexual relationship is a waste of time: anyone who is prepared so to do has already thought through the possibilities of exploitation. Those who are exploiters will not feel a discussion with anyone to be appropriate. Otherwise the

guidance is good; think carefully before doing something that will affect the lives of all parties involved, which ought to apply in every aspect of doctor-patient relations.

The editorial presents a more alarmist view, apparently backed up by a very brief quotation from Graeme Catto, implying that any relationship following that of doctor and patient will be viewed as inappropriate, unless the circumstances are exceptional. Although Sheather calls for balance, there is no reason why this balance cannot be reached without breach of the published guidance. Happily, the lay press do not seem to have latched on to this subject, but it is easy to see headlines of the “Doctors wait to abuse ex-patients” variety appearing as a result of it. Perhaps I have misread the editorial, but it was more alarming than the published guidance deserves, which is not ideal in such an authoritative organ of the profession.

Gerald Freshwater *occupational physician*
Shetland Medical Services, Lerwick ZE1 0EL
freshwater@zetnet.co.uk

Competing interests: GF is subject to registration with the GMC.

¹ Sheather J. Sexual relationships between doctors and former patients. *BMJ* 2006;333:1132. (2 December.)

Dating patients

EDITOR—I am often struck by the difference in attitude taken to doctors who form relationships with patients and nurses who do the same.¹ If the classic instance of a doctor-patient relationship is a male doctor and a female patient, the classic instance of a nurse-patient relationship is that of a female nurse with a male patient. The former attracts censure and possible disqualification. The latter attracts warm hearted details and best wishes published in the local press, particularly when there are wedding bells.

I have often felt that the power differential lies not so much between a male doctor and a competent and mature female patient. The power differential is between doctors and other men, who are conscious of the advantage a white coat brings in the mating ritual. I wonder how many cases are brought before the General Medical Council by disgruntled husbands.

Chris Jones *senior lecturer*
Faculty of Health, Aintree Complex, Edge Hill University, Liverpool L9 7NL
jonesch@edgehill.ac.uk

Competing interests: None declared.

¹ Sheather J. Sexual relationships between doctors and former patients. *BMJ* 2006;333:1132. (2 December.)

bmj.com

We select the letters for these pages from the rapid responses posted on bmj.com, favouring those received within five days of publication of the article to which they refer.

Letters are thus an early selection of rapid responses on a particular topic. Readers should consult the website for the full list of responses and any authors’ replies, which usually arrive after our selection.