Published in final edited form as:

Neurosci Lett. 2006 November 20; 408(3): 209–213.

Mutations in *DJ-1* are rare in familial Parkinson disease

Nathan Pankratz, Ph.D.¹, Michael W. Pauciulo, M.B.A.², Veronika E. Elsaesser, B.S.², Diane K. Marek, B.S.², Cheryl A. Halter, M.S.¹, Joanne Wojcieszek, M.D.¹, Alice Rudolph, Ph.D.³, Clifford W. Shults, M.D.⁴, Tatiana Foroud, Ph.D.¹, William C. Nichols Ph.D.^{2,5}, and the Parkinson Study Group – PROGENI Investigators

¹ Indiana University Medical Center, Indianapolis, IN;

Correspondence to: William C. Nichols, Ph.D., Associate Professor of Pediatrics, Division of Human Genetics, Cincinnati Children's Hospital Medical Center, 3333 Burnet Avenue, Cincinnati, OH 45229, Phone: 513 636-2438, Fax: 513 636-3486, bill.nichols@cchmc.org.

Parkinson Study Group Investigators:

PROGENI Steering Committee:

UNIVERSITY OF CALIFORNIA-SAN DIEGO: C Shults; CLINICAL TRIALS COORDINATION CENTER, UNIVERSITY OF ROCHESTER: F Marshall; D Oakes; A Rudolph; A Shinaman; COLUMBIA UNIVERSITY MEDICAL CENTER: K Marder; INDIANA UNIVERSITY SCHOOL OF MEDICINE: PM Conneally, T Foroud, C Halter; KANSAS MEDICAL CENTER: K Lyons; ELI LILLY & COMPANY: E Siemers.

PSG Investigators and Coordinators:

ALBANY MEDICAL COLLEGE: S Factor, D Higgins, S Evans; BARROW NEUROLOGICAL INSTITUTE: H Shill, M Stacy, J Danielson, L Marlor, K Williamson; BAYLOR COLLEGE OF MEDICINE: J Jankovic, C Hunter; BETH ISRAEL DEACONESS MEDICAL CENTER: D Simon, P Ryan, L Scollins; BETH ISRAEL MEDICAL CENTER: R Saunders-Pullman, K Boyar, C Costant-Toth, E Ohmann; BRIGHAM & WOMEN'S HOSPITAL: L Sudarsky, C Joubert; BROWN UNIVERSITY (MEMORIAL HOSPITAL OF RI): J Friedman, K Chou, H Fernandez, M Lannon; CLEVELAND CLINIC FLORIDA-WESTON: N Galvez-Jimenez, A Podichetty; CLINICAL NEUROSCIENCE CENTER: P Lewitt, M DeAngelis; COLORADO NEUROLOGICAL INSTITUTE: C O'Brien, L Seeberger, C Dingmann, D Judd; COLUMBIA UNIVERSITY MEDICAL CENTER: K Marder, J Fraser, J Harris; CREIGHTON UNIVERSITY: J Bertoni, C Peterson; HOTEL-DIEU HOSPITAL-CHUM: S Chouinard, M Panisset, J Hall, H Poiffaut; HUNTER HOMES MCGUIRE VETERANS MEDICAL CENTER: V Calabrese, P Roberge; INDIANA UNIVERSITY SCHOOL OF MEDICINE: J Wojcieszek, J Belden, C Halter; INSTITUTE FOR NEURODEGENERATIVE DISORDERS: D Jennings, K Marek, S Mendick; JOHNS HOPKINS UNIVERSITY: S Reich, B Dunlop; LONDON HEALTH SCIENCES CENTRE: M Jog, C Horn; LSU MEDICAL CENTER: J Rao, M Cook; MAYO CLINIC JACKSONVILLE: R Uitti, M Turk; MCFARLAND NEUROSCIENCES: T Ajax, J Mannetter; MCGILL CENTRE FOR STUDIES IN AGING: M Panisset, J Hall; MEDICAL COLLEGE OF GEORGIA: K Sethi, J Carpenter, K Ligon, S Narayan, L Woodward; MEDICAL COLLEGE OF WISCONSIN: K Blindauer, J Petit; MEDICAL UNIVERSITY OF OHIO: L Elmer, E Aiken, K Davis, C Schell, S Wilson; MOUNT SINAI SCHOOL OF MEDICINE NEW YORK: M Velickovic, W Koller, S Phipps; NORTH SHORE-LIJ HEALTH SYSTEM: A Feigin, M Gordon, J Hamann, E Licari, M Marotta-Kollarus, B Shannon, R Winnick; NORTHWESTERN UNIVERSITY: T Simuni, A Kaczmarek, K Williams, M Wolff; OHIO STATE UNIVERSITY: M Fernandez, J Hubble, S Kostyk, A Campbell, C Reider; OREGON HEALTH & SCIENCE UNIVERSITY: R Camicioli, J Carter, P Andrews, S Morehouse, C Stone; OTTAWA HOSPITAL CIVIC SITE: T Mendis, D Grimes, P Gray, K Haas; PACIFIC NEUROSCIENCE MEDICAL GROUP: J Sutton, B Hutchinson, J Young; SASKATOON DIST HEALTH BOARD ROYAL UNIV HOSP: A Rajput, A Rajput, L Klassen, T Shirley; SCOTT & WHITE HOSPITAL/TEXAS A&M UNIVERSITY: B Manyam, P Simpson, J Whetteckey, B Wulbrecht; THE PARKINSON'S & MOVEMENT DISORDER INSTITUTE: D Truong, M Pathak, N Luong, T Tra, A Tran, J Vo; TORONTO WESTERN HOSPITAL, UNIVERSITY HEALTH: A Lang, G Kleiner-Fisman, A Nieves, J So; UMDNJ-SCHOOL OF OSTEOPATHIC MEDICINE: G Podskalny, L Giffin; UNIVERSITY OF ALABAMA AT BIRMINGHAM: P Atchison, C Allen; UNIVERSITY OF ALBERTA: W Martin, M Wieler; UNIVERSITY OF CALGARY: O Suchowersky, M Klimek; UNIVERSITY OF CALIFORNIA IRVINE: N Hermanowicz, S Niswonger; UNIVERSITY OF CALIFORNIA SAN DIEGO: C Shults, D Fontaine; UNIVERSITY OF CALIFORNIA SAN FRANCISCO: M Aminoff, C Christine, M Diminno, J Hevezi; UNIVERSITY OF CHICAGO: A Dalvi, U Kang, J Richman, S Uy, J Young; UNIVERSITY OF CINCINNATI: A Dalvi, A Sahay, D Schwieterman; UNIVERSITY OF COLORADO HEALTH SCIENCES CENTER: M Leehey, S Culver, T Derian; UNIVERSITY OF CONNECTICUT: T Demarcada, S Belber; UNIVERSITY OF IOWA: R Rodnitzky, J Dobson; UNIVERSITY OF KANSAS MEDICAL CENTER: R Pahwa, K Lyons, T Gales, S Thomas; UNIVERSITY OF MARYLAND SCHOOL OF MEDICINE: L Shulman, S Reich, W Weiner, K Dustin; UNIVERSITY OF MIAMI: C Singer, W Koller, K Lyons, W Weiner, L Zelaya; UNIVERSITY OF MINNESOTA: P Tuite, V Hagen, S Rolandelli, R Schacherer; UNIVERSITY OF NEW MEXICO: P Gordon, J Werner; UNIVERSITY OF PUERTO RICO SCHOOL OF MEDICINE: C Serrano, S Roque; UNIVERSITY OF ROCHESTER: R Kurlan, D Berry, I Gardiner; UNIVERSITY OF SOUTH FLORIDA: R Hauser, J Sanchez-Ramos, T Zesiewicz, H Delgado, K Price, P Rodriguez; UNIVERSITY OF TENNESSEE-MEMPHIS: R Pfeiffer, L Davis, B Pfeiffer; UNIVERSITY OF TEXAS SOUTHWESTERN MEDICAL CENTER: R Dewey, B Hayward, M Meacham; WAKE FOREST UNIVERSITY SCHOOL OF MEDICINE: F Walker, V Hunt; WASHINGTON UNIVERSITY: B Racette, L Good, M Rundle

Biostatistics and Clinical Trials Coordination Center Staff:

D Oakes, A Watts, A Wang, T Ross, S Bennett, Dietgard Kamp, E Julian-Baros.

2 Cincinnati Children's Hospital Medical Center, Cincinnati, OH;

3 University of Rochester, Rochester, NY;

4 University of California, La Jolla, CA; VA San Diego Healthcare System, San Diego, CA;

5 University of Cincinnati College of Medicine, Cincinnati, OH

Abstract

Mutations in *DJ-1* (*PARK7*) are one cause of early-onset autosomal-recessive parkinsonism. We screened for *DJ-1* mutations in 93 affected individuals from the 64 multiplex Parkinson disease (PD) families in our sample that had the highest family-specific multipoint LOD scores at the *DJ-1* locus. In addition to sequencing all coding exons for alterations, we used multiplex ligation-dependent probe amplification (MLPA) to examine the genomic copy number of *DJ-1* exons. A known polymorphism (R98Q) was found in five PD subjects, once as a homozygote and in the other four cases as heterozygotes. No additional missense mutations and no exon deletions or duplications were detected. Our results, in combination with those of previous studies, suggest that alterations in *DJ-1* are not a common cause of familial PD.

Keywords

Parkinson disease; DJ-1; multiplex ligation-dependent probe amplification; MLPA

Introduction

Parkinson disease (PD) is the second most common neurodegenerative disorder. It is characterized by tremor, rigidity, postural instability, and bradykinesia as well as a good response to levodopa therapy [5]. In the past decade, mutations in several genes, including *SNCA*, *PRKN*, *DJ-1*, *PINK1*, and *LRRK2*, have been shown to cause these clinical symptoms [8].

Each of the known PD genes was identified using a positional cloning approach. Linkage to chromosome 1p36 (*PARK7* locus) was initially reported in a large consanguineous Dutch kindred [25]. Subsequently, deletion of exons 1–5 of *DJ-1* was reported in this family [3]. Despite screening at least one individual from nearly 1,200 families [1–4,6,9–14,17–19,21–22,24], only six families have been reported to harbor disease producing mutations in *DJ-1*. Affected individuals in five of the six families were homozygous for their particular mutation, with two of these families known to be consanguineous. Each of these six families had a unique *DJ-1* mutation. A total of four point mutations and three genomic rearrangements, including partial or entire exon duplication or deletion, have been reported. The clinical presentation of individuals with *DJ-1* mutations is similar to that of patients with homozygous *PRKN* mutations. Patients with mutations in either gene typically have an age of onset under 45 years and a slower disease progression

In the current study, we have ascertained a large cohort of multiplex PD families and have used an approach based on the linkage evidence to chromosome 1p36 to prioritize families for *DJ-1* screening, an approach we have previously used to prioritize families for *PRKN* mutation screening [7]. We have performed extensive screening of *DJ-1*, testing for coding sequence point mutations as well as gene dosage alterations (whole exon deletions or duplications).

Materials and Methods

Subjects

A sample of 287 multiplex PD families was recruited through 59 Parkinson Study Group sites located throughout North America. Families were ascertained through an affected sibling pair, and each participating affected family member underwent a standardized neurological evaluation after appropriate written informed consent approved by each individual institution's institutional review board was completed. The evaluation included the Unified Parkinson's Disease Rating Scale (UPDRS) Parts II and III [20] and a diagnostic checklist which was used to provide both inclusion criteria highly associated with autopsy-confirmed PD and exclusion criteria associated with non-PD pathology [15–16]. Peripheral blood for DNA extraction was obtained from all individuals.

Data from a previously completed genome screen [23] was used to prioritize families based on their linkage to the region containing *DJ-1*. The 93 PD individuals from the 64 families with the highest family-specific multipoint LOD scores (range: 0.1 – 1.2) at this locus were then screened for *DJ-1* mutations. The markers used to calculate the family-specific multipoint LOD scores were those closest to the PARK7 locus (D1S468, D1S214, D1S450, and D1S2667) from the 10 cM genome screen (ABI Prism Linkage Mapping Set, Applied Biosystems, Foster City, CA, USA)

A control sample of 96 subjects was also available. All individuals were clinically evaluated and did not have a diagnosis or symptoms of PD or any other neurological disorder. Fifty-two controls were recruited in Indiana and had an average age at exam of 68.8 years (range 55–82 years). The remaining 44 controls were obtained through the National Cell Repository for Alzheimer's Disease (NCRAD) and had an average age at exam of 76.7 years (range 58–91 years). All control subjects were Caucasian and provided written informed consent.

DNA Sequence Analysis

Coding exons (exons 2–7) of *DJ-1* were PCR amplified using primers designed by the authors (WCN, MWP) (see Table 1). Following an initial denaturation at 95°C for 5 minutes, PCR was carried out for 35 cycles (30s at 95°C, 30s at anneal temperature, 30s at 72°C) followed by a 10 min extension at 72°C. PCR products were purified using the QIAquick 96 PCR purification kit (Qiagen, Santa Clara, CA) and sequenced on both strands using an ABI Prism 3700 DNA analyzer (Applied Biosystems, Foster City, CA) and the Applied Biosystems BigDye Terminator version 2.0 kit. Sequences were aligned and analyzed using Sequencer v. 4.0.5 software (Gene Codes Corporation, Ann Arbor, MI). Sequence variants were identified according to GenBank accession number NM_007262.3 with nucleotides numbered with the A of the ATG initiation codon designated as +1.

DNA Dosage Analysis

Multiplex ligation-dependent probe amplification (MLPA) was performed with 100 ng of genomic DNA according to manufacturer's instructions using the P051 Salsa MLPA Parkinson probe set (MRC-Holland, Amsterdam, The Netherlands). This probe set includes probes for all *DJ-1* exons except 2 and 4. Probe amplification products were run on an ABI 3730xl DNA Analyzer using GS500 size standard (Applied Biosystems, Foster City, CA). MLPA peak plots were visualized using Genemapper Software v3.7 (Applied Biosystems, Foster City, CA). Non-normalized values for peak height and peak area were then exported from Genemapper Software v3.7 to an Excel template. Normalization of data and calculation of dosage ratios was performed as described at www.mrc-holland.com/MLPA%20analysis.html. The control value for each probe was an average of 3 control samples run with the PD samples. Due to variation

in assay performance, we used dosage ratio values of ≤ 0.7 and ≥ 1.35 as our boundaries for deletions and duplications, respectively.

Results

The 93 individuals screened for *DJ-1* mutations were either Caucasian (95%) or Hispanic (5%). The average age of onset of these individuals was 58.0 years (median: 59 years, range: 18–80 years), which is slightly younger than the average age of onset in the full sample (61.2 years). The 93 individuals sequenced represented 64 different families. One affected individual was sequenced from 45 of the families while two affected individuals were sequenced from 11 families and three affected individuals were sequenced from eight families. Some families had more than one individual sequenced due to family properties and genotypes of the markers around the *DJ-1* locus. Three different sequence alterations were identified in seven of the 93 individuals screened for mutations. Two of the alterations were synonymous substitutions (480C > A and 501A > G), and the remaining alteration (293G > A) changed an arginine at amino acid 98 to a glutamine. This amino acid substitution (R98Q) was the most common sequence alteration found in our sample. We identified four PD patients who were heterozygous R98Q and one patient who was homozygous R98Q. Three of the four R98Q heterozygotes were affected members of the same family. The fourth heterozygote was from an unrelated family. The two synonymous substitutions (480C > A and 501A > G) were each found in the heterozygous state in a single individual with PD. No change in exon copy number, either as a whole exon deletion or duplication, was detected for exons 1, 3, 5, 6, or 7 in any of the 93 PD samples screened.

The panel of 96 control subjects was screened for the R98Q amino acid substitution as well. Three of the controls were found to carry this polymorphism in the heterozygous state; none of the controls were homozygous for R98Q. Therefore, the frequency of the R98Q polymorphism in our control sample is 1.6%. The controls were not screened for the two synonymous substitutions since neither is predicted to result in an amino acid change.

Discussion

We have performed thorough screening of *DJ-1* to detect mutations which may cause PD. Rather than screening our entire sample of multiplex PD families, we have employed a more efficient approach that utilized data from our previous genome screen to prioritize families for testing based upon their family specific LOD score at the *DJ-1* locus. We did not identify any gene dosage alterations in our 93 PD cases. We identified only one missense mutation, which has been shown in previous studies to be a benign polymorphism [4,13,22], even in the homozygous state [13]. We found the R98Q substitution at similar frequencies in our cases and controls, further confirming that this polymorphism is unlikely to be disease producing.

Previous studies of *DJ-1* have often limited the scope of screening to detect only coding sequence changes. Only one third of the previously completed studies [6,12,19,21], which represent less than one quarter of the 1200 total patients screened to date, employed methods to detect any alternations in gene dosage. Therefore, it is possible that the frequency of *DJ-1* mutations might have been underestimated in these studies. However, our analyses would suggest that even when a cohort is enriched for families more likely to carry a *DJ-1* mutation and methods include screening for gene dosage alterations, mutations in this gene are rare in familial PD.

There are several notable strengths of our study. First, thorough screening of the *DJ-1* gene was performed, including sequencing of all translated exons as well as testing for genomic rearrangements by MLPA analysis. Second, families were selected for testing based on their

evidence of linkage to the *DJ-1* region on chromosome 1p36. We have previously successfully utilized this approach to prioritize screening of our patient cohort for mutations at the *PRKN* locus [7]. Thus, we have augmented our tested sample for families most likely to have a mutation in *DJ-1*. Conversely, our study also has several limitations. First, due to the likely paucity of *DJ-1* mutations, we did not screen our entire familial PD sample. However, we did screen more than 15% of our sample which included those most likely to harbor a *DJ-1* mutation. Second, we did not screen our entire sample of early onset PD cases, which is typically defined as having an age of onset of 45 or earlier. Rather we used a more cost effective approach to only screen families with evidence of linkage to this genomic region. Finally, the dosage methods did not interrogate either exon 2 or 4. In addition, positive controls carrying DJ-1 exon deletions were not available to us. While false negatives cannot be ruled out, MLPA analysis was performed at least twice for each sample minimizing this risk. However, based on the results of the screening in our sample as well as the results from other studies, mutations in *DJ-1* do not appear to be a common cause of PD, and are much rarer than mutations in the *PRKN* or *LRRK2* genes.

Acknowledgements

This project was supported by R01 NS37167 and the National Cell Repository for Alzheimer's Disease (U24 AG021886). We thank Kathleen Miller and Michele Goodman for their assistance. We thank the subjects for their participation in this research study.

References

1. Abou-Sleiman PM, Healy DG, Quinn N, Lees AJ, Wood NW. The role of pathogenic DJ-1 mutations in Parkinson's disease. *Ann Neurol* 2003;54(3):283–286. [PubMed: 12953260]
2. Annesi G, Savettieri G, Pugliese P, D'Amelio M, Tarantino P, Ragonese P, La Bella V, Piccoli T, Civitelli D, Annesi F, Fierro B, Piccoli F, Arabia G, Caracciolo M, Cirò Candiano IC, Quattrone A. DJ-1 mutations and parkinsonism-dementia-amyotrophic lateral sclerosis complex. *Ann Neurol* 2005;58(5):803–807. [PubMed: 16240358]
3. Bonifati V, Rizzu P, van Baren MJ, Schaap O, Breedveld GJ, Krieger E, Dekker MC, Squitieri F, Ibanez P, Joosse M, van Dongen JW, Vanacore N, van Swieten JC, Brice A, Meco G, van Duijn CM, Oostra BA, Heutink P. Mutations in the DJ-1 gene associated with autosomal recessive early-onset parkinsonism. *Science* 2003;299(5604):256–259. [PubMed: 12446870]
4. Clark LN, Afridi S, Mejia-Santana H, Harris J, Louis ED, Cote LJ, Andrews H, Singleton A, Wavrant De-Vrieze F, Hardy J, Mayeux R, Fahn S, Waters C, Ford B, Frucht S, Ottman R, Marder K. Analysis of an early-onset Parkinson's disease cohort for DJ-1 mutations. *Mov Disord* 2004;19(7):796–800. [PubMed: 15254937]
5. de Rijk MC, Tzourio C, Breteler MM, Dartigues JF, Amaducci L, Lopez-Pousa S, Manubens-Bertran JM, Alperovitch A, Rocca WA. Prevalence of parkinsonism and Parkinson's disease in Europe: the EUROPARKINSON Collaborative Study. European Community Concerted Action on the Epidemiology of Parkinson's disease. *J Neurol Neurosurg Psychiatry* 1997;62(1):10–15. [PubMed: 9010393]
6. Djarmati A, Hedrich K, Svetel M, Schäfer N, Juric V, Vukosavic S, Hering R, Riess O, Romac S, Klein C, Kostic V. Detection of Parkin (PARK2) and DJ1 (PARK7) mutations in early-onset Parkinson disease: Parkin mutation frequency depends on ethnic origin of patients. *Hum Mutat* 2004;23(5):525. [PubMed: 15108293]
7. Foroud T, Uniacke SK, Liu L, Pankratz N, Rudolph A, Halter C, Shults C, Marder K, Conneally PM, Nichols WC, and the Parkinson Study Group. Heterozygosity for a mutation in the parkin gene leads to later onset Parkinson disease. *Neurology* 2003;60(5):796–801. [PubMed: 12629236]
8. Gasser T. Genetics of Parkinson's disease. *Curr Opin Neurol* 2005;18(4):363–369. [PubMed: 16003110]
9. Guo JF, Tang BS, Zhang YH, Xia K, Cai F, Pan Q, Shen L, Jiang H, Zhao GH, Yan XX, Cao L. Mutation analysis of DJ1 gene in patients with autosomal recessive early-onset Parkinsonism. *Zhonghua Yi Xue Yi Chuan Xue Za Zhi* 2005;22(6):641–643. [PubMed: 16331561]

10. Hague S, Rogaeva E, Hernandez D, Gulick C, Singleton A, Hanson M, Johnson J, Weiser R, Gallardo M, Ravina B, Gwinn-Hardy K, Crawley A, St George-Hyslop PH, Lang AE, Heutink P, Bonifati V, Hardy J, Singleton A. Early-onset Parkinson's disease caused by a compound heterozygous DJ-1 mutation. *Ann Neurol* 2003;54(2):271–274. [PubMed: 12891685]
11. Healy DG, Abou-Sleiman PM, Valente EM, Gilks WP, Bhatia K, Quinn N, Lees AJ, Wood NW. DJ-1 mutations in Parkinson's disease. *J Neurol Neurosurg Psychiatry* 2004;75(1):144–145. [PubMed: 14707326]
12. Hedrich K, Djarmati A, Schäfer N, Hering R, Wellenbrock C, Weiss PH, Hilker R, Vieregge P, Ozelius LJ, Heutink P, Bonifati V, Schwinger E, Lang AE, Noth J, Bressman SB, Pramstaller PP, Riess O, Klein C. DJ-1 (PARK7) mutations are less frequent than Parkin (PARK2) mutations in early-onset Parkinson disease. *Neurology* 2004;62(3):389–394. [PubMed: 14872018]
13. Hedrich K, Schäfer N, Hering R, Hagenah J, Lanthaler AJ, Schwinger E, Kramer PL, Ozelius LJ, Bressman SB, Abbruzzese G, Martinelli P, Kostic V, Pramstaller PP, Vieregge P, Riess O, Klein C. The R98Q variation in DJ-1 represents a rare polymorphism. *Ann Neurol* 2004;55(1):145. [PubMed: 14705128]
14. Hering R, Strauss KM, Tao X, Bauer A, Woitalla D, Mietz EM, Petrovic S, Bauer P, Schaible W, Müller T, Schöls L, Klein C, Berg D, Meyer PT, Schulz JB, Wollnik B, Tong L, Krüger R, Riess O. Novel homozygous p.E64D mutation in DJ1 in early onset Parkinson disease (PARK7). *Hum Mutat* 2004;24(4):321–329. [PubMed: 15365989]
15. Hughes AJ, Ben-Shlomo Y, Daniel SE, Lees AJ. What features improve the accuracy of clinical diagnosis in Parkinson's disease: a clinicopathologic study. 1992. *Neurology* 2001;57(10 Suppl 3):S34–S38. [PubMed: 11775598]
16. Hughes AJ, Daniel SE, Kilford L, Lees AJ. Accuracy of clinical diagnosis of idiopathic Parkinson's disease: a clinico-pathological study of 100 cases. *J Neurol Neurosurg Psychiatry* 1992;55(3):181–184. [PubMed: 1564476]
17. Ibáñez P, De Michele G, Bonifati V, Lohmann E, Thobois S, Pollak P, Agid Y, Heutink P, Dürr A, Brice A. French Parkinson's Disease Genetics Study Group. Screening for DJ-1 mutations in early onset autosomal recessive parkinsonism. *Neurology* 2003;61(10):1429–1431. [PubMed: 14638971]
18. Karamohamed S, Golbe LI, Mark MH, Lazzarini AM, Suchowersky O, Labelle N, Guttman M, Currie LJ, Wooten GF, Stacy M, Saint-Hilaire M, Feldman RG, Liu J, Shoemaker CM, Wilk JB, DeStefano AL, Latourelle JC, Xu G, Watts R, Growdon J, Lew M, Waters C, Vieregge P, Pramstaller PP, Klein C, Racette BA, Perlmuter JS, Parsian A, Singer C, Montgomery E, Baker K, Gusella JF, Herbert A, Myers RH. Absence of previously reported variants in the SCNA (G88C and G209A), NR4A2 (T291D and T245G) and the DJ-1 (T497C) genes in familial Parkinson's disease from the GenePD study. *Mov Disord* 2005;20(9):1188–1191. [PubMed: 15966003]
19. Klein C, Djarmati A, Hedrich K, Schäfer N, Scaglione C, Marchese R, Kock N, Schüle B, Hiller A, Lohnau T, Winkler S, Wiegers K, Hering R, Bauer P, Riess O, Abbruzzese G, Martinelli P, Pramstaller PP. PINK1, Parkin, and DJ-1 mutations in Italian patients with early-onset parkinsonism. *Eur J Hum Genet* 2005;13(9):1086–1093. [PubMed: 15970950]
20. Lang, AE.; Fahn, S. Assessment of Parkinson's disease. In: Munsat, T., editor. Quantification of neurologic deficit. Butterworth; Boston: 1989. p. 285-309.
21. Lockhart PJ, Bounds R, Hulihan M, Kachergus J, Lincoln S, Lin CH, Wu RM, Farrer MJ. Lack of mutations in DJ-1 in a cohort of Taiwanese ethnic Chinese with early-onset parkinsonism. *Mov Disord* 2004;19(9):1065–1069. [PubMed: 15372597]
22. Lockhart PJ, Lincoln S, Hulihan M, Kachergus J, Wilkes K, Bisceglia G, Mash DC, Farrer MJ. DJ-1 mutations are a rare cause of recessively inherited early onset parkinsonism mediated by loss of protein function. *J Med Genet* 2004;41(3):e22. [PubMed: 14985393]
23. Pankratz N, Uniacke SK, Halter CA, Rudolph A, Shults CW, Conneally PM, Foroud T, Nichols WC. Parkinson Study Group. Genes influencing Parkinson disease onset: replication of PARK3 and identification of novel loci. *Neurology* 2004;62(9):1616–1618. [PubMed: 15136695]
24. Tan EK, Tan C, Zhao Y, Yew K, Shen H, Chandran VR, Teoh ML, Yih Y, Pavanni R, Wong MC. Genetic analysis of DJ-1 in a cohort Parkinson's disease patients of different ethnicity. *Neurosci Lett* 2004;367(1):109–112. [PubMed: 15308309]
25. van Duijn CM, Dekker MC, Bonifati V, Galjaard RJ, Houwing-Duistermaat JJ, Snijders PJ, Testers L, Breedveld GJ, Horstink M, Sandkuijl LA, van Swieten JC, Oostra BA, Heutink P. Park7, a novel

locus for autosomal recessive early-onset parkinsonism, on chromosome 1p36. *Am J Hum Genet* 2001;69(3):629–634. [PubMed: 11462174]

Table 1

Primers used for PCR amplification

Exon	Forward Primer	Reverse Primer
2	TCAGTTGTCTATGAAAACCG	GAGCAGTGGATCAACAGGTA
3	AGGATCATTGAGCCAGGA	TCCTCACCTCTTAATCTGT
4	GTCTATTGGCAGATAGGCT	ACAGCCTCCTCCCGAAATAT
5	ATAGGTCAGAGAGCTTGTGG	CAGATGCCCTGCATGCTTTA
6	GCCAGGCACTATTGCGATTT	AGAAGAATCGCTTGAACCCG
7	CTTGGAGTGCCTAGTAAATG	TTCAGTGCCATATGGACAGT