Antibiotics in surgical treatment of acute abscesses

SIR,—I am pleased to see that primary closure, under antibiotic cover, of acute superficial abscesses continues to be advocated (12 July, p 111). However, it is surprising that the investigators of the trial prescribed a five-day course of clindamycin for one-quarter of their patients, since the Committee on Safety of Medicines, in June 1979, recommended that this drug "should in general be reserved for serious or life-threatening conditions where other antibiotic therapy is ineffective or undesirable."¹ This recommendation was made because of the well-known association between clindamycin and antibiotic-associated colitis.

Dr P W H Blick and others have, however, shown that a single preoperative dose of clindamycin provides adequate antibiotic cover when closing an abscess cavity by primary suture. This is an important finding as a single-dose method might reduce the (admittedly small) risk of inducing pseudomembranous colitis to infinitesimal proportions and allow the method of primary closure, with this antibiotic, to be advocated widely once again.

While I too have found that the time needed for skin healing after incision, curettage, and primary suture under antibiotic cover is rather less than that after incision and open drainage, I believe that the main advantage of the former technique is that it allows the patient to return to work sooner; there is no need for him to remain at home for nursing visits or to make daily trips to hospital for dressings.²

JOHN MURIE

Department of Surgery,
Kilmarnock Infirmary,
Kilmarnock KA3 2DJ

¹ Committee on Safety of Medicines. *Adverse reaction series*, No 17. London:HMSO, 1979.

² Murie JA. *J R Coll Surg Edinb* (in press).

*We sent this letter to the authors, and Mr Wilson's reply is printed below.—ED, BMJ.

SIR,—I do not think that Mr Murie would want his comments about our choice of antibiotic to detract from the main purpose of the paper. This was to report our findings that a single preoperative dose of an effective antibiotic is sufficient chemotherapy to permit primary closure of an acute abscess after thorough curettage, and that a course of postoperative antibiotic therapy does not further enhance the healing process.

However, referring to the Committee on Safety of Medicine's circular of June 1979, Mr Murie expresses surprise that clindamycin was the antibiotic which we used to establish this surgical principle. On reflection I am sure he will appreciate that the protocol for the trial had been prepared much earlier than June 1979. In fact, the trial was almost complete by the time the letter was issued. Nevertheless, I must report that, despite the Committee's letter, we still use clindamycin for the treatment of acute soft tissue infections because of the excellent clinical results it provides for our patients. This decision is based on a five-year study of 15 019 patients treated in our own department, which revealed that only 1 in 1000 patients developed diarrhoea of five or more motions in 24 hours, and in only two of

these patients were there symptoms of pseudomembranous colitis.¹

Most of the fatalities from pseudomembranous colitis associated with clindamycin therapy have been in elderly, debilitated patients with severe chronic illness. Many of them were already suffering from diarrhoea before clindamycin or lincomycin was added to their therapy in a forlorn hope of reversing terminal symptoms. These unfortunate patients are in stark contrast to the ambulant, young patients (mean age 26 years) who attend our department suffering from an acute soft tissue infection.

In view of the recent advances in our understanding of the nature of antibiotic-associated colitis, especially the role of *Clostridium difficile*, and the discovery of vancomycin and possibly metranidazole as specific therapeutic agents, we feel justified in continuing to use this valuable antibiotic for the treatment of acute soft tissue infections, but with certain reservations. It should not be used for patients who have chronic colitis and it is possibly unwise to use it for elderly, frail patients. Provided that any young, otherwise healthy patient who develops severe diarrhoea seeks immediate treatment, we consider the risk to be minimal.

D H WILSON

Accident and Emergency Department,
General Infirmary at Leeds,
Leeds LS1 3EX

¹ Wilson DH. *Br J Surg* 1980;67:93-6.

Jejunioleal tuberculosis

SIR,—I was pleased to see our article in the *BMJ* (12 July, p 118). It has been brought to my attention, however, that we have not properly acknowledged the help which we received from our pathologist colleague in the preparation of the figure, and I would just like to take this opportunity to thank him for his expert assistance (Dr C Burrows, consultant pathologist, Walton Hospital, Rice Lane, Liverpool 9).

P WAKE

Royal Liverpool Hospital,
Liverpool L7 8XP

Genital herpes

SIR,—Your recent excellent leading article (7 June, p 1335) on genital herpes contained two statements worthy of comment: "Genital herpes is commoner than primary syphilis, from which it must be differentiated" and "Syphilis must be excluded." The implication is one of mutual exclusiveness. In fact, sexually transmitted diseases frequently coexist, and even when virologically proved genital herpes has been diagnosed all genital ulcers must be submitted to dark-ground microscopy. The fluorescent treponemal antibody absorption test can be misleading. False-positive results occur in genital herpes,¹ and if dark ground microscopy is not used cases of syphilis may be misdiagnosed.

Occam's razor does not apply in venereology. The presence of one sexually transmitted infection implies either promiscuity or intercourse with a promiscuous partner, and promiscuity implies the risk of acquiring multiple infections. Multiple causes of venereal infection must be actively sought as many of these infections may be asymptomatic. In one series² of virologically proved genital herpes

seven also had gonorrhoea, two had trichomoniasis, three had candidiasis, and one had warts. In another series¹ of 32 virologically proved cases of genital herpes one had gonorrhoea, one had non-specific genital infection, two had trichomoniasis, and two had candidiasis.

It therefore follows that the management of genital herpes should include an active search for any associated, possibly asymptomatic, and potentially damaging sexually transmitted disease.

D P MURRAY

British Military Hospital,
Münster, West Germany

¹ Wright JT, Cremer AW, Ridgway GL. *Br J Vener Dis* 1975;51:329-30.

² Beilby JOW, Cameron CH, Catterall RD. *Lancet* 1968;i:1065-6.

Children in wheelchairs

SIR,—In Contemporary Themes the important topic of children in wheelchairs was discussed by Alison Wisbeach and Professor K S Holt (19 July, p 199). Much of what was stated applies equally to the adult wheelchair users in corresponding categories. May I draw the attention of your readers to that excellent book by Dr Phillipa Russell,¹ which I have recently read, because it gives a first-hand, comprehensive account of the care of children in wheelchairs and enables one to appreciate the problems by putting them in perspective. Although I am myself an adult in a wheelchair, I learned much from the author, who I understand is the mother of a wheelchair child. From a technical standpoint much more can be gained from the proceedings of the seminar on rehabilitation of the disabled held at the University of Strathclyde, Glasgow, in 1978, one of the Strathclyde bioengineering seminars. It appears that there is indeed more co-ordination between the workers caring for the disabled in Scotland than is apparent in England and, in addition, more educational facilities for those who elect to do this work—for instance, a National Centre for Training in Prosthetics and Orthotics. With the Year of the Disabled in mind (1981) may I suggest that such an educational project be made available in this country.

MABEL L HAIGH

Wetherby, W Yorks LS22 5BX

¹ Russell, P. *The wheelchair child*. 2nd ed. London: Souvenir Press, 1978.

Seating the disabled

SIR,—We were delighted to read the leading article on seating the disabled (5 April, p 963), which raises the following points.

(1) The complex factors that maintain stability and protect from injury—that is, promote tissue viability—have received a good deal of attention. It may seem otherwise because the published work is scattered in time and space, across many disciplines (not necessarily medical), in various specialist journals. General journals such as yours can correct this.

(2) You say in your editorial, "As is so frequently the case, no single organisation has clear responsibility for initiating and seeing through the creation of a seat support for the disabled." Seating includes cushions and chairs and is one part of "support systems." Support systems form one aspect among many of the