are exceptionally difficult to make from geographical data.³

(3) Furthermore, the analysis of data for towns need not provide any useful guide to the results obtained by analysing the same data at the individual level. The aggregation of data for individuals into towns creates all manner of spurious geographical associations, which statistical techniques are very adept at identifying.

(4) Statistical analysis of geographical data suffers from a number of problems that have been completely ignored. In this respect the study repeats many of the mistakes that characterised geographical studies in the 1960s.

(5) Excessive emphasis is placed on testing the parameters of linear regression models for significance. Unless the data were collected by simple random sampling, which they were not, significance tests are meaningless. Statements such as "... a highly significant regression coefficient ($p < 0.001$)" are totally misleading.

(6) To make matters worse, it is recognised that the residuals are spatially autocorrelated: "Some towns ... had a higher cardiovascular mortality than the model predicted ... such towns tended to occur in geographic clusters." Yet the serious consequence this has for regression models is overlooked.⁴

(7) Many other aspects of the statistical analysis are unsatisfactory. For example, no indication of the form of the data frequency distributions is given; it is admitted that the relationship is really non-linear, so there is specification error; the choice of variables appears indiscriminate, as is also the global logarithmic transformation; and there may well be multicollinearity problems owing to the use of highly intercorrelated variables in regression models.

(8) No attempt is made to account for the surprisingly high correlations between cardiovascular disease and both latitude and longitude ($r = +0.74$, $+0.68$). If correlations of this magnitude occur for what appear to be nonsense relationships, what hope is there for finding a real water quality effect ($r = -0.67$)?

(9) Finally, the models used are poor from a theoretical point of view. They assume that people who died in 1969-73 have experienced all their lives a uniform water quality of the type that occurred in the town where they happened to be living when death occurred.

Few would deny that geographical mapping of medical data can reveal all kinds of interesting distributional patterns. However, it is essential that the analyses of these patterns are carried out with an appropriate degree of rigour. Regrettably it appears that medical researchers often lack sufficient experience of, or have no expertise in, geographical analysis. The availability of geographical data and statistical packages seems to have encouraged all manner of madness.

STAN OPENSHAW
MARTIN CHARLTON

University Department of
Town and Country Planning,
Newcastle upon Tyne NE1 7RU

¹ Openshaw S. *Trans Inst Br Geogr* 1977;2:459-72.

² Openshaw S. *Env Plann A* 1978;10:781-94.

³ Blalock HM. *Causal inferences in non-experimental research*. North Carolina: Chapel Hill, 1964.

⁴ Cliff AD, Ord JK. *Spatial autocorrelation*. London: Pion, 1973.

* * We sent this letter to the authors, and Dr Pocock and Professor Shaper reply below. —Ed, *BMJ*.

SIR,—Thank you for the opportunity of replying to the comments on our British regional heart study made by the Newcastle geographers.

We studied towns because they are eminently sensible units for studying geographical variations in cardiovascular mortality. Larger units such as the English regions are few in number and too

heterogeneous, while smaller units such as wards have populations too small to provide accurate estimation of mortality rates. Towns tend to be fairly homogenous in climate and water quality but less so in socioeconomic conditions. However, our field work experience in 25 towns in England, Wales, and Scotland leaves us convinced that quantifying socioeconomic conditions on a town basis is a valuable contribution to our study. Towns may not be theoretically perfect units but it seems reasonable to deal with real problems in a practical framework. We were most concerned about the problem of internal heterogeneity of towns from the point of view of their water supplies and this was carefully investigated. The conclusions of the study are robust to the exclusion of those towns whose water supplies are more variable. Migration between towns is not on a large enough scale to bias the study, and water quality in most towns has been stable over many decades. We have no information from which to draw conclusions regarding the duration of exposure required to produce a cardiovascular effect.

We describe geographical associations in order to attempt to explain which socioeconomic and environmental factors are related to cardiovascular mortality. Of course we realise that causal inferences are not readily sustained by this type of data, which is precisely why we have taken so much care to determine whether the observed relationship between water hardness and cardiovascular mortality could be attributed to other factors.

We are conscious of the differences between group-based and individual-based data. However, disagreement between these two types of data does not necessarily invalidate the findings within either type of data. But agreement may occur and our study shows an association between socioeconomic factors and cardiovascular mortality which is also seen in many individual-based studies. Phases 2 and 3 of the regional heart study are evidence of our concern to produce more specific cross-sectional and prospective data on cardiovascular risk factors by using individuals rather than towns.

Statistical tests of significance are useful in assessing whether the relationship between two variables can be attributed to chance. Observational studies like ours, which by their nature preclude random sampling, obviously require caution in their interpretation. Statements such as "... highly significant regression coefficient ($p < 0.001$)" should not be misleading to those who understand the statistical method and its function as an objective guideline to rational thought.

We have spent considerable time investigating the impact of spatial autocorrelation on our regression models. This required the development of new statistical theory as models previously used in this area appeared implausible. We have now completed analysis using these new models and, as anticipated, the results and conclusions in our paper are not altered to any important degree. These new methods and results will be submitted to a more technical journal in the near future.

We have shown that geographical variations in cardiovascular mortality (including correlations with latitude and longitude) can be related to water hardness, climate, and socioeconomic factors. No single observational study can establish causal relationships but the association between water hardness and cardiovascular mortality has been reported in many different countries. On the available evidence, we consider it reasonable to conclude that some factor in drinking water closely related to hardness is directly affecting cardiovascular mortality.

Dr Openshaw and Mr Charlton are clearly disturbed by our report on the regional heart study, as evidenced by their somewhat intemperate comments. We have spent considerable time in discussion over many years with geographers, geologists, geochemists, and others concerned with environmental studies and we will continue to do so. We welcome comment on the regional heart study but would suggest that in future it be phrased

in the usual scientific language of rational discussion.

S J POCOCK
A G SHAPER

Department of Clinical
Epidemiology and General
Practice,
Royal Free Hospital School of
Medicine,
London NW3 2QG

Cholesterol and mortality rates

SIR,—Mr A R P Walker (31 May, p 1320) correctly warns that only total mortality can be used as an end point to test the value of community-based intervention. However, both he and Dr Robert Beaglehole and others (2 February, p 285) fail to see the significance in the observed "inverse relation between serum cholesterol level and overall mortality" in the New Zealand Maoris. The relation between serum cholesterol and mortality is a U-shaped curve when a human population has a life style which allows a wider range of cholesterol levels. One prospective study of United States adults published over a decade ago showed the lowest overall mortality at a cholesterol level of 5.83-6.45 mmol/l (225-249 mg/100 ml), with higher mortality above and below.¹ It appears that the Maoris represent only the lower limb of this U-shaped curve. Obviously, a change in their habits which would allow a wider range of values would show a similar U-shaped curve. A similar U-shaped mortality curve can be seen with a wide range of body weights.² Extremes of leanness and obesity are more hazardous than an average weight.

We have observed that cardiac patients in marathon training can easily move their body weights and cholesterol levels down to extremely low values in a short time, and we have expressed concern over the higher risk these low values may represent.³ The necropsy findings differ at these extremes of body weight and cholesterol. Complicated atherosclerosis is more common at the upper end of the scale of obesity and elevated serum lipids; however, rhythm deaths are more common at the lower extremes of underweight and low lipids. Since dietary restrictions were common among the cases of rhythm death we have seen, we have been using the term "nutritional arrhythmias" for these cases. We caution that intervention efforts directed against one disease process should not increase overall mortality by introducing a new risk.

Necropsy studies offer a more precise method of classifying deaths. In the future we should try to separate the complicated atherosclerotic deaths (thrombosis, infarction, etc) from the rhythm deaths, which show no acute lesions.

T J BASSLER

Centinela Hospital,
Inglewood, California 90307,
USA

¹ Stamler J, Lilienfeld AM. *Circulation* 1970;42A:55-95.

² Sorlie P, Gordon T, Kannel WB. *JAMA* 1980;243:1828-31.

³ Bassler TJ. *JAMA* (in press).

Heart disease in different ethnic groups

SIR,—In your leading article about heart disease in different ethnic groups (16 August, p 469) you raise the question "What do the ethnic differences mean?" In reply you refer to "a large genetic component" but add that

this might not be the whole story: "The answer may well be more complicated." A very recent review¹ of the epidemiology of heart disease supports both comments.

In the course of the review I analyse the sex and age patterns of mortality from "ischaemic heart disease" (8th ICD 410-414) in five populations: England and Wales, Scotland, US whites, US non-whites, and Japan. Defined in this way, ischaemic heart disease is heterogeneous, comprising three main components. Most deaths over the age range 30-75 years can, however, be attributed to one component, "acute myocardial infarction" (8th ICD 410). I deduce that genetic factors play two distinctive roles where acute myocardial infarction is concerned. Firstly, genes determine whether or not a person is at risk to acute myocardial infarction but, once the disease process has started, other genes are the main factors involved in determining its average duration, or latent period, from initiation to death. In males the average latent period ranges from about seven years in US non-whites to 13 years in England and Wales; in females these intervals are twice as long. Seasonally dependent environmental factors can exacerbate pathogenesis and hasten death but they have little influence on the average duration of the latent period.

A most interesting feature of the epidemiology of ischaemic heart disease, and one that has attracted much attention, is the recent fall in mortality recorded in several countries, including the United States. By comparing the secular trends in overall mortality with those recorded for mortality from ischaemic heart disease in US whites and US non-whites, I conclude that the fall is probably largely or wholly genuine. If it is, then the characteristics of the decrease in relation to sex and age implicate one or more extrinsic precipitators, whose impact has been declining for a decade or more. Classical precipitators are infective micro-organisms and allergic agents.

The evidence can be readily interpreted in terms of autoaggressive theory.¹ Once random intrinsic changes have occurred in a genetically predisposed person to produce a mutant stem cell, a precipitator appears to be needed to trigger the growth of a forbidden clone, the peripheral cells of which attack the endothelial lining of coronary arteries. A search for the precipitator or precipitators of ischaemic heart disease might well prove rewarding.

My analysis implies, I should add, that the "risk factors"—hypertension, hypercholesterolaemia, obesity, smoking, and lack of exercise, have little or no causal action; their association with ischaemic heart disease appears to be predominantly genetic in origin.

PHILIP R J BURCH

Department of Medical Physics,
General Infirmary,
Leeds LS1 3EX

¹ Burch PRJ. *Cardiovasc Res* 1980;14:307-38.

Hyperlipidaemia advances and retreats

SIR,—As co-ordinator of the Persantine (dipyridamole) Aspirin Reinfarction Study and as co-principal co-ordinator of the Aspirin Myocardial Infarction Study referred to in your leading article entitled "Hyperlipidaemia advances and retreats" (2 August, p 340), I would like to offer the following corrections to

the statement that "Both the Aspirin Myocardial Infarction Study and the Persantine Aspirin Reinfarction Study had failed to show any benefit, and the main positive finding was a surprisingly high incidence of aspirin-related side effects." The Aspirin Myocardial Infarction Study showed no benefit in total mortality over a period of 42 months but did show modest benefits in coronary mortality and coronary incidence that were not statistically significant.¹ The Persantine Aspirin Reinfarction Study showed favourable effects both for the pure aspirin group and for the combination group for total mortality and coronary mortality of 17% and 25% respectively.² By our conservative statistical standards—that is, $z \leq 2.6$ ($p \leq .01$)—these differences were not significant. For coronary incidence, however, the Persantine plus aspirin combination was highly significantly different from placebo for the first two years of treatment for the endpoint coronary incidence (coronary mortality plus definite non-fatal myocardial infarction). Aspirin alone, while less effective, showed significant differences for the same endpoint at months 8 and 24 of treatment.

Side effects due to a daily dose of 1 g of aspirin were indeed not inconsiderable, mainly owing to increased bleeding tendency and some laboratory indications of kidney damage. However, no patient's life was lost because of that. Side reactions attributable to Persantine were limited to an excess of headaches.

C R KLIMT

Maryland Medical Research Institute,
Baltimore, Maryland 21210, USA

¹ Persantine-Aspirin Reinfarction Study Research Group. *Circulation* 1980;62:449-61.

² Aspirin Myocardial Infarction Study Research Group. *JAMA* 1980;243:661-9.

Sodium and potassium intake and blood pressure

SIR,—One of the conclusions of Dr Peter Brustyn and colleagues (23 August, p 537) was that small reductions of sodium intakes by Western populations are unlikely to lower blood pressure levels. Although agreeing with this opinion it should be pointed out that this conclusion was not derived from their experiments, for these did not include the decreasing of sodium intakes.

A recent comprehensive review¹ of population sodium intakes suggested four groups having a different range of consumption levels: (1) Very low sodium intakes, less than 30 mmol sodium (NaCl 2 g) per day, consumed only by hunter gatherers and primitive peasant groups who add no salt to their food; among them blood pressures do not rise with age and essential hypertension is virtually unknown. (2) Customary sodium intakes, ranging from 90 to 200 mmol per day, as in all Western and almost all Eastern populations who add salt to their food; among them blood pressures rise with age. This range of sodium intake permits the expression of hypertension in the genetically susceptible minority, roughly 10-20% of middle-aged persons. (3) Very high sodium intakes of 600-800 mmol per day; these levels have been consumed only for a few days as experimental procedures; they cause blood pressure levels to rise. This review had less to report concerning group 4, the intermediate group having sodium intakes ranging from 30 to 90 mmol per day—that is, intermediate in sodium consumption between groups 1 and 2. This intermediate level of sodium intake has been reported in Samburu and Maasai herdsmen who drank much sodium-containing milk, in Tarahumara Mexican Indians, in Ojong Java Pacific islanders, in Hadza Tanzanian newly

settled nomads who took salty drinking water, and formerly in Kikuyu Kenyan women who added plant ashes from brackish swamps to a no-added-salt diet.² All these groups had virtually no hypertension and no rise of blood pressure in adult life.

Considerable interest must therefore be taken in the results of experiments to study hypertensive patients who had eaten the customary sodium intakes of group 2 and were switched to the lower (intermediate) sodium intakes of group 4. They were not switched to the intolerable low sodium levels of group 1; these are known to reduce hypertensive blood levels. Three reports have recently stated that the majority of essential hypertension patients who had intermediate sodium intakes reverted to normal diastolic blood pressures and ceased all hypotensive drugs, usually within four to eight weeks, while consuming 50-70 mmol of sodium a day.³⁻⁵ These diets involved the exclusion of a short list of highly salted foods. Patients came to accept the change, and even refused to revert to their previous dietary habits when offered the chance; eventually most came to dislike highly salted foods. It is hoped that these preliminary data will be assessed in more sophisticated trials and also will investigate protective factors such as potassium intake, supported by one reference to an experimental study in the review.¹

HUGH TROWELL

Woodgreen,
Nr Fordingbridge, Hants SP6 2AZ

¹ Tobian L. *Am J Clin Nutr* 1979;32:2739-48.

² Trowell HC. *E Afr Med J* 1980;57:167-73.

³ Magnani B, Ambrosioni E, Agosta R, Racco F. *Clin Sci Mol Med* 1976;51, suppl:625-6.

⁴ Dodson PM. *Br Med J* 1980;280:564.

⁵ Mannerberg D. *Chest Heart Stroke J* 1979;3:62-5.

Three types of health education

SIR,—Ever since St Gregory the Great ("Non angli sed angeli"), it has been traditional in this country to debate classifications proposed by others. In this vein may we question the typology of health education proposed by Dr Peter Draper and others (16 August, p 493), which we find confusing? Surely it is important to distinguish clearly the target group, the individual or society, from the objective—be it primary, secondary, or tertiary prevention. Rather than three types of health education, conceptually are there not a minimum of six?

Whatever the terminology, we are in total agreement that "social" rather than "personal" health education, promoting social responsibility in policy affecting health, is an unexploited activity. Part of the reason may be that the ground rules have changed. One-man campaigns in the nineteenth century public health tradition are insufficient against the increasingly resourceful forces at work today. New approaches are certainly required and perhaps one should take the form of highly sophisticated regional initiatives.

The authors call for a reappraisal of the most effective means of organising health education. In Wessex a multidisciplinary regionwide positive health team has just been formed on an experimental basis. Utilising the media to the full, it aims to emphasise individual and collective responsibility for good health. As a first exercise, it has investigated the effectiveness of the very best "personal" health education in promoting the voluntary use of car seat belts in a defined community.¹ The disappointing response observed confirms the need for "social" health education