norphine 3 µg and fentanyl 1 µg/kg body weight peroperatively with minimal intravenous induction agents and halogenated vapours. Postoperatively I give papaveretum 10-20 mg intramuscularly as soon as full consciousness returns but before severe pain is felt.³ Occasionally severe spasms of pain require injection of orphenadrine.⁴ This technique provides analgesia comparable, as assessed by hourly pain scores, with continuous infusion. There was no detectable depression of respiration measured by counting, by Wright's respirometer, or by random blood gas samples.

E N S FRY

Department of Anaesthesia,
North Tees General Hospital,
Stockton-on-Tees,
Cleveland TS19 8PE

¹ Fry ENS. *Br Med J* 1976;ii:817.

² Fry ENS. *Ann R Coll Surg Engl* 1979;61:371-2.

³ Fry ENS. *Br J Anaesth* (in press).

⁴ Fry ENS. *Br J Anaesth* 1977;50:841.

SIR,—Too much narcotic analgesic, like too little gruel, is not a good thing. The opportunity to ask for less would have been of benefit to the group of patients given continuous papaveretum infusions by Dr J A Catling and others (16 August, p 478).

Disturbances in respiration, apnoeic episodes, and low respiratory rates seen in five of seven patients, combined with high overall pain scores suggest that constant infusion results in either overdosing or underdosing. This is an inevitable consequence of inflexible administration techniques, where analgesic requirements for up to 48 hours postoperatively are calculated on the effect of a single loading dose given early in the postanesthetic period. No allowance can be made for the varying requirement for analgesia which come with elimination of anaesthetic agents, the effect of antiemetic drugs, physiotherapy, or the onset of sleep.

The only logical way to control analgesic dosage is by reference to the degree of pain felt by the patient. While on-demand systems have been shown to result in lower overall pain score it is now important to know whether patients using systems like the Cardiff Palliator are demonstrably "better" using the respiratory parameters which Catling and others have defined.

P J SLATTERY
M D VICKERS

Welsh National School of Medicine,
Department of Anaesthetics,
Cardiff CF4 4XN

Impotence in diabetic and non-diabetic hospital outpatients

SIR,—Dr Eva Lester and others (2 August, p 354) demonstrated a 23% incidence of impotence in 83 male diabetic outpatients as compared with a similar incidence of 20% in 50 non-diabetic outpatients. This would seem to indicate that the high incidence of impotence in diabetic men previously claimed has been exaggerated.

Recently we have shown that hypoxia in men with chronic obstructive airways disease is associated with low testosterone values¹ and that the degree of testosterone depression is related to the severity of arterial hypoxia.² The majority of men thus affected are impotent and lack early morning penile erections.³ In view of this and the fact that

patients with chronic obstructive airways disease are one of the largest medical outpatient groups it would be of interest to know from which non-diabetic clinics Dr Lester and her colleagues took their "controls." For if these included a significant proportion of men with chronic obstructive airways disease it might suggest that diabetic men are indeed more frequently impotent than non-diabetic patients excepting those with hypoxia and chronic obstructive airways disease.

PETER D'A SEMPLE
G H BEASTALL
ROBERT HUME

Chest Clinic,
Inverclyde Royal Hospital,
Greenock PA16 0XN

¹ Semple Pd'A, Watson WS, Beastall GH, et al. *Thorax* 1979;34:783-8.

² Semple Pd'A, Beastall GH, Watson WS, Hume R. *Clin Sci* 1980;58:105-6.

³ Semple Pd'A, Beastall GH, Hume R. *Br J Sex Med* 1980;62:48 and 53.

* * * We sent a copy of this letter to the authors, whose reply is printed below.—ED, *BMJ*.

SIR,—Our 50 controls came from a variety of medical and surgical clinics. None of them were from the large and active chest clinic here. Although we did not record detailed clinical findings in the controls it is therefore very unlikely that a significant number had chronic obstructive airways disease.

We do feel that the prevalence of impotence in diabetics has been exaggerated. The figure of 55% which is quoted from paper to paper is an overall figure from a study of 198 men of whom 111 were under 60 years and 87 were aged 60 to 90 years.¹ The prevalence of impotence in the men under 60 years in this study was 26.4% which is very similar to our own figure of 23%.

The findings of Dr Semple and his colleagues of impotence in men with chronic obstructive airways disease and severe hypoxia confirm our feeling that "there is a lot of it about" in men who are unwell. Although four of their seven patients with chronic obstructive airways disease and erectile difficulty had low or low normal testosterone concentrations, they found normal gonadotrophin concentrations so that evidence for a hypothalamic-pituitary mechanism is lacking. This seems to indicate that the question as to whether impotence in patients attending hospital with a variety of disorders including diabetes mellitus is organic, functional, or perhaps even "normal" remains open and we cannot claim to have answered it.

EVA LESTER
F J WOODROFFE

Diabetic Clinic,
North Middlesex Hospital,
London N18 1QX

¹ Rubin A, Babbott D. *JAMA* 1958;168:498-500.

Is methotrexate carcinogenic?

SIR,—We wish to comment on the assertion of Mr R S Knowles and Mr J E Virden (30 August, p 589) that methotrexate is carcinogenic.

Five comprehensive methotrexate toxicology studies in different species of laboratory animals have failed to demonstrate carcino-

genic activity.¹⁻⁵ Roschlau and Justus found that methotrexate induced lung tumours in mice.⁶ These findings are questionable, however, since there were no concurrent untreated controls and spontaneous tumour development in the species used is well documented.

While a number of case histories describing tumour occurrence after methotrexate treatment for psoriasis and one report of a hepatocellular carcinoma found at autopsy in a patient with acute lymphoblastic leukaemia have appeared in the literature, definitive cause-and-effect judgments cannot be made on the basis of isolated reports, especially if one considers that some 25% of the population may be expected to develop malignant disease in the course of their lives. Analysis of a series of 205 psoriasis patients treated with methotrexate for periods of two to seven years with a minimum follow-up of seven years showed no increase in the incidence of internal tumours over a matched population control.⁷ Furthermore, Bagshawe has reported only one second tumour in over 400 methotrexate-treated choriocarcinoma patients followed up since the late 1950s⁸; similarly follow-up of choriocarcinoma patients treated at the United States National Institutes of Health over the last 22 years shows no evidence of increased susceptibility to develop malignant disease after successful chemotherapy, which presumably included methotrexate.⁹

Logically, one would not expect methotrexate to be carcinogenic since, unlike most cytotoxic drugs, it neither reacts chemically with, nor is incorporated into, nucleic acids.

In view of the substantial animal toxicology data and clinical evidence, we submit that it is unjustified to describe methotrexate as carcinogenic and that it should be considered a non-carcinogenic drug.

COLIN TURNBULL
MARYANNE ROACH

Lederle Laboratories,
Gosport, Hants PO13 0AS

¹ Ryan TJ, Buddington MM, Spriggs AI. *Br J Dermatol* 1965;77:541-55.

² Von Schmoll D, Osswald H. *Arzneim Forsch* 1970;20:1461-7.

³ Rustia M, Shubik P. *Toxicol Appl Pharmacol* 1973;26:329-38.

⁴ Data on file, Lederle Laboratories.

⁵ Weisberger EK. *Cancer* 1977;40:1935-49.

⁶ Roschlau G, Justus J. *Dtsch Gesundheitsw* 1971;26:219-22.

⁷ Bailin PL, Tindall JP, Roenigk HH, Hogan MD. *JAMA* 1975;232:359-62.

⁸ Bagshawe KD. *Proc Roy Soc Med* 1977;70:303-6.

⁹ Lewis JL. *Am J Obstet Gynaecol* 1980;136:163-72.

Rugger player's ear

SIR,—Your reply to the question "what is rugger player's ear?" (Any Questions, 16 August, p 498) could perhaps have mentioned two other conditions in addition to the "cauliflower ear" described. Chondrodermatitis helices nodularis is generally considered to be caused by repeated minor trauma to the ear, and I have seen the typical painful papules of this condition in four patients who had played rugby football until after the age of 40. All of them had been forwards and therefore involved with the scrummage, and there was no other likely cause of trauma in their cases. In addition, they all had cauliflower ears to a greater or lesser degree.

More commonly, I have seen multiple herpes simplex lesions on the ears and adjacent areas of neck and scalp in several medical students following rugby football