the first time in weeks, and even go out for meals with his friends, for the simple reason that his pain was effectively controlled. He went home (a prospect which had seemed impossible) and spent 10 good weeks there before his final admission.

Dr Lloyd also includes dextromoramide and pethidine in his chart. These are both very short-acting and strongly habituating drugs, which many of us feel have little or no place in the management of terminal pain.

Please don't let us withhold effective pain relief from this group of patients, for whom after all there is often little else that we can offer medically. And let us dispel all myths about these drugs and ensure they are used not only effectively but in good time by the coming generation of doctors on the wards and in the community.

LESLEY A M WILLS

Haslemere, Surrey

¹ Saunders C. *Nursing Times* July 22 1976: 1133.

² Twycross RG. *Int J Clin Pharmacol* 1974;93:184.

³ Wills LAM. *Postgrad Med J* 1978;54:391-4.

A better system for polio vaccination in developing countries?

SIR,—I read the paper by Dr T Jacob John and others (23 August, p 542) with interest, especially the findings that the oral vaccines tested seemed to have very low potency.

The authors state, "Two batches of imported oral polio vaccine were purchased from the supplier and brought to the laboratory in insulated containers containing ice. They were stored at -20°C and tested for potency just before use. . . ." It is well known that freezing polio vaccine largely inactivates it. This alone could have explained the low potency discovered. I am familiar with the oral vaccine manufactured by Wellcome, which clearly states that freezing is harmful to the vaccine. I assume that other manufacturers also issue the same warnings. This would make the cluster vaccination technique even more impressive than it would have been had Dr John and his colleagues used vaccine which had not been frozen.

DION BELL

Department of Tropical Medicine,
Liverpool School of Tropical
Medicine,
Liverpool L3 5QA

Conversion of thyroxine to triiodothyronine

SIR,—We have read with interest the report of Dr Paul Heyma and colleagues (5 July, p 24), who suggested that the effect of propranolol on thyroxine (T4) conversion to triiodothyronine (T3) was related not to its β -adrenergic blocking activity but to its membrane-stabilising effect. Dr Jones and others in a subsequent letter (p 453) reported that they were unable to show any alteration in T4 conversion to T3 with another (albeit less pronounced) membrane-stabilising β -blocker, oxprenolol. Our recent work with a β_2 -agonist, salbutamol, may help to clarify the problem.

Six healthy females aged 26.7 ± 2.4 years (mean $\pm$ SEM) and weighing 60.3 ± 2.1 kg (an average 8.5% above ideal weight) were given orally 4 mg of salbutamol six hourly for seven days. Serum T4 (Amersham RIA PEG Kit 1M921), T3 (Amersham RIA PEG Kit 1M961), and reverse T3 (Serono Kit 1834) were measured before and at

three, five, and seven days during salbutamol medication. Weight did not alter and serum T4 showed no change. Serum T3 and the T3:T4 ratio showed a slight upward trend but this was not statistically significant. However, both serum reverse T3 and the ratio of reverse T3 to T4 significantly decreased. This suggests that a β_2 -agonist has little action on conversion of T4 to T3 but can alter reverse T3 metabolism.

The fall in the ratio of reverse T3 to T4 together with stable values of T4 and T3 imply that the fall in reverse T3 was the result of enhanced catabolism induced by the β_2 -agonist rather than a decrease in conversion of reverse T3 from T4. This is in agreement with our previous work with laevodopa in semi-starvation states, where laevodopa prevented the usual rise in reverse T3 but had no effect on the fall in T3 despite the maintenance of the resting metabolic rate.¹ Thus it appears that conversion of T4 to T3 may not be under the direct control of β -receptor agonists but that these agonists may directly influence reverse T3 catabolism.

R T JUNG

Hammersmith Hospital,
London W12 0HS

N WALKER

Addenbrooke's Hospital,
Cambridge

Medical Research Centre,
Dunn Clinical Nutrition Centre,
Addenbrooke's Hospital,
Cambridge

¹ Shetty PS, Jung RT, James WPT. *Proc Nutrition Soc* 1978;38, 18 (abstract).

Nephrotic syndrome with slight proliferative changes: response to prednisone

SIR,—We read with great interest, the recent article by Dr H Rashid and others (2 August, p 347) on the response of patients with the nephrotic syndrome and slight proliferative changes in the glomeruli on light microscopy to prednisolone.

We have studied 16 adult patients who had the nephrotic syndrome and diffuse mesangial proliferative changes on light microscopy. Eleven were treated with prednisolone and five responded. The six who did not respond and five other patients were treated with a combination of cyclophosphamide, dipyrindamole, and warfarin. Of these, eight responded, including two patients who were non-responsive to prednisolone. Thus our overall response rate to both regimens was 81% (13/16). The mean duration of follow-up after remission was 12 months (range 2-43 months).

Treatment of the nephrotic syndrome with diffuse mesangial proliferative changes has shown variable results with different regimens in various studies. Cyclophosphamide and anticoagulants as single agents have been shown to be of use in several studies.¹⁻³ Our results so far indicate that a combination of these agents (cyclophosphamide, dipyrindamole, and warfarin) can be useful not only in patients not responsive to prednisolone but also as therapy of first choice.

Waldherr⁴ has suggested that diffuse mesangial proliferative glomerulonephritis with the nephrotic syndrome evolves from a minimal lesion with the same clinical syndrome. We were able to document this in one of our patients on clinical presentation and during a relapse of his nephrotic syndrome. He respon-

ded to prednisolone on both occasions. In our group of patients, seven (out of 11 who had immunofluorescent studies) had negative staining to all immunoglobulins. Six responded to treatment.

The implications of a diffuse mesangial proliferative glomerulonephritis on light microscopy in patients with the nephrotic syndrome is still unclear. Negative immunofluorescence may predict response to treatment, not only to steroids but also to cyclophosphamide, dipyrindamole, and warfarin. Further work and controlled trials are definitely needed in this area.

K T WOO

E J C LEE

C H LIM

Department of Renal Medicine,
Singapore General Hospital,
Singapore 0316

¹ German Glomerulonephritis Research Group. *Glomerulonephritis: international conference on pathogenesis, pathology and treatment*. 1977:196-201.

² Balslev JL, Brun C, Halberg F, et al. *Acta Med Scand* 1976;200:31-6.

³ Cade JR, de Quesada DL, Shires DM, et al. *Nephron* 1971;8:67-80.

⁴ Waldherr R, Gubler MC, Levy M, et al. *Clin Nephrol* 1978;10:171-9.

Factor VIII supply and demand

SIR,—Dr J D Cash's comments (23 August, p 565) on my letter (21 June, p 1532) require a brief response.

He states that I failed to mention the availability of plasma for fractionation as an important aspect of the problem of national self-sufficiency in blood products. There was a very good reason for this omission. The Wessex Regional Transfusion Service has recently offered to increase the supply of plasma to the Lister blood products laboratory, only to be told that there was not the capacity to handle the extra plasma.

I think it most important that the repetitive cry for increased plasma does not obscure the very real need for adequate fractionation potential in England today.

A ARONSTAM

Treloar Haemophilia Centre,
Lord Mayor Treloar Hospital,
Alton, Hants

Progesterone, fluid, and electrolytes in premenstrual syndrome

SIR,—Dr Katharina D Dalton (9 July, p 61) draws a distinction between women whose symptoms increase during the premenstrual week but which are none the less present throughout the cycle and women whose symptoms recur regularly during the same phase of each cycle and at no other phase. The latter she considers to be *true* premenstrual syndrome sufferers whereas the former are not. However, this distinction is even more difficult to make than might at first sight appear. For example, in the paper to which Dr Dalton refers¹ the diagnosis of the premenstrual syndrome was sufficiently elastic to include those women whose symptoms begin at or around ovulation, continue throughout the luteal phase, worsen premenstrually, and persist during the menstruum before fading away shortly after the period ceases. How is one to distinguish this "true" premenstrual tension sufferer from the woman whose diary reveals symptoms occurring intermittently during the cycle (but with seven or more days symptom free) with a premenstrual exacerbation?