distress: a double-blind trial of progesterone and placebo." Progesterone is no panacea for all the diagnoses which come under the umbrella of menstrual distress, including premenstrual exacerbations of symptoms present throughout the cycle, as well as spasmodic dysmenorrhoea (which responds to oestrogens and oral contraceptives) and endometriosis (which responds to progestogens).

Dr Clare also refers to the trial by Smith⁵; progesterone suppositories raise the plasma progesterone for 6-24 hours⁴ so medication is required at least once or twice daily and, on an adequate progesterone dosage, there is a cumulative effect during the first four days.⁴ Smith used a small dose of progesterone suppositories 200 mg on alternate days from day 19 until day 26; thus the progesterone was too late, too little, and too infrequent for this to be considered a trial. I would repeat that the common causes of failure with progesterone therapy are incorrect diagnosis and inadequate dosage.¹

K DALTON

London W1N 1AE

¹ Dalton K. *Premenstrual syndrome and progesterone therapy*. London: Heinemann Medical Books, 1977.

² Sampson G. *Br J Psychiat* 1979;135:208-15.

³ Dalton K. *Br J Psychiat* 1980;137:199.

⁴ Nillius SJ, Johansson EDB. *Am J Obstet Gynaec* 1971;110:470.

⁵ Smith SL. In Sachar EJ, ed. *Topics in psychoendocrinology*. New York: Grune and Stratton, 1975.

Gonorrhoea presenting as "sterile" pyuria

SIR,—Dr B Chattopadhyay and Mr I Hall draw attention in their paper (27 September, p 841) to the fact that gonorrhoea may occasionally present in men as "sterile" pyuria. In such circumstances urethral discharge is not noted by the patient (though careful examination after "milking" the anterior urethra will nearly always show evidence of purulent secretion).

The important clinical lesson to be drawn from their account is that symptoms of dysuria and frequency in young men should always be regarded as potentially serious conditions due either to sexually transmitted infections or to urinary tract infections secondary to some underlying abnormality. Among the common sexually transmitted infections, non-specific urethritis is often responsible for frequency and gonorrhoea for dysuria.

I would have thought that gonococcal infection in the female would be an extremely rare cause of "sterile" pyuria if a properly collected clean-catch specimen is sent to the laboratory. The short female urethra is usually adequately "rinsed" free of inflammatory cells by micturition with this technique.

J K OATES

Westminster Hospital,
London SW1P 2AP

SIR,—Dr B Chattopadhyay and Mr I Hall (27 September, p 841) are correct to add gonorrhoea to their list of causes of sterile pyuria in women, but the reason may be perverse. We were recently aided in making the diagnosis of gonorrhoea in a woman who presented complaining of haematuria by the finding of "sterile" pyuria in a midstream urine.

Two urine specimens received two days apart, and containing more than 20 white cells per high-

power field of a centrifuged deposit, showed no growth when cultured on CLED selective medium. Further incubation on non-selective blood agar in the presence of carbon dioxide failed to reveal the presence of significant numbers of fastidious organisms,¹ although no attempt was made to identify *Neisseria gonorrhoea*. Gram staining of the deposits revealed a mixed flora consistent with vaginal contamination. On being further questioned, the patient admitted to the recent onset of a vaginal discharge and *N gonorrhoea* was isolated from a cervical swab.

Women with vaginal discharges may frequently contaminate a midstream urine and the use of a vaginal pack inserted before collection has been recommended (Association of Clinical Pathologists Broadsheet 80). The finding of "sterile" pyuria in women may suggest a vaginal origin for these cells, and thus lead to a diagnosis of gonorrhoea.

M C KELSEY

Department of Microbiology

B DILLON

Department of Obstetrics and Gynaecology,
Barnet General Hospital,
Barnet, Herts EN5 3DJ

¹ Maskell R, Pead L, Allen J. *Lancet* 1979;i:1058-9.

Systematic review of the benzodiazepines

SIR,—The recommendations presented by the Committee on the Review of Medicines in its article "Systematic review of the benzodiazepines" (29 March, p 910) appropriately focuses concern about the use of these drugs. However, I would like to comment on one point that perhaps should be expanded on.

In the opening paragraphs the committee refers to the differences between the "long"-acting and the "short"-acting benzodiazepines, yet in the guidelines for lorazepam, oxazepam, etc, it mentions that paradoxical aggressive outbursts and uncovering of suicidal tendencies have been reported with the benzodiazepines. It should be noted that these reactions have been reported only in relation to the long-acting drugs—namely, diazepam,^{1,2} chlordiazepoxide,^{1,2} clorazepate,³ and flurazepam⁴ but not with oxazepam.^{2,5} In fact, oxazepam reduced anxiety scores in such patients and in this respect may be the drug of choice for the susceptible population. Furthermore, the uncovering of suicidal tendencies is probably a reference to the report by Hall and Joffe,⁶ which implicates only diazepam.

As the opinion of the committee will no doubt exert considerable influence on the labelling and use of benzodiazepines, I feel it would be appropriate to clarify this seeming inconsistency.

T R EINARSON

Pharmacy Department,
General and Marine Hospital,
Owen Sound, Ontario,
Canada N4K 5H3

¹ Anonymous. *Mass Gen Hosp Newsletter* 1978;1:38.

² Gardos G, DiMascio A, Salzman C, Shader RI. *Arch Gen Psychiatry* 1968;18:757-60.

³ Karch F. *Ann Int Med* 1979;91:61-2.

⁴ Einarson TR. *Drug Intell Clin Pharm* 1980;14:291-2.

⁵ Kochansky GE, Salzman C, Shader RI, Harmatz JS, Ogeltree AM. *Am J Psychiatry* 1975;132:861-3.

⁶ Hall RCW, Joffe JR. *Am J Psychiatry* 1972;129:738-42.

Heart rate variations in the Guillain-Barré syndrome

SIR,—We have read with interest the paper by Dr J C Frison and others (6 September,

p 649) reporting a reduction in cardiac beat-to-beat interval variation among patients with Guillain-Barré syndrome, which was taken to indicate a disorder of the autonomic innervation of the heart. We would like to point out that beat-to-beat variation, recorded as the standard deviation, is strongly dependent on both age and size of the mean R-R interval, so that these factors must be considered when comparisons are made between patients and normal groups.

Our measurements of beat-to-beat variation in normal subjects (to be published) show that the standard deviation declines with age and increases with the size of the mean R-R interval. Thus with a mean interval of 600 ms (or 100 beats/min) the standard deviation will average 45 ms at age 7 years but 24 ms at 50 years. Similarly, at age 33 years the predicted standard deviation will be 31 ms at a mean interval of 600 ms and 66 ms at a mean interval of 1000 ms (or 60 beats/min).

Thus although the two groups studied by Dr Frison and his colleagues were matched for age the difference in the size of the standard deviation between the two could be explained, at least in part, by a tachycardia in the Guillain-Barré patients. Since tachycardia is a common feature of the acute phase of the illness, its resolution during recovery should necessarily be accompanied by an increase in the size of the standard deviation.

In conclusion, we do not believe that the reduction in beat-to-beat variation noted by these authors can properly be ascribed to autonomic neuropathy in the absence of a consideration of resting heart rate.

S A SMITH

S E SMITH

Department of Pharmacology,
St Thomas's Hospital Medical
School,
London SE1 7EH

No to euthanasia

SIR,—Having read the statement on euthanasia (6 September, p 691), may I voice the opinion of a geriatric nearing 80 who has had a lengthy experience of hospitalisation in different medical and surgical wards (not strictly geriatric) for painful afflictions, which at times in my weakened state made me feel that I would be better off dead? Despite the almost unbearable pain I sometimes endured I am strongly opposed to the belief, expressed by EXIT in a recent broadcast on Radio 4, that euthanasia should be officially legalised. I find it difficult to appreciate the benefits (and what really are they?) which are claimed.

So far as those who are younger and normally able are concerned, it can seldom be needed, as most can—and in fact do—voluntarily provide their own euthanasia by overdoses of drugs, etc. With regard to the elderly, whether or not the illness is terminal—and indeed how can any doctor be quite sure?—there exists an emotional ebb and flow, with depression deepening more at some phases of the illness than at others. At such low levels the patient is most suggestible, and may quite well give a misconceived consent to euthanasia, whereas in a better phase he might have decided more reasonably otherwise. During this highly suggestible time there is some risk of exposure to the insidious persuasion of anyone who might have personal reasons for not wishing him or her to live longer. Under such circumstances distressed close relatives might reluctantly agree to give their consent, only to