more distress than single-site aspirations so it was routine practice to cover the procedure by intravenous diazepam. I found this to be extremely successful and in subsequent patients who are anxious I have continued with the practice of sedation by this method.

ROBERT LEONARD

University Department of
Medicine,
Royal Victoria Infirmary,
Newcastle upon Tyne NE1 4LP

Urethral catheterisation

SIR,—The article on urethral catheterisation (4 October, p 928) justifies two comments.

Firstly, although the article gave an excellent account of the reasons for catheterisation and the types of catheters to be used and placed a great deal of emphasis on absolute asepsis, it failed to give an adequate description of the actual catheterisation technique. Urethral catheterisation is a very common procedure on medical and surgical wards and this task is usually delegated to the most junior member of a team. It would, therefore, have been more helpful if the steps in the procedure had been demonstrated by a series of diagrams (as is commonly done with urethral dilatation).

Secondly, the authors failed to commit themselves about the amount of sterile water required to inflate a catheter balloon and recommended "an appropriate volume." Most manufacturers recommend a volume in excess of 20 ml, but many practising urologists regard this as an uncomfortable and unnecessary amount and usually use volumes of less than 10 ml.

K HASHEMI

Department of Surgery,
East Birmingham Hospital,
Birmingham B9 5ST

Adverse reaction to bupivacaine

SIR,—Dr A M Henderson's case (18 October, p 1043) of an adverse reaction to bupivacaine used for intravenous regional anaesthesia is a timely reminder that bupivacaine is *not* the drug of choice for this technique. He terms it "unexpected" but we have consistently argued against the use of bupivacaine^{1,2} for this procedure because of the risk of such a reaction. The potential for systemic toxicity is great when the tourniquet is released as a relatively large dose of local anaesthetic rapidly enters the circulation. Therefore the drug of choice is prilocaine³ since it is by far the least toxic of the available local anaesthetics. The incidence of systemic toxicity using bupivacaine is low but with prilocaine it is non-existent.

Plasma protein binding does not confer any safety margin to a drug like bupivacaine, which crosses membranes easily and rapidly. Further, the plasma concentrations mentioned by Dr Henderson are peripheral venous in origin. Of more relevance to the occurrence of systemic toxicity immediately following tourniquet release are the arterial concentrations. The situation is equivalent to that following a bolus injection, when transient high arterial concentrations (and therefore brain concentrations) are not reflected in venous measurements.

We would also suggest that systemic reactions to local anaesthetics are far more effectively treated with increments (50 mg) of intravenous thiopentone than with a combination of diazepam and chlorpromazine. The

reaction would have been controlled much faster and we doubt very much if the patient would have had to be kept in hospital overnight.

J A W WILDSMITH
D H T SCOTT
D B SCOTT

Department of Anaesthetics,
Royal Infirmary,
Edinburgh EH3 9YW

¹ Brown DT, Scott DHT, Wildsmith JAW. *Hospital Update*, 1978; November: 738.

² Wildsmith JAW, Scott DHT, Brown DT. *Anaesthesia* 1979;34:919.

³ Eriksson E, ed. *Illustrated handbook in regional anaesthesia*. Munksgaard: Copenhagen, 1969.

Cancer and the fetus

SIR,—Minerva refers (9 August, p 461) to a review article¹ which pointed out the rarity of cancer occurring during pregnancy and to the amazingly rare transmission of malignant cells to the fetus across the placenta. I personally reviewed the world literature in 1967² and at the time I could find only 22 documented cases recording placental or even transplacental metastases, and I reported only the third case involving carcinoma of the bronchus. Most cases were of malignant melanoma and it is these tumours which appear to metastasise more readily to the placenta and fetus.

In my original case cells morphologically identical to those of the primary oat cell carcinoma were found by my colleague Dr R Seal in the maternal blood spaces of the placenta. The fetus was spared and at follow-up two years later the child was quite normal.

It would be interesting to know why the trophoblast appears to resist tumour cell invasion. Surely this must be an important research area in our entire understanding of the biology of carcinoma in humans.

EGRYN M JONES

Heald Green, Cheshire

¹ Greene M. *JAMA* 1980;243:2241.

² Jones EM. *Br Med J* 1969;iii:491-2.

Controlled trial of cromoglycate and slow-release aminophylline in childhood asthma

SIR,—The paper by Dr A T Edmunds and others (27 September, p 842) raises some questions of ethics. The authors state that both theophylline and cromoglycate provide effective prophylactic treatment in childhood asthma. Why then was it necessary to include a placebo period? These quite severely affected children were on no regular medication for four weeks, during which time their symptoms increased and on eight occasions hospital admission and attendant loss of schooling resulted. The question of venesection at the end of two treatment periods when they were obviously not taking aminophylline is also open to question.

It has been my experience that asthmatic children who are well controlled on regular therapy are reluctant to enter a study of this type. It is therefore surprising that not one of 30 children dropped out of the study during an exacerbation of symptoms when they knew that there was a possibility that they were taking a placebo only. It would be interesting to know exactly what the parents were told before consent was obtained for their children's participation in this study.

The conclusion of this study supports the work of MacDonald and McWilliam, who found cromoglycate and slow-release aminophylline equally effective.¹ I would like to add that when the price of spinhalers and theophylline assay are included in the costing slow-release aminophylline continues to be the most cost-effective drug, which is preferred by the majority of children.

I BLUMENTHAL

St James's University Hospital,
Leeds LS9 7TF

¹ MacDonald TH, McWilliam R. *J Int Med Res* 1979;7, suppl 1:87-92.

*. *We sent this letter to Dr Edmunds, who replies below.—Ed, *BMJ*.

SIR,—We thank Dr Blumenthal for his comments and interest. It was necessary to include a placebo period to demonstrate that the trial patients required the treatment being studied. All our patients were able to take salbutamol freely throughout the trial. Patients in whom unacceptable symptoms persisted despite regular use of salbutamol, and patients who recorded peak expiratory flow rates less than 25% of the expected normal on all three measurements over a period of 24 hours, were reviewed at the hospital. If necessary they were treated with a short course of prednisolone. If a patient required more than two courses of prednisolone in any treatment period he or she would have moved on to the next treatment; however, the need for this did not arise.

The study was approved by the United Bristol Hospitals ethical committee. All patients and their parents were given full details of the trial before they gave consent. Thus they knew that they would not remain ill for long if they did develop acute asthma. They also knew that venepuncture was required at the end of each period as the trial was double blind, and it was necessary to demonstrate that therapeutic doses of slow-release aminophylline were being taken. The majority of parents at the conclusion of the trial felt that they understood the pattern of their children's asthma better, and that their children were subsequently better controlled when maintained on the best-choice treatment as found in the trial.

A T EDMUNDS

University Department of Child Health,
Royal Hospital for Sick Children,
Bristol BS2 8BJ

Family planning

SIR,—In his book review on the development of family planning services in Britain (11 October, p 999) Mr Michael Brudenell begs some questions. Firstly, he infers that family planning is about the "unplanned pregnancy problem" and, secondly, that "the consumer is the ultimate arbiter of what is right for him or her."

The first statement puts clinical family planning into a totally negative role, whereas in reality—when almost half of those using the services are nulliparous—the doctor is in an ideal situation to affect the health of early pregnancy for many of these *future* parents. Health advice on rubella vaccination, smoking in pregnancy, nutrition, post-pill contraception for up to six months before conception, etc, is acceptable to women in this situation.