whereas no increase was found for radiotherapy.

ANTHONY GREEN

Barnet,
Herts EN5 4QA

- ¹ Chabner B. *New Eng J Med* 1977;297:213-4.
² Wall RJ, Clausen KP. *New Eng J Med* 1975;293:271.
³ Reimer RR, et al. *New Eng J Med* 1977;297:171-81.

Cervical cancer screening

SIR,—We were interested to read the opinions expressed in the National Institutes of Health (NIH) consensus statement published in the *BMJ* (8 November, p 1264) concerning cervical cancer screening policy. There is clearly a degree of uncertainty about the appropriate screening interval ("... regular intervals of one to three years"), which parallels the American Cancer Society's change of policy from annual to three-yearly screening.¹

In this country there is debate on the need for regular screening of young women in the face of DHSS policy to reimburse general practitioners only for examinations of women aged 35 or more.^{2,3} The controversy has been fuelled by statistics showing rising mortality from cancer of the cervix in younger age groups (see accompanying table). In fact, since only a minority of smears are taken by general practitioners most examinations are on women under 35, and have been for many years.⁴

Mortality from Cancer of Cervix (per million) England & Wales

Year	Age group	
	15-34 y	Over 34 y
1970	6.0	170
1971	6.6	175
1972	7.6	167
1973	6.9	170
1974	8.2	155
1975	11.4	159
1976	10.1	165
1977	10.5	160
1978	13.3	159
1979	00.0	000

It should be obvious that, if screening is beneficial at all, the more frequently it is carried out the better the results (cancers prevented, life years saved). However, more intensive programmes involve increasing costs, and there are practical and ethical problems in setting up trials to evaluate the corresponding benefits which might result. Practical difficulties would include the need to compare the effects of a variety of screening regimens on different risk categories of women. Any less comprehensive trials, in which few beneficial effects of screening were detected, would result in calls to repeat the trial with a different pattern of screening or on another risk category of women.

We would support Dr Duncan Neuhauser's comments, in his "minority opinion," concerning the value of using computer-based policy models. These models may be constructed for testing a variety of screening policies on mathematical populations of women.^{5,6} Ethical difficulties are avoided, and practical problems are reduced to testing the assumptions made in the model about the natural history of the disease and the effects of screening. We have recently constructed our own interactive model, in which a population of women defined by age, parity, and marital status are allowed mathematically to

be born, age, give birth, be screened, develop cervical cancer, and die of that disease or from other causes. The influence of screening policies can be simulated; and medical, human, and financial indices may be examined.

Validation of such models is inherently difficult. Existing survey data and consensus statements of the type made by NIH are of value to model builders in providing data against which to validate. Confidence in the predictions of these models comes only with sensitivity analysis. Such models do, however, allow students and planners to experiment with alternative policies and get a feel for the dynamics of the situation, which is simply not possible in the analysis phase of a clinical trial.

The impossibility of carrying out correctly planned clinical trials for screening may not be the great disadvantage one might at first consider it.

D M PARKIN

Leeds Area Health Authority
(Teaching),

A D CLAYDEN

Medical Statistics Section,
University Department of
Community Medicine,
Leeds

- ¹ American Cancer Society. CA-A cancer journal for clinicians 1980;30:215-23.
² Anonymous. *Lancet* 1978;ii:1029-30.
³ Cotton R. *World Medicine* 1980;15:21-6.
⁴ Department of Health and Social Security. Health and personal social services statistics for England. London: HMSO, 1980.
⁵ Knox EG. In: McLachlan G, ed. *Problems and progress in medical care*. Vol 9. London: Oxford University Press, 1973.
⁶ Coppleson LW, Brown B. *Am J Obstet Gynec* 1975;122:127-36.

Chasing the unknown primary

SIR,—With reference to your leading article (8 November, p 1232) on adenocarcinoma from unknown primary sites, several points of clinicopathological importance deserve emphasis.

As you state, the diagnosis of adenocarcinoma must be accurately established and other possible types of tumour excluded. Although not mentioned in your editorial, in difficult cases transmission electron microscopy is a desirable and frequently helpful investigation. In this department electron microscopy of suspected adenocarcinoma has revealed tumours such as poorly differentiated squamous carcinoma, melanoma, apudoma (including oat cell carcinoma), lymphoma, mesothelioma, and malignant histiocytoma. The diagnosis of adenocarcinoma can itself be confirmed by finding one or more characteristic ultrastructural features in the tumour cells.¹

The importance of establishing the correct diagnosis is so vital that a strong case can be made for establishing NHS centres with a diagnostic electron microscopy service. In Sheffield, we currently offer a "free open-access" service to surrounding district general hospitals. In most cases an ultrastructural answer is available within 48 hours. A fact not appreciated by many clinicians (and histopathologists) is that the information is frequently obtained from material that has been fixed in formalin or embedded in paraffin wax. Although preservation is not optimal, meaningful answers can be obtained in most instances.

"Special stains" are briefly mentioned in the leading article but their potential importance is not explained. The diagnosis of adenocarcinoma by light microscopy is often dependent on the demonstration of mucin

within the tumour cells. Different primary tumours produce different types of mucin and the mucins can be distinguished by relatively simple histochemical techniques.² With this technique, cases of "bronchial" adenocarcinoma have been shown to be secondaries from the large bowel.

The gall bladder, bile duct, and thyroid can be all sites of occult adenocarcinoma but they do not appear in your lists of possible primary sites. Systemic treatment is also said to be warranted for "carcinoma of the testis." Such terminology is presumably derived from the American references quoted. In Britain a different classification of testicular tumours is used³ and true adenocarcinoma of the testis is a rarity. To help those who are initiating therapy it should be made clear that you are referring to British subtypes of teratoma.

In the future, most hope for solving a case of adenocarcinoma from an unknown primary site must come from immunoperoxidase techniques. A secretory protein hormone/enzyme profile will hopefully pinpoint the primary tumour.

DAVID SLATER

Department of Histopathology,
Royal Hallamshire Hospital,
Sheffield S10 2JF

- ¹ Ghadially FN. *Diagnostic electron microscopy of tumours*. London: Butterworths, 1980:68-77.
² Culling CFA, Reid PE, Burton JD, Dunn WL. *J Clin Path* 1975;28:656-8.
³ Pugh RCB. *Pathology of the testis*. Oxford: Blackwell Scientific Publications, 1976:139-59.

Pneumonia during treatment of acute leukaemia

SIR,—Your recent leading article "Pneumonia during the treatment of acute leukaemia," (8 November, p 1235) provides an interesting, if selective, summary of experience in certain centres in the USA, but has little relevance to the infective problems encountered in leukaemia units in Britain, and may indeed be misleading if interpreted as a guide to practice here. The most common infections in patients with neutropenia, the proportion with positive microbiology, the percentage with proved bacteraemia, and the nature of the causative organisms vary sharply both from time to time and from centre to centre in Britain and internationally. The early management of infections requires a knowledge of the current pathogenic flora locally in the wards concerned rather than in Baltimore or Houston.

Our experience in Cambridge suggests that while pneumonias are not infrequent they are neither the most common nor the most important infections in patients with neutropenia and leukaemia. Bacteraemias of uncertain origin, possibly most often arising from the gastrointestinal tract, are more common and more serious causes of the dangerous state of circulatory collapse associated with infection by Gram-negative organisms, which you refer to—with somewhat dubious accuracy—as endotoxaemia. The combination of tobramycin and a cephalosporin is not, in fact, established to be more effective than a number of other possible combinations—for example, gentamycin and carbenicillin. Indeed, the EORTC trial report¹ given as authority for the tobramycin with cephalosporin recommendation did not use this combination of drugs but did show that of 443 infected sites in 396 patients only 56 were pulmonary, and that of 159 bacteraemias